

## 6. REFERENCES

- Brown, R.M., Jr. (2004). Cellulose Structure and Biosynthesis: what is on the store for the 21st Century? *Journal of Polymer Science: Part A: Polymer Chemistry* 42(3): 487-495.
- Budhiono, A., Rosidi, B., Taher, H. and M. Iguchi (1999). Kinetic aspects of bacterial cellulose formation in nata-de-coco culture system. *Carbohydr Polymers* 40:137-143.
- Cannon, R.E. and Anderson, S.M. (1991). Biogenesis of Bacterial Cellulose. *Critical Reviews in Microbiology* 17(6): 435-447.
- Carlson, R. (1992). *Design and Optimization in the Organic Synthesis*. Elsevier Science B.V. The Netherlands.
- De Gallardo, D.E.J., Andres, R.M. and E.T. Magno (1971). A study on the isolation and screening of microorganisms for production of diverse-textured nata. *Philipp J Sci* 100:41-49
- Halib, N, Amin. M and I. Ahmad (2012). Physicochemical Properties and Characterization of *Nata de coco* from Local Food Industries as a Source of Cellulose. *Sains Malaysiana* 41(2): 205-211
- Hestrin, S. and Schramm, M. (1954). Synthesis of cellulose by *Acetobacter xylinum*: preparation of freeze-dried cells capable of polymer-izing glucose to cellulose. *Biochem J* 58:345-352
- Jonas, R. and Farah, L.F. (1998). Production and application of microbial cellulose. *Polymer Degradation and Stability* 59: 101-106
- Keshk, S and Sameshima, K. (2006). The utilization of sugar cane molasses with/without the presence of lignosulfonate for the production of bacterial cellulose. *Appl Microbiol Biotechnol* 72(2):291-296.
- Lapuz, M.M., Gallardo, E.G. and M.A. Palo (1967). The nata organism cultural requirements, characteristics and identity. *Philipp J Sci* 96(2):91-108
- Montgomery, D.C. *Response Surface Methods and Other Approaches to Process Optimization*. In: D.C. Montgomery (1997). *Design and Analysis of Experiments* (Ed.). John Wiley & Sons, New York, USA pp. 427-510
- Pambayun, R. (2002). *Teknologi Pengolahan Nata de coco*. Kanisius. Yogyakarta.

Parajó, J.C., Santos, V., Domínguez, H., Vázquez, M. (1995). NH<sub>4</sub>OH--based pretreatment for improving the nutritional quality of single-cell protein (SCP). *Appl. Biochem. Biotechnol.* 55:133–149.

Rahman, A. (1992). *Teknologi Fermentasi*. ARCAN Pusat Antar Universitas Pangan dan Gizi IPB. Bandung.

Sanchez, P.C. (2008). *Philippine Fermented Foods*. The University of The Philippines Press. Quezon City, The Philippines.

Sutarminingsih, L. (2004). *Peluang Usaha Nata de coco*. Kanisius. Yogyakarta

Vandamme, E.J., de Baets, S., Vanbaelen, A., Joris, K., and P. de Wulf. (1998) Improved production of bacterial cellulose and its application potential. *Polym Degrad Stab* 59:93–99.

Verschuren, P.G., Carodona, T.D., Nout, M.J.R., de Gooijer, K.D., van den Heuvel, J.C. (2000) Location and limitation of cellulose production by *Acetobacter xylinum* established from oxygen profiles. *J Biosci Bioeng* 89(5):414–419.

Verseput, R. (2000). "Digging into DOE: Selecting the right central composite design for response surface methodology applications." <http://www.qualitydigest.com/june01/html/doe.html>

White, D.G. and Brown, R.M. Jr. (1989). Prospects for the commercialization of the biosynthesis of microbial cellulose. In: Schuerech, C (ed). *Cellulose and wood-chemistry and technology*. Wiley, New York.

Yamanaka, S, Watanabe, K and N. Kitamura (1989). The structure and mechanical properties of sheets prepared from bacterial cellulose. *J Mater Sci* 24:3141–3145.

Zakaria, J. and Nazeri, M.A. (2012). Optimization of Bacterial Cellulose Production from Pineapple Waste: Effect of Temperature, pH and Concentration. *EnCon 2012, 5th Engineering Conference, "Engineering Towards Change - Empowering Green Solutions"*