

6. DAFTAR PUSTAKA

- Aboubakar, Njintang, Y. N., Scher, J., & Mbofung, C. M. F. (2008). Physicochemical, thermal properties and microstructure of six varieties of taro (*Colocasia esculenta* L. Schott) flours and starches. *Journal of Food Engineering*, 86(2), 294–305. <https://doi.org/10.1016/j.jfoodeng.2007.10.006>
- Adeyeye, S. A. O. (2016). Assessment of quality and sensory properties of sorghum – wheat flour cookies. *Cogent Food & Agriculture*, 2(1), 1–10. <https://doi.org/10.1080/23311932.2016.1245059>
- Ahmed, J., Thomas, L., & Arfat, Y. A. (2019). Functional, rheological, microstructural and antioxidant properties of quinoa flour in dispersions as influenced by particle size. *Food Research International*, 116, 302–311. <https://doi.org/10.1016/j.foodres.2018.08.039>
- Altindag, G., Certel, M., Erem, F., & Ilknur Konak, Ü. (2015). Quality characteristics of gluten-free cookies made of buckwheat, corn, and rice flour with/without transglutaminase. *Food Science and Technology International*, 21(3), 213–220. <https://doi.org/10.1177/1082013214525428>
- Amante, P. R., Santos, E. C. Z., Correia, V. T. da V., & Fante, C. A. (2020). Research Notes: Benefits and Possible Food Applications of Arrowroot (*Maranta Arundinaceae* L.). *Journal of Culinary Science and Technology*, 00(00), 1–9. <https://doi.org/10.1080/15428052.2020.1791295>
- Amin, T., Naik, H. R., Hussain, S. Z., Mir, M. A., & Jabeen, A. (2018). In vitro digestion , physicochemical and morphological properties of low glycemic index rice flour prepared through enzymatic hydrolysis. *International Journal of Food Properties*, 21(1), 2632–2645. <https://doi.org/10.1080/10942912.2018.1545789>
- Aprianita, A., Vasiljevic, T., Bannikova, A., & Kasapis, S. (2014). Physicochemical properties of flours and starches derived from traditional Indonesian tubers and roots. *Journal of Food Science and Technology*, 51(12), 3669–3679. <https://doi.org/10.1007/s13197-012-0915-5>
- Arun, K. B., Persia, F., & Aswathy, P. S. (2015). Plantain peel - a potential source of antioxidant dietary fibre for developing functional cookies. *Journal of Food Science and Technology*, 52(10), 6355–6364. <https://doi.org/10.1007/s13197-015-1727-1>
- Asmeda, R., Noorlaila, A., & Norziah, M. H. (2015). Relationships of damaged starch granules and particle size distribution with pasting and thermal profiles of milled MR263 rice flour. *FOOD CHEMISTRY*, 191, 45–51. <https://doi.org/10.1016/j.foodchem.2015.05.095>
- Belorio, M., Sahagún, M., & Gómez, M. (2019). Influence of flour particle size distribution on the quality of maize gluten-free cookies. *Foods*, 8(2).

<https://doi.org/10.3390/foods8020083>

- Bilgiçli, N., & İbanoğlu, Ş. (2015). Effect of pseudo cereal flours on some physical, chemical and sensory properties of bread. *Journal of Food Science and Technology*, 52(11), 7525–7529. <https://doi.org/10.1007/s13197-015-1770-y>
- Bilgiçli, N., Kara, M., Elgün, A., Ertaş, N., & Demir, M. K. (2006). Determination of technologic and sensory properties of cookies prepared with corn flour. *Journal of Food, Agriculture and Environment*, 4(2), 109–111. https://www.researchgate.net/profile/Meryem-Kara-5/publication/235799904_Determination_of_technologic_and_sensory_properties_of_cookies_prepared_with_corn_flour/links/577e7ac808aed807ae7b1ade/Determination-of-technologic-and-sensory-properties-of-cookies-prepared-with-corn-flour.pdf
- Brito, I. L., de Souza, E. L., Felex, S. S. S., Madruga, M. S., Yamashita, F., & Magnani, M. (2014). Nutritional and sensory characteristics of gluten-free quinoa (*Chenopodium quinoa Willd*)-based cookies development using an experimental mixture design. *Journal of Food Science and Technology*, 52(9), 5866–5873. <https://doi.org/10.1007/s13197-014-1659-1>
- BSN. (2011). *Biskuit* (SNI 2973:2). Badan Standarisasi Nasional.
- Cannas, M., Pulina, S., Conte, P., Caro, A. Del, Urgeghe, P. P., Piga, A., & Fadda, C. (2020). Effect of Substitution of Rice Flour with Quinoa Flour on the Chemical-Physical, Nutritional, Volatile and Sensory Parameters of Gluten-Free Ladyfinger Biscuits. *Foods*, 9(6), 808. <https://doi.org/doi:10.3390/foods9060808>
- Caprez, A., Arrigoni, E. V. A., Amado, R., & Neukom, H. (1986). Influence of Different Types of Thermal Treatment on the Chemical Composition and Physical Properties of Wheat Bran. *Journal of Cereal Science*, 4(3), 233–239. [https://doi.org/10.1016/S0733-5210\(86\)80025-X](https://doi.org/10.1016/S0733-5210(86)80025-X)
- Chauhan, A., Saxena, D. C., & Singh, S. (2015). Total dietary fibre and antioxidant activity of gluten free cookies made from raw and germinated amaranth (*Amaranthus spp.*) flour. *LWT - Food Science and Technology*, 63(2), 939–945. <https://doi.org/10.1016/j.lwt.2015.03.115>
- Chevallier, S., Colonna, P., & Bule, A. (2000). Physicochemical Behaviors of Sugars , Lipids , and Gluten in Short Dough and Biscuit. *Journal of Agricultural and Food Chemistry*, 48, 1322–1326. <https://doi.org/10.1021/jf990435+>
- Chung, H., Cho, A., & Lim, S. (2014). Utilization of germinated and heat-moisture treated brown rices in sugar-snap cookies. *LWT - Food Science and Technology*, 57(1), 260–266. <https://doi.org/10.1016/j.lwt.2014.01.018>
- Ciacci, C., Maiuri, L., Caporaso, N., Bucci, C., Del Giudice, L., Rita Massardo, D., Pontieri, P., Di Fonzo, N., Bean, S. R., Ioerger, B., & Londei, M. (2007). Celiac

- disease: In vitro and in vivo safety and palatability of wheat-free sorghum food products. *Clinical Nutrition*, 26(6), 799–805. <https://doi.org/10.1016/j.clnu.2007.05.006>
- Darman, R. D., Sidoine, M. B., & Lenzemo, V. W. (2020). Development of sorghum-based shortbread biscuits from “muskwari” flour. *Food Science and Nutrition*, 8(7), 3181–3189. <https://doi.org/10.1002/fsn3.1574>
- Doescher, L. C., Hosene, R. C., Milliken, G. A., & Rubenthaler, G. L. (1987). Effect of Sugars and Flours on Cookies Spread Evaluated by Time-Lapse Photography. *Cereal Chemistry*, 64(3), 163–167. https://www.cerealsgrains.org/publications/cc/backissues/1987/Documents/64_163.pdf
- Donfrancesco, B. Di, & Koppel, K. (2017). Sensory Characteristics and Volatile Components of Dry Dog Foods Manufactured with Sorghum Fractions. *Molecules*, 22(6). <https://doi.org/10.3390/molecules22061012>
- Dovi, K. A. P., Constance Chiremba, & Taylor, J. R. N. & H. L. de K. (2018). Rapid sensory profiling and hedonic rating of whole grain sorghum-cowpea composite biscuits by low income consumers. *Journal of the Science of Food and Agriculture*, 98(3), 905–913. <https://doi.org/10.1002/jsfa.8536>
- Fustier, P., Castaigne, F., Turgeon, S. L., & Biliaderis, C. G. (2008). Flour constituent interactions and their influence on dough rheology and quality of semi-sweet biscuits: A mixture design approach with reconstituted blends of gluten, water-solubles and starch fractions. *Journal of Cereal Science*, 48(1), 144–158. <https://doi.org/10.1016/j.jcs.2007.08.015>
- Garzón, A. G., Erben, M., Osella, C. A., & Drago, S. R. (2020). Effects of baking on γ -aminobutyric acid and free phenolic acids from gluten-free cookies made with native and malted whole sorghum flours. *Journal of Food Processing and Preservation*, 44(8), 1–10. <https://doi.org/10.1111/jfpp.14571>
- Giarnetti, M., Paradiso, V. M., Caponio, F., Summo, C., & Pasqualone, A. (2015). Fat replacement in shortbread cookies using an emulsion filled gel based on inulin and extra virgin olive oil. *LWT - Food Science and Technology*, 63(1), 339–345. <https://doi.org/10.1016/j.lwt.2015.03.063>
- Giri, N. A., & Sajeev, M. S. (2020). Physico-Mechanical and Nutritional Evaluation of Taro (*Colocasia esculenta*) Flour-based Gluten-free Cookies. *Agricultural Research*, 9(1), 125–131. <https://doi.org/10.1007/s40003-019-00411-z>
- Giuberti, G., Rocchetti, G., Sigolo, S., Fortunati, P., Lucini, L., & Gallo, A. (2018). Exploitation of alfalfa seed (*Medicago sativa* L.) flour into gluten-free rice cookies: Nutritional, antioxidant and quality characteristics. *Food Chemistry*, 239, 679–687. <https://doi.org/10.1016/j.foodchem.2017.07.004>

- Gordillo, C. A. S., Valencia, G. A., Zapata, R. A. V., & Henao, A. C. A. (2014). Physicochemical Characterization of Arrowroot Starch (*Maranta arundinacea* Linn) and Glycerol / Arrowroot Starch Membranes. *International Journal of Food Engineering*, 10(4), 727–735. <https://doi.org/10.1515/ijfe-2014-0122>
- Goyat, J., Passi, S. J., Suri, S., & Dutta, H. (2018). Development of Chia (*Salvia hispanica*, L.) and Quinoa (*Chenopodium quinoa*, L.) seed flour substituted cookies-physicochemical, nutritional and storage studies. *Current Research in Nutrition and Food Science*, 6(3), 757–769. <https://doi.org/10.12944/CRNFSJ.6.3.18>
- Hazelton, J. L., DesRochers, J. L., & Walker, C. E. (2003). BISCUITS, COOKIES, AND CRACKERS | Chemistry of Biscuit Making. In *Encyclopedia of Food Sciences and Nutrition* (second ed, pp. 533–539). Elsevier. <https://doi.org/10.1016/b0-12-227055-x/00105-x>
- Indrastati, N., & Anjani, G. (2016). Snack Bar Kacang Merah Dan Tepung Umbi Garut Sebagai Alternatif Makanan Selingan Dengan Indeks Glikemik Rendah. *Journal of Nutrition College*, 5(4), 546–554. <https://ejournal3.undip.ac.id/index.php/jnc/article/view/16471/15870>
- Jafari, M., Koocheki, A., & Milani, E. (2017). Effect of extrusion cooking on chemical structure, morphology, crystallinity and thermal properties of sorghum flour extrudates. *Journal of Cereal Science*, 75, 324–331. <https://doi.org/10.1016/j.jcs.2017.05.005>
- Jan, K. N., Panesar, P. S., & Singh, S. (2017). Textural, in vitro antioxidant activity and sensory characteristics of cookies made from blends of wheat-quinoa grown in India. *Journal of Food Processing and Preservation*, 42(3). <https://doi.org/10.1111/jfpp.13542>
- Jan, K. N., Panesar, P. S., & Singh, S. (2018). Optimization of antioxidant activity, textural and sensory characteristics of gluten-free cookies made from whole indian quinoa flour. *Lwt*, 93, 573–582. <https://doi.org/10.1016/j.lwt.2018.04.013>
- Jan, R., Saxena, D. C., & Singh, S. (2016a). Effect of Germination on Nutritional, Functional, Pasting and Microstructural Properties of Chenopodium (*Chenopodium album*) Flour. *Journal of Food Processing and Preservation*, 41(3), 1–10. <https://doi.org/10.1111/jfpp.12959>
- Jan, R., Saxena, D. C., & Singh, S. (2016b). Physico-chemical, textural, sensory and antioxidant characteristics of gluten - free cookies made from raw and germinated Chenopodium (*Chenopodium album*) flour. *LWT - Food Science and Technology*, 71, 281–287. <https://doi.org/10.1016/j.lwt.2016.04.001>
- Jan, R., Saxena, D. C., & Singh, S. (2018). Comparative study of raw and germinated Chenopodium (*Chenopodium album*) flour on the basis of thermal, rheological, minerals, fatty acid profile and phytochemicals. *Food Chemistry*, 269, 173–180.

<https://doi.org/10.1016/j.foodchem.2018.07.003>

- Jan, U., Gani, A., Ahmad, M., Shah, U., Baba, W. N., Masoodi, F. A., Maqsood, S., Gani, A., Wani, I. A., & Wani, S. M. (2015). Characterization of cookies made from wheat flour blended with buckwheat flour and effect on antioxidant properties. *Journal of Food Science and Technology*, 52(10), 6334–6344. <https://doi.org/10.1007/s13197-015-1773-8>
- Julianti, E., Rusmarilin, H., & Yusraini, E. (2017). Functional and rheological properties of composite flour from sweet potato, maize, soybean and xanthan gum. *Journal of the Saudi Society of Agricultural Sciences*, 16(2), 171–177. <https://doi.org/10.1016/j.jssas.2015.05.005>
- Kaur, M., Kaushal, P., & Sandhu, K. S. (2011). Studies on physicochemical and pasting properties of Taro (*Colocasia esculenta* L.) flour in comparison with a cereal, tuber and legume flour. *Journal of Food Science and Technology*, 50(1), 94–100. <https://doi.org/10.1007/s13197-010-0227-6>
- Kaushal, P., Kumar, V., & Sharma, H. K. (2012). Comparative study of physicochemical, functional, antinutritional and pasting properties of taro (*Colocasia esculenta*), rice (*Oryza sativa*) flour, pigeonpea (*Cajanus cajan*) flour and their blends. *LWT - Food Science and Technology*, 48(1), 59–68. <https://doi.org/10.1016/j.lwt.2012.02.028>
- Kementerian Pertanian Republik Indonesia. (2018). *Statistik Konsumsi Pangan Tahun 2018*. http://epublikasi.setjen.pertanian.go.id/epublikasi/StatistikPertanian/2018/Konsumsi/Statistik_Konsumsi_Pangan_Tahun_2018/files/assets/basic-html/page124.html
- Khan, M. A., Mahesh, C., Semwal, A. D., & Sharma, G. K. (2013). Effect of spinach powder on physico-chemical, rheological, nutritional and sensory characteristics of chapati premixes. *Journal of Food Science and Technology*, 52(4), 2359–2365. <https://doi.org/10.1007/s13197-013-1198-1>
- Korus, A., Gumul, D., Krystyan, M., Juszczak, L., & Korus, J. (2017). Evaluation of the quality, nutritional value and antioxidant activity of gluten-free biscuits made from corn-acorn flour or corn-hemp flour composites. *European Food Research and Technology*, 243(8), 1429–1438. <https://doi.org/10.1007/s00217-017-2853-y>
- Krystyan, M., Gumul, D., Korus, A., Korus, J., & Sikora, M. (2018). Physicochemical properties and sensory acceptance of biscuits fortified with *Plantago psyllium* flour. *Emirates Journal of Food and Agriculture*, 30(9), 758–763. <https://doi.org/10.9755/ejfa.2018.v30.i9.1799>
- Kumar, K. A., Sharma, G. K., Khan, M. A., Govindaraj, T., & Semwal, A. D. (2015). Development of multigrain premixes—its effect on rheological, textural and microstructural characteristics of dough and quality of biscuits. *Journal of Food Science and Technology*, 52(12), 7759–7770. <https://doi.org/10.1007/s13197-015-1950-9>

- Laguna, L., Salvador, A., Sanz, T., & Fiszman, S. M. (2011). Performance of a resistant starch rich ingredient in the baking and eating quality of short-dough biscuits. *LWT - Food Science and Technology*, 44(3), 737–746. <https://doi.org/10.1016/j.lwt.2010.05.034>
- Ledl, F., & Schleicher, E. (1990). New Aspects of The Maillard Reaction in Foods and In The Human Body. *Angewandte Chemie - International Edition*, 29(6), 565–706. <https://doi.org/10.1002/anie.199005653>
- Lestari, L. A., Huriyati, E., & Marsono, Y. (2017). The development of low glycemic index cookie bars from foxtail millet (*Setaria italica*), arrowroot (*Maranta arundinacea*) flour, and kidney beans (*Phaseolus vulgaris*). *Journal of Food Science and Technology*, 54(6), 1406–1413. <https://doi.org/10.1007/s13197-017-2552-5>
- Lindeboom, N., Chang, P. R., Falk, K. C., & Tyler, R. T. (2005). Characteristics of starch from eight quinoa lines. *Cereal Chemistry*, 82(2), 216–222. <https://doi.org/10.1094/CC-82-0216>
- Liu, Q., Donner, E., Yin, Y., Huang, R. L., & Fan, M. Z. (2006). The physicochemical properties and in vitro digestibility of selected cereals, tubers and legumes grown in China. *Food Chemistry*, 99(3), 470–477. <https://doi.org/10.1016/j.foodchem.2005.08.008>
- Man, J., Cai, J., Cai, C., Xu, B., Huai, H., & Wei, C. (2012). Comparison of physicochemical properties of starches from seed and rhizome of lotus. *Carbohydrate Polymers*, 88(2), 676–683. <https://doi.org/10.1016/j.carbpol.2012.01.016>
- Mancebo, C. M., Picón, J., & Gómez, M. (2015). Effect of flour properties on the quality characteristics of gluten free sugar-snap cookies. *LWT - Food Science and Technology*, 64(1), 264–269. <https://doi.org/10.1016/j.lwt.2015.05.057>
- Mancebo, C. M., Rodriguez, P., & Manuel, G. (2016). Assessing rice flour-starch-protein mixtures to produce gluten free sugar-snap cookies. *LWT - Food Science and Technology*, 67, 127–132. <https://doi.org/10.1016/j.lwt.2015.11.045>
- Mohsen, S. M., Fadel, H. H. M., Bekhit, M. A., Edris, A. E., & Ahmed, M. Y. S. (2009). Effect of substitution of soy protein isolate on aroma volatiles, chemical composition and sensory quality of wheat cookies. *International Journal of Food Science and Technology*, 44(9), 1705–1712. <https://doi.org/10.1111/j.1365-2621.2009.01978.x>
- Palavecino, P. M., Penci, M. C., Calderón-Domínguez, G., & Ribotta, P. D. (2016). Research Article. *Starch - Stärke*, 68((11-12)), 1055–1064. <https://doi.org/10.1002/star.201600111>

- Pasqualone, A., Maria, A., Michele, V., Summo, C., Gambacorta, G., Caponio, F., & Blanco, A. (2015). Production and characterization of functional biscuits obtained from purple wheat. *FOOD CHEMISTRY*, 180, 64–70. <https://doi.org/10.1016/j.foodchem.2015.02.025>
- Pico, J., Tapia, J., Bernal, J., & Gómez, M. (2017). Comparison of different extraction methodologies for the analysis of volatile compounds in gluten-free flours and corn starch by GC/QTOF. *Food Chemistry*, 267, 303–312. <https://doi.org/10.1016/j.foodchem.2017.06.157>
- Rababah, T. M., Al-Mahasneh, M. A., & Ereifej, K. I. (2006). Effect of Chickpea, Broad Bean, or Isolated Soy Protein Additions on the Physicochemical and Sensory Properties of Biscuits. *Journal of Food Science*, 71(6), S438–S442. <https://doi.org/10.1111/j.1750-3841.2006.00077.x>
- Rai, S., Kaur, A., & Singh, B. (2014). Quality characteristics of gluten free cookies prepared from different flour combinations. *Journal of Food Science and Technology*, 51(4), 785–789. <https://doi.org/10.1007/s13197-011-0547-1>
- Rao, B. D., Anis, M., Kalpana, K., Sunooj, K. V., Patil, J. V., & Ganesh, T. (2015). Influence of milling methods and particle size on hydration properties of sorghum flour and quality of sorghum biscuits. *LWT - Food Science and Technology*, 67, 8–13. <https://doi.org/10.1016/j.lwt.2015.11.033>
- Saeed, S. M. G., Tayyaba, S., Ali, S. A., Tayyab, S., Sayeed, S. A., Ali, R., Mobin, L., & Naz, S. (2020). Evaluation of the potential of Lotus root (*Nelumbo nucifera*) flour as a fat mimetic in biscuits with improved functional and nutritional properties. *CYTA - Journal of Food*, 18(1), 624–634. <https://doi.org/10.1080/19476337.2020.1812727>
- Sakač, M., Pestorić, M., Mišan, A., Nedeljković, N., Jambrec, D., Jovanov, P., Banjac, V., Torbica, A., Hadnadev, M., & Mandić, A. (2015). Antioxidant capacity, mineral content and sensory properties of gluten-free rice and buckwheat cookies. *Food Technology and Biotechnology*, 53(1), 38–47. <https://doi.org/10.17113/ftb.53.01.15.3633>
- Sarabhai, S., Indrani, D., Vijaykrishnaraj, M., Milind, Arun Kumar, V., & Prabhasankar, P. (2015). Effect of protein concentrates, emulsifiers on textural and sensory characteristics of gluten free cookies and its immunochemical validation. *Journal of Food Science and Technology*, 52(6), 3763–3772. <https://doi.org/10.1007/s13197-014-1432-5>
- Shahzad, M. A., Ahmad, N., Ismail, T., Manzoor, M. F., Ismail, A., Ahmed, N., & Akhtar, S. (2020). Nutritional composition and quality characterization of lotus (*Nelumbo nucifera* Gaertn.) seed flour supplemented cookies. *Journal of Food Measurement and Characterization*. <https://doi.org/10.1007/s11694-020-00622-x>

- Singh, P., Arora, A., Strand, T. A., Leffler, D. A., Catassi, C., Green, P. H., Kelly, C. P., Ahuja, V., & Makharia, G. K. (2018). Global Prevalence of Celiac Disease: Systematic Review and Meta-analysis. *Clinical Gastroenterology and Hepatology*, *16*(6), 823-836.e2. <https://doi.org/10.1016/j.cgh.2017.06.037>
- Singthong, J., & Meesit, U. (2017). Characteristic and functional properties of Thai lotus seed (*Nelumbo nucifera*) flours. *International Food Research Journal*, *24*(4), 1414–1421. [http://www.ifrj.upm.edu.my/24 \(04\) 2017/\(10\).pdf](http://www.ifrj.upm.edu.my/24%20(04)%202017/(10).pdf)
- Suarni. (2004). Pemanfaatan Tepung Sorgum untuk Produk Olahan. *Jurnal Litbang Pertanian*, *23*(4), 145–151. <http://203.190.36.42/publikasi/p3234045.pdf>
- Suarni, S. (2009). Prospek Pemanfaatan Tepung Jagung Untuk Kue Kering (Cookies). *Jurnal Penelitian Dan Pengembangan Pertanian*, *28*(2), 63–71. <http://ejurnal.litbang.pertanian.go.id/index.php/jppp/article/view/7774/6738>
- Sukhija, S., Singh, S., & Riar, C. S. (2016). Isolation of starches from different tubers and study of their physicochemical, thermal, rheological and morphological characteristics. *Starch/Stärke*, *68*(1–2), 160–168. <https://doi.org/10.1002/star.201500186>
- Sulieman, A. A., Zhu, K., Peng, W., Hassan, H. A., Obadi, M., Siddeeg, A., & Zhou, H. (2018). Rheological and quality characteristics of composite gluten-free dough and biscuits supplemented with fermented and unfermented *Agaricus bisporus* polysaccharide flour. *Food Chemistry*, *271*, 193–203. <https://doi.org/10.1016/j.foodchem.2018.07.189>
- Sun, Q., Han, Z., Wang, L., & Xiong, L. (2014). Physicochemical differences between sorghum starch and sorghum flour modified by heat-moisture treatment. *Food Chemistry*, *145*, 756–764. <https://doi.org/10.1016/j.foodchem.2013.08.129>
- Tang, Y., Li, X., Zhang, B., Chen, P. X., Liu, R., & Tsao, R. (2015). Characterisation of phenolics, betanins and antioxidant activities in seeds of three *Chenopodium quinoa* Willd. genotypes. *Food Chemistry*, *166*, 380–388. <https://doi.org/10.1016/j.foodchem.2014.06.018>
- Taylor, J. R. N., & Anyango, J. O. (2011). Sorghum Flour and Flour Products: Production, Nutritional Quality, and Fortification. In *Flour and Breads and their Fortification in Health and Disease Prevention*. Elsevier Inc. <https://doi.org/10.1016/B978-0-12-380886-8.10012-1>
- Tegeye, M., Kaur, A., Kaur, J., & Singh, H. (2019). Value added convenience food from composite sorghum-maize-sweet potato flour blends. *Indian Journal of Agricultural Sciences*, *89*(11), 1906–1910. https://www.researchgate.net/publication/337242961_Value_added_convenience_food_from_composite_sorghum-maize-sweet_potato_flour_blends#fullTextFileContent

- Torbica, A., Hadnadev, M., & Dapčević Hadnadev, T. (2012). Rice and buckwheat flour characterisation and its relation to cookie quality. *Food Research International*, 48(1), 277–283. <https://doi.org/10.1016/j.foodres.2012.05.001>
- USDA. (2019a). *Arrowroot Flour*. Food Data Central. <https://fdc.nal.usda.gov/fdc-app.html#/food-details/170684/nutrients>
- USDA. (2019b). Buckwheat flour, whole groat. *Food Data Central*. <https://fdc.nal.usda.gov/fdc-app.html#/food-details/170687/nutrients>
- USDA. (2019c). *Corn flour, whole grain, yellow*. Food Data Central. <https://fdc.nal.usda.gov/fdc-app.html#/food-details/170290/nutrients>
- USDA. (2019d). *Rice flour, white, unenriched*. Food Data Central. <https://fdc.nal.usda.gov/fdc-app.html#/food-details/169714/nutrients>
- USDA. (2019e). *Sorghum flour, whole grain*. Food Data Central. <https://fdc.nal.usda.gov/fdc-app.html#/food-details/168943/nutrients>
- Wang, C., Kovacs, M. I. P., Fowler, D. B., & Holley, R. (2004). Effects of Protein Content and Composition on White Noodle Making Quality: Color 1. *Cereal Chemistry*, 81(6), 777–784. <https://doi.org/10.1094/CCHEM.2004.81.6.777>
- Wang, S., Opassathavorn, A., & Zhu, F. (2015). Influence of Quinoa Flour on Quality Characteristics of Cookie, Bread and Chinese Steamed Bread. *Journal of Texture Studies*, 46(4), 281–292. <https://doi.org/10.1111/jtxs.12128>
- Wang, W., Zheng, B., & Tian, Y. (2020). Functional group changes and chemical bond-dependent dielectric properties of lotus seed flour with microwave vacuum drying. *Journal of Food Science*, 0(0), 1–8. <https://doi.org/10.1111/1750-3841.15492>
- Wang, X., Reddy, C. K., & Xu, B. (2018). A systematic comparative study on morphological, crystallinity, pasting, thermal and functional characteristics of starches resources utilized in China. *Food Chemistry*, 259(2018), 81–88. <https://doi.org/10.1016/j.foodchem.2018.03.121>
- Wong, K. C., Chong, F. N., & Chee, S. G. (1998). Volatile Constituents of Taro (*Colocasia esculents* (L .) Schott) Volatile Constituents of Taro (*Colocasia esculenta* (L .) Schott). *Journal of Essential Oil Research*, 10(1), 93–95. <https://doi.org/10.1080/10412905.1998.9700849>
- Wulandari, F. (2016). Analisis Kandungan Gizi, Nilai Energi, Dan Uji Organoleptik Cookies Tepung Beras Dengan Substitusi Tepung Sukun. *Jurnal Aplikasi Teknologi Pangan*, 5(3), 107–112. <https://doi.org/10.17728/jatp.183>
- Xu, D., Zhou, X., Lei, C., Shang, Y., Zhao, Y., Wang, Z., Zeng, F., & Liu, G. (2020). Development of biscuits and cookies using raw dehydrated potato flour and its nutritional quality and volatile aroma compounds evaluation. *Journal of Food*

Processing and Preservation, 44(7), 1–16. <https://doi.org/10.1111/jfpp.14528>

Yildiz, E., & Gocmen, D. (2020). Use of almond flour and stevia in rice-based gluten-free cookie production. *Journal of Food Science and Technology*, 58, 940–951. <https://doi.org/10.1007/s13197-020-04608-x>

Yilmaz, M. T., Yurt, B., Toker, O. S., Karaman, S., & Bas, A. (2015). A mixture design study to determine interaction effects of wheat, buckwheat, and rice flours in an aqueous model system. *LWT - Food Science and Technology*, 61, 583–589. <https://doi.org/10.1016/j.lwt.2014.11.045>

Yu, D., Chen, J., Ma, J., Sun, H., Yuan, Y., Ju, Q., Teng, Y., Yang, M., Li, W., Fujita, K., Tatsumi, E., & Luan, G. (2018). Effects of different milling methods on physicochemical properties of common buckwheat flour. *LWT - Food Science and Technology*, 92(2018), 220–226. <https://doi.org/10.1016/j.lwt.2018.02.033>

Zhu, F. (2017). Structures, properties, and applications of lotus starches. *Food Hydrocolloids*, 63, 332–348. <https://doi.org/10.1016/j.foodhyd.2016.08.034>

