

## Daftar Pustaka

- Abdel-Moemin, A. R. (2016). Effect of Roselle calyces extract on the chemical and sensory properties of functional cupcakes. *Food Science and Human Wellness*, 5(4), 230–237. <https://doi.org/10.1016/j.fshw.2016.07.003>
- Adriana, L., Pranata, F. S., & Swasti, Y. R. (2020). Kualitas Permen Jelly Sari Buah Pepaya (*Carica papaya* L.) dengan Penambahan Ekstrak Kelopak Bunga Rosella (*Hibiscus sabdariffa* L.). *Jurnal Gizi Dan Pangan Soedirman*, 4(1), 26. <https://doi.org/10.20884/1.jgps.2020.4.1.2601>
- Alappat, B., & Alappat, J. (2020). Anthocyanin pigments: Beyond aesthetics. *Molecules*, 25(23). <https://doi.org/10.3390/molecules25235500>
- Alfian, R., & Susanti, H. (2012). Penetapan Kadar Fenolik Total Ekstrak Metanol Kelopak Bunga Rosella Merah (*Hibiscus sabdariffa* Linn) dengan Variasi Tempat Tumbuh secara Spektrofotometri. *Jurnal Ilmiah Kefarmasian*, 2(1). <https://doi.org/10.12928/pharmaciana.v2i1.655>
- Ali, F., Ferawati, & Arqomah, R. (2013). Ekstraksi Zat Warna dari Kelopak Bunga Rosella (Study Pengaruh Konsentrasi Asam Asetat dan Asam Sitrat). *Jurnal Teknik Kimia*, 19(1), 26–34. Retrieved from <http://jtk.unsri.ac.id/index.php/jtk/article/view/126>
- Amogne, N. Y., Ayele, D. W., & Tsigie, Y. A. (2020). Recent Advances in Anthocyanin Dyes Extracted from Plants for Dye Sensitized Solar Cell. *Materials for Renewable and Sustainable Energy*, 9(4), 1–16. <https://doi.org/10.1007/s40243-020-00183-5>
- Amr, A., & Al-Tamimi, E. (2007). Stability of the Crude Extracts of *Ranunculus asiaticus* Anthocyanins and their Use as Food Colourants. *International Journal of Food Science and Technology*, 42(8), 985–991. <https://doi.org/10.1111/j.1365-2621.2006.01334.x>
- Andzi Barhé, T., & Feuya Tchouya, G. R. (2016). Comparative study of the anti-oxidant activity of the total polyphenols extracted from *Hibiscus Sabdariffa* L., *Glycine max* L. Merr., yellow tea and red wine through reaction with DPPH free radicals. *Arabian Journal of Chemistry*, 9(1), 1–8. <https://doi.org/10.1016/j.arabjc.2014.11.048>
- Attahmid, N. F. U., Yusuf, M., Muhtar, I., & Indriati, S. (2020). Karakteristik Fisikokimia Permen Jelly Bengkuang (*Pachyrhizus erosus*) dengan Penambahan Sorbitol dan Ekstrak Rosella (*Hibiscus sabdariffa*). *Jurnal Sains Dan Teknologi Pangan*, 5(2), 2786–2802. Retrieved from [https://www.researchgate.net/profile/Muhammad-Yusuf-40/publication/341272348\\_Physicochemical\\_Characteristics\\_of\\_Yam\\_Pachyrhizus\\_erosus\\_Jelly\\_Candy\\_with\\_Addition\\_of\\_Sorbitol\\_and\\_Rosella\\_Hibiscus\\_sabdariffa\\_Extract/links/5eb6e2224585152169c1239c/Physicochem](https://www.researchgate.net/profile/Muhammad-Yusuf-40/publication/341272348_Physicochemical_Characteristics_of_Yam_Pachyrhizus_erosus_Jelly_Candy_with_Addition_of_Sorbitol_and_Rosella_Hibiscus_sabdariffa_Extract/links/5eb6e2224585152169c1239c/Physicochem)
- Awe, F. B., Fagbemi, T. N., Ifesan, B. O. T., & Badejo, A. A. (2013). Antioxidant properties of cold and hot water extracts of cocoa, Hibiscus flower extract, and ginger beverage blends. *Food Research International*, 52(2), 490–495. <https://doi.org/10.1016/j.foodres.2013.01.021>
- Baydar, H., Erbaş, S., & Kazaz, S. (2016). Variations in floral characteristics and scent

- composition and the breeding potential in seed-derived oil-bearing Roses (*Rosa damascena* mill.). *Turkish Journal of Agriculture and Forestry*, 40(4), 560–569. <https://doi.org/10.3906/tar-1512-57>
- Bayram, O., Sagdic, O., & Ekici, L. (2015). Natural food colorants and bioactive extracts from some edible flowers. *Journal of Applied Botany and Food Quality*, 88, 170–176. <https://doi.org/10.5073/JABFQ.2015.088.024>
- Bechoff, A., Cissé, M., Fliedel, G., Declémey, A. L., Ayessou, N., Akissoe, N., ... Tomlins, K. I. (2014). Relationships between anthocyanins and other compounds and sensory acceptability of Hibiscus drinks. *Food Chemistry*, 148, 112–119. <https://doi.org/10.1016/j.foodchem.2013.09.132>
- Benvenuti, S., Bortolotti, E., & Maggini, R. (2016). Antioxidant power, anthocyanin content and organoleptic performance of edible flowers. *Scientia Horticulturae*, 199, 170–177. <https://doi.org/10.1016/j.scienta.2015.12.052>
- Borrás-Linares, I., Fernández-Arroyo, S., Arráez-Roman, D., Palmeros-Suárez, P. A., Del Val-Díaz, R., Andrade-González, I., ... Segura-Carretero, A. (2015). Characterization of phenolic compounds, anthocyanidin, antioxidant and antimicrobial activity of 25 varieties of Mexican Roselle (*Hibiscus sabdariffa*). *Industrial Crops and Products*, 69, 385–394. <https://doi.org/10.1016/j.indcrop.2015.02.053>
- Boskabady, M. H., Shafei, M. N., Saberi, Z., & Amini, S. (2011). Pharmacological effects of *Rosa damascena*. *Iranian Journal of Basic Medical Sciences*, 14(4), 295–307. <https://doi.org/10.22038/ijbms.2011.5018>
- Branen, A. L. (1975). Toxicology and biochemistry of butylated hydroxyanisole and butylated hydroxytoluene. *Journal of the American Oil Chemists' Society*, 52(2), 59–63. <https://doi.org/10.1007/BF02901825>
- Caleja, C., Barros, L., Antonio, A. L., Oliveira, M. B. P. P., & Ferreira, I. C. F. R. (2017). A comparative study between natural and synthetic antioxidants: Evaluation of their performance after incorporation into biscuits. *Food Chemistry*, 216, 342–346. <https://doi.org/10.1016/j.foodchem.2016.08.075>
- Carocho, M., Barreiro, M. F., Morales, P., & Ferreira, I. C. F. R. (2014). Adding molecules to food, pros and cons: A review on synthetic and natural food additives. *Comprehensive Reviews in Food Science and Food Safety*, 13(4), 377–399. <https://doi.org/10.1111/1541-4337.12065>
- Carocho, M., Morales, P., & Ferreira, I. C. F. R. (2015). Natural food additives: Quo vadis? *Trends in Food Science and Technology*, 45(2), 284–295. <https://doi.org/10.1016/j.tifs.2015.06.007>
- Castañeda-Ovando, A., Pacheco-Hernández, M. de L., Páez-Hernández, M. E., Rodríguez, J. A., & Galán-Vidal, C. A. (2009). Chemical studies of anthocyanins: A review. *Food Chemistry*, 113(4), 859–871. <https://doi.org/10.1016/j.foodchem.2008.09.001>
- Celli, G. B., & Brooks, M. S. L. (2017). Impact of extraction and processing conditions on betalains and comparison of properties with anthocyanins — A current review.

*Food Research International*, 100, 501–509.  
<https://doi.org/10.1016/j.foodres.2016.08.034>

- Chairunnisa, S., Wartini, N. M., & Suhendra, L. (2019). Pengaruh Suhu dan Waktu Maserasi terhadap Karakteristik Ekstrak Daun Bidara (*Ziziphus mauritiana* L.) sebagai Sumber Saponin. *Jurnal Rekayasa Dan Manajemen Agroindustri*, 7(4), 551. <https://doi.org/10.24843/jrma.2019.v07.i04.p07>
- Chaiyasut, C., Sivamaruthi, B. S., Pengkumsri, N., Sirilun, S., Peerajan, S., Chaiyasut, K., & Kesika, P. (2016). Anthocyanin profile and its antioxidant activity of widely used fruits, vegetables, and flowers in Thailand. *Asian Journal of Pharmaceutical and Clinical Research*, 9(6), 218–224. <https://doi.org/10.22159/ajpcr.2016.v9i6.14245>
- Chanukya, B. S., & Rastogi, N. K. (2016). A Comparison of Thermal Processing, Freeze Drying and Forward Osmosis for the Downstream Processing of Anthocyanin from Rose Petals. *Journal of Food Processing and Preservation*, 40(6), 1289–1296. <https://doi.org/10.1111/jfpp.12714>
- Chanwitheesuk, A., Teerawutgulrag, A., & Rakariyatham, N. (2005). Screening of antioxidant activity and antioxidant compounds of some edible plants of Thailand. *Food Chemistry*, 92(3), 491–497. <https://doi.org/10.1016/j.foodchem.2004.07.035>
- Chayaratanaasin, P., Barbieri, M. A., Suanpairintr, N., & Adisakwattana, S. (2015). Inhibitory effect of *Clitoria ternatea* flower petal extract on fructose-induced protein glycation and oxidation-dependent damages to albumin in vitro. *BMC Complementary and Alternative Medicine*, 15(1), 1–9. <https://doi.org/10.1186/s12906-015-0546-2>
- Chung, C., Rojanasasithara, T., Mutilangi, W., & McClements, D. J. (2015). Enhanced stability of anthocyanin-based color in model beverage systems through whey protein isolate complexation. *Food Research International*, 76, 761–768. <https://doi.org/10.1016/j.foodres.2015.07.003>
- Chung, C., Rojanasasithara, T., Mutilangi, W., & McClements, D. J. (2016). Enhancement of colour stability of anthocyanins in model beverages by gum arabic addition. *Food Chemistry*, 201, 14–22. <https://doi.org/10.1016/j.foodchem.2016.01.051>
- Chusak, C., Ying, J. A. Y., Zhien, J. L., Pasukamonset, P., Henry, C. J., Ngamukote, S., & Adisakwattana, S. (2019). Impact of *Clitoria ternatea* (butterfly pea)flower on in vitro starch digestibility, texture and sensory attributes of cooked rice using domestic cooking methods. *Food Chemistry*, 295(May), 646–652. <https://doi.org/10.1016/j.foodchem.2019.05.157>
- Clydesdale, F. M. (1993). Color as a factor in food choice. *Critical Reviews in Food Science and Nutrition*, 33(1), 83–101. <https://doi.org/10.1080/10408399309527614>
- Connor, A. M., Luby, J. J., Tong, C. B. S., Finn, C. E., & Hancock, J. F. (2002). Genotypic and environmental variation in antioxidant activity, total phenolic content, and anthocyanin content among blueberry cultivars. *Journal of the American Society for Horticultural Science*, 127(1), 89–97. <https://doi.org/10.21273/jashs.127.1.89>

- Cortez, R., Luna-Vital, D. A., Margulis, D., & Gonzalez de Mejia, E. (2017). Natural Pigments: Stabilization Methods of Anthocyanins for Food Applications. *Comprehensive Reviews in Food Science and Food Safety*, 16(1), 180–198. <https://doi.org/10.1111/1541-4337.12244>
- Coultate, T., & Blackburn, R. S. (2018). Food colorants: their past, present and future. *Coloration Technology*, 134(3), 165–186. <https://doi.org/10.1111/cote.12334>
- Da-Costa-Rocha, I., Bonnlaender, B., Sievers, H., Pischel, I., & Heinrich, M. (2014). Hibiscus sabdariffa L. - A phytochemical and pharmacological review. *Food Chemistry*, 165, 424–443. <https://doi.org/10.1016/j.foodchem.2014.05.002>
- Dangles, O., & Fenger, J. A. (2018). The chemical reactivity of anthocyanins and its consequences in food science and nutrition. *Molecules*, 23(8). <https://doi.org/10.3390/molecules23081970>
- Domínguez-Rodríguez, G., Marina, M. L., & Plaza, M. (2017). Strategies for the extraction and analysis of non-extractable polyphenols from plants. *Journal of Chromatography A*, 1514, 1–15. <https://doi.org/10.1016/j.chroma.2017.07.066>
- Fajarwati, N. H., Parnanto, N. H. R., & Manuhara, G. J. (2017). Pengaruh konsentrasi asam sitrat dan suhu pengeringan terhadap karakteristik fisik, kimia dan sensoris manisan kering labu siam (*Sechium edule* Sw.) dengan pemanfaatan pewarna alami Dari Ekstrak Rosela Ungu (*Hibiscus sabdariffa* L.). *Jurnal Teknologi Hasil Pertanian*, 10(1), 50–66. Retrieved from <https://jurnal.uns.ac.id/ilmupangan/article/view/17494>
- Figueroa-Robles, A., Antunes-Ricardo, M., & Guajardo-Flores, D. (2021). Encapsulation of phenolic compounds with liposomal improvement in the cosmetic industry. *International Journal of Pharmaceutics*, 593, 120125. <https://doi.org/10.1016/j.ijpharm.2020.120125>
- Fizriani, A., Quddus, A. A., & Hariadi, H. (2021). Pengaruh Penambahan Ekstrak Bunga Telang Terhadap Sifat Kimia dan Organoleptik Pada Produk Minuman Cendol. *Jurnal Ilmu Pangan Dan Hasil Pertanian*, 4(2), 136–145. <https://doi.org/10.26877/jiph.v4i2.7516>
- Ge, J., Yue, P., Chi, J., Liang, J., & Gao, X. (2018). Formation and stability of anthocyanins-loaded nanocomplexes prepared with chitosan hydrochloride and carboxymethyl chitosan. *Food Hydrocolloids*, 74, 23–31. <https://doi.org/10.1016/j.foodhyd.2017.07.029>
- Havananda, T., & Luengwilai, K. (2019). Variation in floral antioxidant activities and phytochemical properties among butterfly pea (*Clitoria ternatea* L.) germplasm. *Genetic Resources and Crop Evolution*, 66(3), 645–658. <https://doi.org/10.1007/s10722-018-00738-6>
- He, J., & Monica Giusti, M. (2010). Anthocyanins: Natural colorants with health-promoting properties. *Annual Review of Food Science and Technology*, 1(1), 163–187. <https://doi.org/10.1146/annurev.food.080708.100754>
- Hnin, K. K., Zhang, M., & Wang, B. (2021). Development of nutritional properties in cookies with the incorporation of different levels of rose flower powder by

- microwave-vacuum drying. *Drying Technology*, 39(9), 1136–1148. <https://doi.org/10.1080/07373937.2020.1725035>
- Hu, Z., Tang, X., Liu, J., Zhu, Z., & Shao, Y. (2017). Effect of parboiling on phytochemical content, antioxidant activity and physicochemical properties of germinated red rice. *Food Chemistry*, 214, 285–292. <https://doi.org/10.1016/j.foodchem.2016.07.097>
- Ihsanto, E., & Hidayat, S. (2014). Rancang Bangun Sistem Pengukuran Ph Meter dengan Menggunakan Mikrokontroller Arduino Uno. *Jurnal Teknologi Elektro*, 5(3). <https://doi.org/10.22441/jte.v5i3.769>
- Isnaini, L. (2010). Ekstraksi Pewarna Merah Cair Alami Berantiosidan dari Kelopak Bunga Rosella (*Hibiscus sabdariffa L.*) dan Aplikasinya pada Produk Pangan. *Jurnal Teknologi Pertanian*, 11(1), 18–27. Retrieved from <https://jtp.ub.ac.id/index.php/jtp/article/view/303>
- Iwashina, T. (2015). Contribution to flower colors of flavonoids including anthocyanins: A review. *Natural Product Communications*, 10(3), 529–544. <https://doi.org/10.1177/1934578x1501000335>
- Jabeur, I., Pereira, E., Barros, L., Calhelha, R. C., Soković, M., Oliveira, M. B. P. P., & Ferreira, I. C. F. R. (2017). *Hibiscus sabdariffa L.* as a source of nutrients, bioactive compounds and colouring agents. *Food Research International*, 100, 717–723. <https://doi.org/10.1016/j.foodres.2017.07.073>
- Jayawardana, B. C., Yanagihara, M., Han, K. H., Fukushima, M., Sekikawa, M., & Shimada, K. (2012). Effects of anthocyanin-rich colored potato flakes on lipid oxidation, instrumental color evaluation and sensory characteristics in cooked pork sausages. *Food Science and Technology Research*, 18(3), 455–460. <https://doi.org/10.3136/fstr.18.455>
- Jing, P., & Giusti, M. M. (2005). Characterization of anthocyanin-rich waste from purple corncobs (*Zea mays L.*) and its application to color milk. *Journal of Agricultural and Food Chemistry*, 53(22), 8775–8781. <https://doi.org/10.1021/jf051247o>
- José, A., Queiroz, D. M., & Moreira, S. (2019). Production and characterization of mixed pineapple and hibiscus jam Produção e caracterização de geleia mista de abacaxi com hibisco. *Nota Técnica*, 13(83), 155–163. Retrieved from <https://revista.ufrr.br/agroambiente/article/view/5523>
- Juniarka, I. G., Lukitaningsih, E., & Noegrohati, S. (2011). Analisis Aktivitas Antiosidan dan Kandungan Antosianin Total Ekstrak dan Liposom Kelopak Bunga Rosella (*Hibiscus Sabdariffa L.*). *Majalah Obat Tradisional*, 16(3), 115–123. Retrieved from <https://journal.ugm.ac.id/TradMedJ/article/view/8048>
- Kähkönen, M. P., & Heinonen, M. (2003). Antioxidant activity of anthocyanins and their aglycons. *Journal of Agricultural and Food Chemistry*, 51(3), 628–633. <https://doi.org/10.1021/jf025551i>
- Kaisoon, O., Siriamornpun, S., Weerapreeyakul, N., & Meeso, N. (2011). Phenolic compounds and antioxidant activities of edible flowers from Thailand. *Journal of Functional Foods*, 3(2), 88–99. <https://doi.org/10.1016/j.jff.2011.03.002>

- Khaleghi, A., & Khadivi, A. (2020). Morphological characterization of Damask rose (*Rosa × damascena* Herrm.) germplasm to select superior accessions. *Genetic Resources and Crop Evolution*, 67(8), 1981–1997. <https://doi.org/10.1007/s10722-020-00954-z>
- Khoo, H. E., Azlan, A., Tang, S. T., & Lim, S. M. (2017). Anthocyanidins and anthocyanins: Colored pigments as food, pharmaceutical ingredients, and the potential health benefits. *Food and Nutrition Research*, 61(1). <https://doi.org/10.1080/16546628.2017.1361779>
- Kogawa, K., Kazuma, K., Kato, N., Noda, N., & Suzuki, M. (2007). Biosynthesis of malonylated flavonoid glycosides on the basis of malonyltransferase activity in the petals of *Clitoria ternatea*. *Journal of Plant Physiology*, 164(7), 886–894. <https://doi.org/10.1016/j.jplph.2006.05.006>
- Kohno, Y., Kitamura, S., Yamada, T., Sugihara, K., & Ohta, S. (2005). Production of superoxide radical in reductive metabolism of a synthetic food-coloring agent, indigocarmine, and related compounds. *Life Sciences*, 77(6), 601–614. <https://doi.org/10.1016/j.lfs.2004.08.045>
- Kristiana, H. D., Ariviani, S., & Khasanah, L. U. (2012). Ekstraksi Pigmen Antosianin Buah Senggani (*Melastoma malabathricum* Auct. non Linn) dengan Variasi Jenis Pelarut. *Jurnal Teknossains Pangan*, 1(1), 105–109. Retrieved from <https://jurnal.uns.ac.id/teknossains-pangan/article/viewFile/4198/3622>
- Kumari, P., Ujala, & Bhargava, B. (2021). Phytochemicals from edible flowers: Opening a new arena for healthy lifestyle. *Journal of Functional Foods*, 78(January), 104375. <https://doi.org/10.1016/j.jff.2021.104375>
- Laksmi, R. T., M, L. A., & Kusrahayu. (2012). Daya Ikat Air, pH dan Sifat Organoleptik Chicken Nugget yang Disubtitusi dengan Telur Rebus. *Animal Agriculture Journal*, 1(1), 453–460. Retrieved from <https://ejournal3.undip.ac.id/index.php/aaj/article/viewFile/686/687>
- Loghmani-Khouzani, H., Sabzi Fini, O., & Safari, J. (2007). Essential oil composition of *Rosa damascena* Mill cultivated in central Iran. *Scientia Iranica*, 14(4), 316–319. Retrieved from [http://scientiariranica.sharif.edu/article\\_2858.html](http://scientiariranica.sharif.edu/article_2858.html)
- Lu, B., Li, M., & Yin, R. (2016). Phytochemical Content, Health Benefits, and Toxicology of Common Edible Flowers: A Review (2000–2015). *Critical Reviews in Food Science and Nutrition*, 56(October), S130–S148. <https://doi.org/10.1080/10408398.2015.1078276>
- Lung, J. P., & Destiani, D. . (2018). Uji Aktivitas Antioksidan Vitamin A, C, E dengan Metode DPPH. *Farmaka*, 15(1), 53–62. Retrieved from <http://journal.unpad.ac.id/farmaka/article/view/12805>
- Mahboubi, M. (2016). *Rosa damascena* as holy ancient herb with novel applications. *Journal of Traditional and Complementary Medicine*, 6(1), 10–16. <https://doi.org/10.1016/j.jtcme.2015.09.005>
- Maleta, H. S., Indrawati, R., Limantara, L., & Broto Sudarmo, T. H. P. (2018). Ragam Metode Ekstraksi Karotenoid dari Sumber Tumbuhan dalam Dekade Terakhir

- (Telaah Literatur). *Jurnal Rekayasa Kimia & Lingkungan*, 13(1), 40–50. <https://doi.org/10.23955/rkl.v13i1.10008>
- Martins, N., Roriz, C. L., Morales, P., Barros, L., & Ferreira, I. C. F. R. (2016). Food colorants: Challenges, opportunities and current desires of agro-industries to ensure consumer expectations and regulatory practices. *Trends in Food Science and Technology*, 52, 1–15. <https://doi.org/10.1016/j.tifs.2016.03.009>
- Matyjaszczyk, E., & Śmiechowska, M. (2019). Edible flowers. Benefits and risks pertaining to their consumption. *Trends in Food Science and Technology*, 91(March), 670–674. <https://doi.org/10.1016/j.tifs.2019.07.017>
- Mgaya-Kilima, B., Remberg, S. F., Chove, B. E., & Wicklund, T. (2015). Physiochemical and antioxidant properties of roselle-mango juice blends; Effects of packaging material, storage temperature and time. *Food Science and Nutrition*, 3(2), 100–109. <https://doi.org/10.1002/fsn3.174>
- Monteiro, M. J. P., Costa, A. I. A., Fliedel, G., Cissé, M., Bechoff, A., Pallet, D., ... Pintado, M. M. E. (2017). Chemical-sensory properties and consumer preference of hibiscus beverages produced by improved industrial processes. *Food Chemistry*, 225, 202–212. <https://doi.org/10.1016/j.foodchem.2016.11.127>
- Mortensen, A. (2006). Carotenoids and other pigments as natural colorants. *Pure and Applied Chemistry*, 78(8), 1477–1491. <https://doi.org/10.1351/pac200678081477>
- Mukherjee, P. K., Kumar, V., Kumar, N. S., & Heinrich, M. (2008). The Ayurvedic medicine Clitoria ternatea-From traditional use to scientific assessment. *Journal of Ethnopharmacology*, 120(3), 291–301. <https://doi.org/10.1016/j.jep.2008.09.009>
- Nguyen, T. T., Phan-Thi, H., Pham-Hoang, B. N., Ho, P. T., Tran, T. T. T., & Waché, Y. (2018). Encapsulation of Hibiscus sabdariffa L. anthocyanins as natural colours in yeast. *Food Research International*, 107, 275–280. <https://doi.org/10.1016/j.foodres.2018.02.044>
- Nn, A. (2015). A Review on the Extraction Methods Use in Medicinal Plants, Principle, Strength and Limitation. *Medicinal & Aromatic Plants*, 04(03), 3–8. <https://doi.org/10.4172/2167-0412.1000196>
- Nooshkam, M., Varidi, M., & Bashash, M. (2019). The Maillard reaction products as food-born antioxidant and antibrowning agents in model and real food systems. *Food Chemistry*, 275(September 2018), 644–660. <https://doi.org/10.1016/j.foodchem.2018.09.083>
- Palimbong, S., & Pariama, A. S. (2020). Potensi Ekstrak Bunga Telang (Clitoria ternatea Linn) sebagai Pewarna pada Produk Tape Ketan. *Jurnal Sains Dan Kesehatan*, 2(3), 228–235. Retrieved from <https://jsk.farmasi.unmul.ac.id/index.php/jsk/article/view/147>
- Pasukamonset, P., Kwon, O., & Adisakwattana, S. (2017). Oxidative Stability of Cooked Pork Patties Incorporated with Clitoria ternatea Extract (Blue Pea Flower Petal) During Refrigerated Storage. *Journal of Food Processing and Preservation*, 41(1), 1–10. <https://doi.org/10.1111/jfpp.12751>

- Pasukamonset, P., Pumalee, T., Sanguansuk, N., Chumyen, C., Wongvasu, P., Adisakwattana, S., & Ngamukote, S. (2018). Physicochemical, antioxidant and sensory characteristics of sponge cakes fortified with *Clitoria ternatea* extract. *Journal of Food Science and Technology*, 55(8), 2881–2889. <https://doi.org/10.1007/s13197-018-3204-0>
- Pham, T. N., Le, X. T., Nguyen, P. T. N., Tran, T. H., Dao, T. P., Nguyen, D. H., ... Anh, H. L. T. (2020). Effects of storage conditions on total anthocyanin content of Butterfly pea flower (*Clitoria ternatea* L.). *IOP Conference Series: Materials Science and Engineering*, 736(6). <https://doi.org/10.1088/1757-899X/736/6/062005>
- Pires, T. C. S. P., Barros, L., Santos-Buelga, C., & Ferreira, I. C. F. R. (2019). Edible flowers: Emerging components in the diet. *Trends in Food Science and Technology*, 93, 244–258. <https://doi.org/10.1016/j.tifs.2019.09.020>
- Pires, T. C. S. P., Dias, M. I., Barros, L., Barreira, J. C. M., Santos-Buelga, C., & Ferreira, I. C. F. R. (2018). Incorporation of natural colorants obtained from edible flowers in yogurts. *Lwt*, 97, 668–675. <https://doi.org/10.1016/j.lwt.2018.08.013>
- Prabawati, N. B., Oktavirina, V., Palma, M., & Setyaningsih, W. (2021). Edible flowers: Antioxidant compounds and their functional properties. *Horticulturae*, 7(4), 1–22. <https://doi.org/10.3390/horticulturae7040066>
- Pratimasari. (2018). Optimasi Zat Warna Bunga Telang (*Clitoria ternatea*) Sebagai Pewarna Alami Pada Sirup Parasetamol. *Jurnal Ilmiah Manuntung*, 4(2), 89–97. Retrieved from <http://jurnal.akfarsam.ac.id/index.php/jim/article/view/187>
- Prior, R. L., Wu, X., & Schaich, K. (2005). Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements. *Journal of Agricultural and Food Chemistry*, 53(10), 4290–4302. <https://doi.org/10.1021/jf0502698>
- Priska, M., Peni, N., Carvallo, L., & Ngapa, Y. D. (2018). Antosianin dan Pemanfaatannya. *Cakra Kimia (Indonesian E-Journal of Applied Chemistry)*, 6(2), 79–97. Retrieved from <https://ojs.unud.ac.id/index.php/cakra/article/download/46629/28101>
- Purwati, I., Yuwanti, S., & Sari, P. (2016). Karakterisasi Tablet Effervescent Sarang Semut (*Myrmecodia tuberosa*) – Rosella (*Hibiscus sabdariffa* L.) Berbahan Pengisi Maltodekstrin dan Dekstrin. *Jurnal Agroteknologi*, 10(1), 63–72. Retrieved from <http://jurnal.unej.ac.id/index.php/JAGT/article/view/4292>
- Putra, I. R., Asterina, & Isrona, L. (2012). Gambaran Zat Pewarna Merah pada Saus Cabai yang Terdapat pada Jajanan yang Dijual di Sekolah Dasar Negeri Kecamatan Padang Utara. *Jurnal Kesehatan Andalas*, 3(3), 297–303. Retrieved from <https://doi.org/10.25077/jka.v3i3.108>
- Rajbhar, K., Dawda, H., & Mukundan, U. (2015). Polyphenols: methods of extraction. *Sci. Revs. Chem. Commun.*, 5(1), 1–6. Retrieved from [https://www.researchgate.net/profile/Karishma-Rajbhar-2/publication/312554206\\_Polyphenols\\_methods\\_of\\_extraction/links/5881ef7592851c21ff42cf88/Polyphenols-methods-of-extraction.pdf](https://www.researchgate.net/profile/Karishma-Rajbhar-2/publication/312554206_Polyphenols_methods_of_extraction/links/5881ef7592851c21ff42cf88/Polyphenols-methods-of-extraction.pdf)

- Ramirez, M. M., Wysocki, A. F., Ramirez, M. A., Sims, C. A., & Balaban, M. O. (2010). Sensory and marketing characteristics of a hibiscus beverage. *Journal of Food Distribution Research*, 41(3), 52–62. Retrieved from [https://ageconsearch.umn.edu/record/139067/files/Ramirez\\_41\\_3.pdf](https://ageconsearch.umn.edu/record/139067/files/Ramirez_41_3.pdf)
- Ramli, M. E., Salleh, R. M., Tajarudin, H. A., & Zulkurnain, M. (2021). Influence of amylose content on phenolics fortification of different rice varieties with butterfly pea (*Clitoria ternatea*) flower extract through parboiling. *Lwt*, 147(November 2020), 111493. <https://doi.org/10.1016/j.lwt.2021.111493>
- Riaz, G., & Chopra, R. (2018). A review on phytochemistry and therapeutic uses of *Hibiscus sabdariffa* L. *Biomedicine and Pharmacotherapy*, 102(May 2017), 575–586. <https://doi.org/10.1016/j.biopha.2018.03.023>
- Rowe, K. S., & Rowe, K. J. (1994). Synthetic food coloring and behavior: A dose response effect in a double-blind, placebo-controlled, repeated-measures study. *The Journal of Pediatrics*, 125(5 PART 1), 691–698. [https://doi.org/10.1016/S0022-3476\(06\)80164-2](https://doi.org/10.1016/S0022-3476(06)80164-2)
- Różyło, R., Szymańska-Chągot, M., Gawlik-Dziki, U., & Dziki, D. (2021). Spectroscopic, mineral, and antioxidant characteristics of blue colored powders prepared from cornflower aqueous extracts. *Food Chemistry*, 346(July 2020). <https://doi.org/10.1016/j.foodchem.2020.128889>
- Safaei-Ghomí, J., Akhoondi, S., Batooli, H., & Dackhili, M. (2009). Chemical variability of essential oil components of two Rosa x damascena genotypes growing in Iran. *Chemistry of Natural Compounds*, 45(2), 262–264. <https://doi.org/10.1007/s10600-009-9281-6>
- Sengul, M., Sener, D., & Ercisli, S. (2017). The determination of antioxidant capacities and chemical properties of rosa (Rosa damascena mill.) products. *Acta Scientiarum Polonorum, Hortorum Cultus*, 16(4), 63–72. <https://doi.org/10.24326/asphc.2017.4.7>
- Setyaningrum, A., Sumarni, N. K., & Hardi, J. (2017). Sifat Fisiko-Kimia Edible Film Agar – Agar Rumput Laut (*Gracilaria* sp.) Tersubtitusi Glyserol. *Natural Science: Journal of Science and Technology*, 6(2), 136–143. <https://doi.org/10.22487/25411969.2017.v6.i2.8661>
- Setyasih, M., Pranata, F. S., Purwiantiningsih, E., & Kerja, C. (2008). Kualitas Keripik Bengkuang (*Pachyrhizus erosus* (L.) Urb.) dengan Variasi Konsentrasi Ekstrak Bunga Rosela (*Hibiscus sabdariffa* Linn.). *Biota*, 13(1), 14–23. Retrieved from <https://ojs.uajy.ac.id/index.php/biota/article/view/2616>
- Shahid, M., Shahid-Ul-Islam, & Mohammad, F. (2013). Recent advancements in natural dye applications: A review. *Journal of Cleaner Production*, 53, 310–331. <https://doi.org/10.1016/j.jclepro.2013.03.031>
- Sharma, N., Kumari, A., Chunduri, V., Kaur, S., Banda, J., Goyal, A., & Garg, M. (2022). Anthocyanin biofortified black, blue and purple wheat exhibited lower amino acid cooking losses than white wheat. *Lwt*, 154(November 2021), 112802. <https://doi.org/10.1016/j.lwt.2021.112802>

- Shikov, V., Kammerer, D. R., Mihalev, K., Mollov, P., & Carle, R. (2012). Antioxidant capacity and colour stability of texture-improved canned strawberries as affected by the addition of rose (*Rosa damascena* Mill.) petal extracts. *Food Research International*, 46(2), 552–556. <https://doi.org/10.1016/j.foodres.2011.04.004>
- Sigurdson, G. T., Tang, P., & Giusti, M. M. (2017). Natural Colorants: Food Colorants from Natural Sources. *Annual Review of Food Science and Technology*, 8(December 2016), 261–280. <https://doi.org/10.1146/annurev-food-030216-025923>
- Silva, S., Costa, E. M., Calhau, C., Morais, R. M., & Pintado, M. E. (2017). Anthocyanin extraction from plant tissues: A review. *Critical Reviews in Food Science and Nutrition*, 57(14), 3072–3083. <https://doi.org/10.1080/10408398.2015.1087963>
- Sineiro, J., Rubilar, M., Franco, D., & Jablonsky, M. (2008). Polyphenols from plant materials: Extraction and antioxidant power. *Electronic Journal of Environmental, Agricultural and Food Chemistry*, 7(8), 3211–3216. Retrieved from [https://www.researchgate.net/profile/Daniel-Franco-Ruiz/publication/228546690\\_Polyphenols\\_from\\_plant\\_materials\\_Extraction\\_and\\_antioxidant\\_power/links/0912f50748b6aeb892000000/Polyphenols-from-plant-materials-Extraction-and-antioxidant-power.pdf](https://www.researchgate.net/profile/Daniel-Franco-Ruiz/publication/228546690_Polyphenols_from_plant_materials_Extraction_and_antioxidant_power/links/0912f50748b6aeb892000000/Polyphenols-from-plant-materials-Extraction-and-antioxidant-power.pdf)
- Tanaka, Y., Sasaki, N., & Ohmiya, A. (2008). Biosynthesis of plant pigments: Anthocyanins, betalains and carotenoids. *Plant Journal*, 54(4), 733–749. <https://doi.org/10.1111/j.1365-313X.2008.03447.x>
- Terahara, N., Toki, K., Saito, N., Honda, T., Matsui, T., & Osajima, Y. (1998). Eight new anthocyanins, ternatins C1-C5 and D3 and preternatins A3 and C4 from young *Clitoria ternatea* flowers. *Journal of Natural Products*, 61(11), 1361–1367. <https://doi.org/10.1021/np980160c>
- Thaipong, K., Boonprakob, U., Crosby, K., Cisneros-Zevallos, L., & Hawkins Byrne, D. (2006). Comparison of ABTS, DPPH, FRAP, and ORAC assays for estimating antioxidant activity from guava fruit extracts. *Journal of Food Composition and Analysis*, 19(6–7), 669–675. <https://doi.org/10.1016/j.jfca.2006.01.003>
- Tsai, P. J., McIntosh, J., Pearce, P., Camden, B., & Jordan, B. R. (2002). Anthocyanin and antioxidant capacity in Roselle (*Hibiscus sabdariffa* L.) extract. *Food Research International*, 35(4), 351–356. [https://doi.org/10.1016/S0963-9969\(01\)00129-6](https://doi.org/10.1016/S0963-9969(01)00129-6)
- Türker, N., & Erdogdu, F. (2006). Effects of pH and temperature of extraction medium on effective diffusion coefficient of anthocyanin pigments of black carrot (*Daucus carota* var. L.). *Journal of Food Engineering*, 76(4), 579–583. <https://doi.org/10.1016/j.jfoodeng.2005.06.005>
- Visita, B. F., & Putri, W. D. R. (2014). Pengaruh Penambahan Bubuk Mawar Merah (*Rosa damascena* Mill) dengan Jenis Bahan Pengisi Berbeda pada Cookies. *Jurnal Pangan Dan Agroindustri*, 2(1), 39–46. Retrieved from <https://jpa.ub.ac.id/index.php/jpa/article/view/20>
- Wang, W. (2000). *Lyophilization and development of solid protein pharmaceuticals*. *International Journal of Pharmaceutics* (Vol. 203). [https://doi.org/10.1016/S0378-5173\(00\)00423-3](https://doi.org/10.1016/S0378-5173(00)00423-3)

- Wicaksono, C. D. G., Crisviantoro, N., Kurnianto, B., & Harismah, K. (2019). Kajian Pembuatan Permen Lunak Rosella Rendah Glukosa Dengan Ekstrak Daun Stevia. *Prosiding SNPBS (Seminar Nasional Pendidikan Biologi Dan Saintek) Ke-4*, (2010), 66–71. Retrieved from <https://publikasiilmiah.ums.ac.id/bitstream/handle/11617/11294/p.66-71.pdf?sequence=1>
- Wulandari, R. (2016). Pengaruh Berbagai Konsentrasi Ekstrak Bunga Mawar Merah (*Rosa damascena* Mill) Terhadap Stabilitas Warna Antosianin Agar-Agar Sebagai Sumber Belajar Biologi. *Jurnal Pendidikan Biologi Indonesia*, 2(1), 48–56. <https://doi.org/10.22219/jpbi.v2i1.3371>
- Yazdani, A. A., & Tavakkoli-Moghaddam, R. (2012). Integration of the fish bone diagram, brainstorming, and AHP method for problem solving and decision making-a case study. *International Journal of Advanced Manufacturing Technology*, 63(5–8), 651–657. <https://doi.org/10.1007/s00170-012-3916-7>
- Zhao, L., Fan, H., Zhang, M., Chitrakar, B., Bhandari, B., & Wang, B. (2019). Edible flowers: Review of flower processing and extraction of bioactive compounds by novel technologies. *Food Research International*, 126, 108660. <https://doi.org/10.1016/j.foodres.2019.108660>

