

BAB VII DAFTAR PUSTAKA

- Alasalvar, C., Pelvan, E., Özdemir, K. S., Kocadagılı, T., Mogol, B. A., Paslı, A. A., Özcan, N., Özçelik, B., & Gökmen, V. (2013). Compositional, nutritional, and functional characteristics of instant teas produced from low- and high-quality black teas. *Journal of Agricultural and Food Chemistry*, *61*(31), 7529–7536. <https://doi.org/10.1021/jf4015137>
- Anderson, R. A., & Polansky, M. M. (2002). Tea enhances insulin activity. *Journal of Agricultural and Food Chemistry*, *50*(24), 7182–7186. <https://doi.org/10.1021/jf020514c>
- Bai, H., Jiang, W., Wang, X., Hu, N., Liu, L., Li, X., Xie, Y., & Wang, S. (2021). Component changes of mulberry leaf tea processed with honey and its application to in vitro and in vivo models of diabetes. *Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment*, *38*(11), 1840–1852. <https://doi.org/10.1080/19440049.2021.1953709>
- Bøhn, S. K., Croft, K. D., Burrows, S., Puddey, I. B., Mulder, T. P. J., Fuchs, D., Woodman, R. J., & Hodgson, J. M. (2014). Effects of black tea on body composition and metabolic outcomes related to cardiovascular disease risk: A randomized controlled trial. *Food and Function*, *5*(7), 1613–1620. <https://doi.org/10.1039/c4fo00209a>
- Chandra, J. P., Yuwono, E. C., Mardiono, B., Studi, P., Komunikasi, D., Seni, F., Petra, U. K., & Surabaya, J. S. (2015). *Perancangan Buku Interaktif Pengenalan dan Pemanfaatan Tanaman Obat Tradisional Menjadi Minuman Teh Herbal*. 1–11.
- Chen, L., Mulder, P. P. J., Peijnenburg, A., & Rietjens, I. M. C. M. (2019). Risk assessment of intake of pyrrolizidine alkaloids from herbal teas and medicines following realistic exposure scenarios. *Food and Chemical Toxicology*,

130(May), 142–153. <https://doi.org/10.1016/j.fct.2019.05.024>

da Silva, L. A., Wouk, J., Weber, V. M. R., Eltchechem, C. da L., de Almeida, P., Martins, J. C. L., Malfatti, C. R. M., & Osiecki, R. (2017). Mechanisms and biological effects of Caffeine on substrate metabolism homeostasis: A systematic review. *Journal of Applied Pharmaceutical Science*, 7(6), 215–221. <https://doi.org/10.7324/JAPS.2017.70632>

Dardashti pour, E., Yaghobian, F., Dehghan, F., & Azarbayjani, M. A. (2021). Forecast of ameliorating effect of dietary flavonol consumption in white tea with or without aerobic training on type 2 diabetes (T2D) in females. *Clinical Nutrition ESPEN*, 45(xxxx), 134–140. <https://doi.org/10.1016/j.clnesp.2021.08.025>

Dekant, W., Fujii, K., Shibata, E., Morita, O., & Shimotoyodome, A. (2017). Safety assessment of green tea based beverages and dried green tea extracts as nutritional supplements. *Toxicology Letters*, 277, 104–108. <https://doi.org/10.1016/j.toxlet.2017.06.008>

Feng, Z., Li, Y., Li, M., Wang, Y., Zhang, L., Wan, X., & Yang, X. (2019). Tea aroma formation from six model manufacturing processes. *Food Chemistry*, 285, 347–354. <https://doi.org/10.1016/j.foodchem.2019.01.174>

Fu, C., Jiang, Y., Guo, J., & Su, Z. (2016). Natural Products with Anti-obesity Effects and Different Mechanisms of Action. *Journal of Agricultural and Food Chemistry*, 64(51), 9571–9585. <https://doi.org/10.1021/acs.jafc.6b04468>

Fu, Y., Yang, J. chao, Cunningham, A. B., Towns, A. M., Zhang, Y., Yang, H. ying, Li, J. wen, & Yang, X. fei. (2018). A billion cups: The diversity, traditional uses, safety issues and potential of Chinese herbal teas. *Journal of Ethnopharmacology*, 222, 217–228. <https://doi.org/10.1016/j.jep.2018.04.026>

Ghuniem, M. M. (2019). Assessment of the Potential Health Risk of Heavy Metal Exposure from the Consumption of Herbal, Black and Green Tea. *Biomedical*

Journal of Scientific & Technical Research, 16(1).
<https://doi.org/10.26717/bjstr.2019.16.002806>

Governa, P., Manetti, F., Miraldi, E., & Biagi, M. (2022). Effects of in vitro simulated digestion on the antioxidant activity of different *Camellia sinensis* (L.) Kuntze leaves extracts. *European Food Research and Technology*, 248(1), 119–128. <https://doi.org/10.1007/s00217-021-03864-1>

Grove, K. A., & Lambert, J. D. (2010). Laboratory, epidemiological, and human intervention studies show that tea (*Camellia sinensis*) may be useful in the prevention of obesity. *Journal of Nutrition*, 140(3), 446–453. <https://doi.org/10.3945/jn.109.115972>

Gumantara, M. P. B., & Oktarlina, R. Z. (2017). Perbandingan Monoterapi dan Kombinasi Terapi Sulfonilurea-Metformin terhadap Pasien Diabetes Melitus Tipe 2. *Majority*, 6(1), 55–59.

Guo, T., Song, D., Cheng, L., & Zhang, X. (2019). Interactions of tea catechins with intestinal microbiota and their implication for human health. *Food Science and Biotechnology*. <https://doi.org/10.1007/s10068-019-00656-y>

Guo, X., Cheng, M., Zhang, X., Cao, J., Wu, Z., & Weng, P. (2017). Green tea polyphenols reduce obesity in high-fat diet-induced mice by modulating intestinal microbiota composition. *International Journal of Food Science and Technology*, 52(8), 1723–1730. <https://doi.org/10.1111/ijfs.13479>

Guyenet, S. J., & Schwartz, M. W. (2012). Regulation of food intake, energy balance, and body fat mass: Implications for the pathogenesis and treatment of obesity. *Journal of Clinical Endocrinology and Metabolism*, 97(3), 745–755. <https://doi.org/10.1210/jc.2011-2525>

Han, M. J., Kim, S. H., Shin, J. ho, & Hwang, J. H. (2021). Caffeine-induced hypokalemia: a case report. *BMC Nephrology*, 22(1), 4–9. <https://doi.org/10.1186/s12882-021-02465-0>

Hardoko, Nafi'ah, B., Sasmito, B. B., & Halim, Y. (2019). Antidiabetic activity of

herbal green tea extract from white mangrove (*Avicennia marina*) leaves towards blood glucose level of diabetic wistar rats (*Rattus norvegicus*). *International Journal of Food Studies*, 8(2), 43–52. <https://doi.org/10.7455/ijfs/8.2.2019.a5>

Hastuty, Y. D. (2018). Perbedaan Kadar Kolesterol Orang Yang Obesitas Dengan Orang Yang Non Obesitas. *AVERROUS: Jurnal Kedokteran Dan Kesehatan Malikussaleh*, 1(2), 47. <https://doi.org/10.29103/averrous.v1i2.407>

Henríquez, S., Jara, N., Bunout, D., Hirsch, S., de la Maza, M. P., Leiva, L., & Barrera, G. (2013). Variabilidad de las fórmulas que evalúan la sensibilidad a la insulina y su asociación con el índice matsuda. *Nutricion Hospitalaria*, 28(5), 1594–1598. <https://doi.org/10.3305/nh.2013.28.5.6512>

Hidayanto, F., & Ardi, D. S. (2015). Tanaman herbal sebagai tanaman hias dan tanaman obat. *Jurnal Inovasi Dan Kewirausahaan*, 4(1), 1–4.

Hosseini, S. A., Gorjian, M., Rasouli, L., & Shirali, S. (2015). A comparison between the effect of green tea and Kombucha prepared from green tea on the weight of diabetic rats. *Biomedical and Pharmacology Journal*, 12(April 2015), 141–146. <https://doi.org/10.13005/bbra/1616>

Huang, H., Liao, D., Dong, Y., & Pu, R. (2019). Clinical effectiveness of quercetin supplementation in the management of weight loss: A pooled analysis of randomized controlled trials. *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*, 12, 553–563. <https://doi.org/10.2147/DMSO.S199830>

Iqbal Akbar Asfar, A. M. (2017). Efektifitas Penurunan Kadar Kafein pada Teh Hitam dengan Metode Ekstraksi. *INTEK: Jurnal Penelitian*, 4(2), 100. <https://doi.org/10.31963/intek.v4i2.150>

Jin, D., Xu, Y., Mei, X., Meng, Q., Gao, Y., Li, B., & Tu, Y. (2013). Antiobesity and lipid lowering effects of theaflavins on high-fat diet induced obese rats. *Journal of Functional Foods*, 5(3), 1142–1150. <https://doi.org/10.1016/j.jff.2013.03.011>

- Júdice, P. B., Sardinha, L. B., & Silva, A. M. (2021). Variance in respiratory quotient among daily activities and its association with obesity status. *International Journal of Obesity*, 45(1), 217–224. <https://doi.org/10.1038/s41366-020-0591-x>
- Kadam, R. A., Dhumal, N. D., & Khyade, V. B. (2019). The Mulberry, *Morus alba* (L.): The Medicinal Herbal Source for Human Health. *International Journal of Current Microbiology and Applied Sciences*, 8(04), 2941–2964. <https://doi.org/10.20546/ijcmas.2019.804.341>
- Kim, G. N., & Jang, H. D. (2011). Flavonol Content in the Water Extract of the Mulberry (*Morus alba* L.) Leaf and Their Antioxidant Capacities. *Journal of Food Science*, 76(6), 869–873. <https://doi.org/10.1111/j.1750-3841.2011.02262.x>
- Lee, M. K., Kim, H. W., Lee, S. H., Kim, Y. J., Asamenew, G., Choi, J., Lee, J. W., Jung, H. A., Yoo, S. M., & Kim, J. B. (2019). Characterization of catechins, theaflavins, and flavonols by leaf processing step in green and black teas (*Camellia sinensis*) using UPLC-DAD-QToF/MS. *European Food Research and Technology*, 245(5), 997–1010. <https://doi.org/10.1007/s00217-018-3201-6>
- Lewis, W. H., & Elvin-Lewis, M. P. F. (2003). *Medical botany : plants affecting human health*.
- Li, X., Zhang, Y., Zhao, C., Zhang, B., Peng, B., Zhang, Y., Wang, J., & Wang, S. (2022). Positive effects of Epigallocatechin-3-gallate (EGCG) intervention on insulin resistance and gut microbial dysbiosis induced by bisphenol A. *Journal of Functional Foods*, 93(February), 105083. <https://doi.org/10.1016/j.jff.2022.105083>
- Li, Y., Zheng, M., Zhai, X., Huang, Y., Khalid, A., Malik, A., Shah, P., Karim, S., Azhar, S., & Hou, X. (2015). Effect of *Gymnema sylvestre*, *Citrullus colocynthis* and *Artemisia absinthium* on blood glucose and lipid profile in diabetic human. *Acta Poloniae Pharmaceutica - Drug Research*, 72(5), 981–

985.

- Lim, H. S., Hwang, J. Y., Choi, J. C., & Kim, M. (2015). Assessment of caffeine intake in the Korean population. *Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment*, 32(11), 1786–1798. <https://doi.org/10.1080/19440049.2015.1077396>
- Liu, Y., Ma, T., Chen, D., Zhang, N., Si, B., Deng, K., Tu, Y., & Diao, Q. (2019). Effects of tea saponin supplementation on nutrient digestibility, methanogenesis, and ruminal microbial flora in dorper crossbred ewe. *Animals*, 9(1). <https://doi.org/10.3390/ani9010029>
- Lubis, S. H., Saputri, M., & Hasanah, N. (2021). ORIGINAL ARTICEL (Guazuma ulmifolia Lamk) AGAINST REDUCTION BLOOD TRIGLYCERIDE LEVELS OF WHITE RATS MALE (Rattus norvegicus). *Journal of Pharmaceuticsl and Science*, 4(2), 41–52.
- Ma, Z. W., Zhang, K. N., Zou, Z. J., & Lü, Q. F. (2021). High specific area activated carbon derived from chitosan hydrogel coated tea saponin: One-step preparation and efficient removal of methylene blue. *Journal of Environmental Chemical Engineering*, 9(3), 105251. <https://doi.org/10.1016/j.jece.2021.105251>
- Menezes, I. R. A., Santana, T. I., Varela, V. J. C., Saraiva, R. A., Matias, E. F. F., Boligon, A. A., Athayde, M. L., Coutinho, H. D. M., Costa, J. G. M., & Rocha, J. B. T. (2015). Chemical composition and evaluation of acute toxicological, antimicrobial and modulatory resistance of the extract of *Murraya paniculata*. *Pharmaceutical Biology*, 53(2), 185–191. <https://doi.org/10.3109/13880209.2014.913068>
- Mousavi, A., Vafa, M., Neyestani, T., Khamseh, M., & Hoseini, F. (2014). The effects of green tea consumption on metabolic and anthropometric indices in patients with type 2 diabetes. *Journal of Research in Medical Sciences*, 19(12), 1203–1204.

- Mukhriani. (2014). Ekstraksi, Pemisahan Senyawa, dan Identifikasi Senyawa Aktif. *Jurnal Kesehatan*, 7(2). <https://doi.org/10.17969/agripet.v16i2.4142>
- Murase, T., Haramizu, S., Shimotoyodome, A., Tokimitsu, I., & Hase, T. (2006). Green tea extract improves running endurance in mice by stimulating lipid utilization during exercise. *American Journal of Physiology - Regulatory Integrative and Comparative Physiology*, 290(6), 1550–1556. <https://doi.org/10.1152/ajpregu.00752.2005>
- Nabavi, S. F., Russo, G. L., Daglia, M., & Nabavi, S. M. (2015). Role of quercetin as an alternative for obesity treatment: You are what you eat! *Food Chemistry*, 179, 305–310. <https://doi.org/10.1016/j.foodchem.2015.02.006>
- Nahar, L., Xiao, J., & Sarker, S. D. (2020). Introduction of Phytonutrients. *Handbook of Dietary Phytochemicals*, 1–17. https://doi.org/10.1007/978-981-13-1745-3_2-1
- Naim, F., Marianti, A., & Susanti, R. (2011). Aktifitas Ekstrak Daun Jati Belanda terhadap Kadar Kolesterol HDL dan LDL pada Tikus Hiperkolesterolemia. *Life Science* 9, 5(1), 18–24. <http://journal.unnes.ac.id/sju/index.php/LifeSci>
- Nakov, G., Ivanova, N., Jukic, M., Koceva Komlenic, D., Daniloski, D., & Lukinac, J. (2020). Perceptions of Consumer for Preparation and Consumption of Tea and Analysis of Biologically Active Compounds of Black Tea. *Scientific Papers-Series Management Economic Engineering in Agriculture and Rural Development*, 20(1), 389–398.
- Nuri, N., Puspitasari, E., Hidayat, M. A., Ningsih, I. Y., Triatmoko, B., & Dianasari, D. (2020). Pengaruh Metode Ekstraksi terhadap Kadar Fenol dan Flavonoid Total, Aktivitas Antioksidan serta Antilipase Daun Jati Belanda (*Guazuma ulmifolia*). *Jurnal Sains Farmasi & Klinis*, 7(2), 143. <https://doi.org/10.25077/jsfk.7.2.143-150.2020>
- Pamungkas, J. D., Anam, K., & Kusriani, D. (2016). Penentuan Total Kadar Fenol dari Daun Kersen Segar, Kering dan Rontok (*Muntingia calabura L.*) serta Uji

- Aktivitas Antioksidan dengan Metode DPPH. *Jurnal Kimia Sains Dan Aplikasi*, 19(1), 15. <https://doi.org/10.14710/jksa.19.1.15-20>
- Pan, M. H., Tung, Y. C., Yang, G., Li, S., & Ho, C. T. (2016). Molecular mechanisms of the anti-obesity effect of bioactive compounds in tea and coffee. *Food and Function*, 7(11), 4481–4491. <https://doi.org/10.1039/c6fo01168c>
- Perslami, R., Puruhito, E. F., & Agil, M. (2017). THE EFFECT OF SLIMMING HERBS TEA WITH DIETARY ADJUSTMENT TO THE RESULT OF ANTROPOMETRIC MEASUREMENT AMONG WOMEN MEMBER OF AEROBIC STUDIO. *Journal of Vocasional Health Studies*, 01(6), 32–38. www.e-journal.unair.ac.id/index.php/JVHS
- Pfeuffer, M., Auinger, A., Bley, U., Kraus-Stojanowic, I., Laue, C., Winkler, P., Rüfer, C. E., Frank, J., Bösch-Saadatmandi, C., Rimbach, G., & Schrezenmeir, J. (2013). Effect of quercetin on traits of the metabolic syndrome, endothelial function and inflammation in men with different APOE isoforms. *Nutrition, Metabolism and Cardiovascular Diseases*, 23(5), 403–409. <https://doi.org/10.1016/j.numecd.2011.08.010>
- Quesille-Villalobos, A. M., Torricoa, J. S., & Ranilla, L. G. (2013). Phenolic compounds, antioxidant capacity, and in vitro α -amylase inhibitory potential of tea infusions (*Camellia sinensis*) commercialized in Chile. *CYTA - Journal of Food*, 11(1), 60–67. <https://doi.org/10.1080/19476337.2012.688219>
- Ramappa, V. K., Srivastava, D., Singh, P., Kumar, U., & Singh, V. (2020). Mulberry 1-deoxynojirimycin (DNJ): an exemplary compound for therapeutics. *Journal of Horticultural Science and Biotechnology*, 95(6), 679–686. <https://doi.org/10.1080/14620316.2020.1760738>
- Ramdhan, Tezar dan Aminah, S. (2014). Pengaruh Pemasakan Terhadap Kandungan Antioksidan Sayuran. *Pengaruh Pemasakan Terhadap Kandungan Antioksidan Sayuran*, 4 Nomor 2, 38–39.

- Rezamela, E., H Syahrian, K., Wulansari, R., Prawira-Atmaja, M. I., Agustian, E., & Adilina, I. B. (2020). Polyphenols Content of Indonesian Tea Clones on Optimum Manuring Condition. *IOP Conference Series: Earth and Environmental Science*, 439(1). <https://doi.org/10.1088/1755-1315/439/1/012055>
- Rismayanthi, C., & Purnama, Y. O. (2021). The effect of green tea (*camellia sinensis*) with aerobic exercise for weight loss in obesity adolescents Pengaruh pemberian teh hijau (*camellia sinensis*) dengan latihan aerobik terhadap penurunan berat badan pada remaja obesitas. *Medikora*, 20(2), 162–171.
- Roghani, M., & Baluchnejadmojarad, T. (2010). Hypoglycemic and hypolipidemic effect and antioxidant activity of chronic epigallocatechin-gallate in streptozotocin-diabetic rats. *Pathophysiology*, 17(1), 55–59. <https://doi.org/10.1016/j.pathophys.2009.07.004>
- Sanlier, N., Atik, İ., & Atik, A. (2018). A minireview of effects of white tea consumption on diseases. *Trends in Food Science and Technology*, 82, 82–88. <https://doi.org/10.1016/j.tifs.2018.10.004>
- Seyedan, A., Alshawsh, M. A., Alshagga, M. A., Koosha, S., & Mohamed, Z. (2015). Medicinal Plants and Their Inhibitory Activities against Pancreatic Lipase: A Review. *Evidence-Based Complementary and Alternative Medicine*, 2015. <https://doi.org/10.1155/2015/973143>
- Shavitri, D. M. (2022). Peran Senyawa Katekin Dalam Penurunan Berat Badan Pada Obesitas. *Jurnal Medika Hutama*, 03(02), 402–406.
- Sineke, F. U., Suryanto, E., & Sudewi, S. (2016). PENENTUAN KANDUNGAN FENOLIK DAN SUN PROTECTION FACTOR (SPF) DARI EKSTRAK ETANOL DARI BEBERAPA TONGKOL JAGUNG (*Zea mays L.*). *Pharmacon*, 5(1), 275–283.
- Suh, S. S., Hwang, J., Park, M., Park, H. S., & Lee, T. K. (2014). Phenol content, antioxidant and tyrosinase inhibitory activity of mangrove plants in

- Micronesia. *Asian Pacific Journal of Tropical Medicine*, 7(7), 531–535.
[https://doi.org/10.1016/S1995-7645\(14\)60089-4](https://doi.org/10.1016/S1995-7645(14)60089-4)
- Sukohar, A., Busman, H., Kurniawaty, E., & Pangestu Catur, M. M. S. (2017). Effect of Consumption Kemunings Leaf (*Murraya Paniculata* (L.) Jack) Infuse To Reduce Body Mass Index, Waist Circumference and Pelvis Circumference on Obese Patients. *International Journal of Research in Ayurveda & Pharmacy*, 8(2), 75–78. <https://doi.org/10.7897/2277-4343.08268>
- Sukohar, A., Pasya, A. V., Soleha, T. U., & Sangging, P. R. A. (2017). The effect of kemuning leaves (*Murraya paniculata* (L.) Jack) infusion on SGOT and SGPT enzym activities in obese patients. *Biomedical and Pharmacology Journal*, 10(2), 953–958. <https://doi.org/10.13005/bpj/1190>
- Sukriket, P., Lookhanumarnjao, S., & Bumrungpert, A. (2016). The effect of mulberry leaf tea on postprandial glycemic control and insulin sensitivity: A randomized, placebo-controlled crossover study. *Journal of Pharmacy and Nutrition Sciences*, 6(1), 33–37. <https://doi.org/10.6000/1927-5951.2016.06.01.6>
- Supriani, A., Rosyidah, N. N., & Hardiyanti, T. (2019). Pengaruh PEMBERIAN TEH DAUN JATI BELANDA TERHADAP PERUBAHAN KADAR KOLESTEROL PADA MASYARAKAT PENDERITA HIPERKOLESTEROL. *Journal Ners Community*, 10(1), 85–96.
- Susanty, S., & Bachmid, F. (2016). COMPARISON OF MACERATION AND REFLUX EXTRACTION METHODS TO PHENOLIC LEVELS OF CORN COB EXTRACT (*Zea mays* L.). *Jurnal Konversi*, 5(2), 87.
- Suter, I. K. (2013). PANGAN FUNGSIONAL DAN PROSPEK PENGEMBANGANNYA. *Trends in Food Science & Technology*, 6(2), 66–67. [https://doi.org/10.1016/s0924-2244\(00\)88964-5](https://doi.org/10.1016/s0924-2244(00)88964-5)
- Swanson, S. M. (2004). *Multiple Stressors: Literature Review and Gap Analysis*. Water Environment Research Foundation.

- Syahidah, & Subekti, N. (2019). Biological Activity of Mangrove Leaves Extract (Rhizophora sp.). *IOP Conference Series: Earth and Environmental Science*, 270(1). <https://doi.org/10.1088/1755-1315/270/1/012051>
- T. R., D., G, T., N.F, T., M.G., A., P.F., O., & B. M., S. (2013). White Tea (Camellia Sinensis (L.)): Antioxidant Properties And Beneficial Health Effects. *International Journal of Food Science, Nutrition and Dietetics*, 04(1), 19–26. <https://doi.org/10.19070/2326-3350-130005>
- Taufik, Y., Widiantara, T., & Garnida, Y. (2016). The effect of drying temperature on the antioxidant activity of black mulberry leaf tea (Morus nigra). *Rasayan Journal of Chemistry*, 9(4), 889–895.
- Teshome, K. (2019). Effect of tea processing methods on biochemical composition and sensory quality of black tea (Camellia sinensis (L.) O. Kuntze): A review. *Journal of Horticulture and Forestry*, 11(6), 84–95. <https://doi.org/10.5897/JHF2019.0588>
- Trabelsi, A., El Kaibi, M. A., Abbassi, A., Horchani, A., Chekir-Ghedira, L., & Ghedira, K. (2020). Phytochemical Study and Antibacterial and Antibiotic Modulation Activity of Punica granatum (Pomegranate) Leaves. *Scientifica*, 2020. <https://doi.org/10.1155/2020/8271203>
- Tresia, G. E., Evvyernie, D., & Tiuria, R. (2016). Phytochemical screening and in vitro ovicidal, larvacidal, and nematicidal effects of murraya paniculata (L.) Jack extract on gastrointestinal parasites of goats. *Media Peternakan*, 39(3), 173–179. <https://doi.org/10.5398/medpet.2016.39.3.173>
- Vanstraelen, S., Rahier, J., & Geubel, A. P. (2008). Jaundice as a misadventure of a green tea (camellia sinensis) lover: A case report. *Acta Gastro-Enterologica Belgica*, 71(4), 409–412.
- Vichasilp, C., Nakagawa, K., Sookwong, P., Higuchi, O., Luemunkong, S., & Miyazawa, T. (2012). Development of high 1-deoxynojirimycin (DNJ) content mulberry tea and use of response surface methodology to optimize tea-

- making conditions for highest DNJ extraction. *LWT - Food Science and Technology*, 45(2), 226–232. <https://doi.org/10.1016/j.lwt.2011.09.008>
- Wang, K., Liu, F., Liu, Z., Huang, J., Xu, Z., Li, Y., Chen, J., Gong, Y., & Yang, X. (2011). Comparison of catechins and volatile compounds among different types of tea using high performance liquid chromatograph and gas chromatograph mass spectrometer. *International Journal of Food Science and Technology*, 46(7), 1406–1412. <https://doi.org/10.1111/j.1365-2621.2011.02629.x>
- Yu, M., Gouvinhas, I., & Barros, A. (2021). Variation of the polyphenolic composition and antioxidant capacity of freshly prepared pomegranate leaf infusions over one-day storage. *Antioxidants*, 10(8). <https://doi.org/10.3390/antiox10081187>
- Yuliantari, N. W. A., Widarta, I. W. R., & Permana, I. D. G. M. (2017). Pengaruh Suhu dan Waktu Ekstraksi Terhadap Kandungan Flavonoid dan Aktivitas Antioksidan Daun Sirsak (*Annona muricata* L.) Menggunakan Ultrasonik The Influence of Time and Temperature on Flavonoid Content and Antioxidant Activity of Sirsak Leaf (*Annona mur.* *Media Ilmiah Teknologi Pangan*, 4(1), 35–42.
- Yusiasih, R., Pitoi, M. M., Ariyani, M., Koesmawati, T. A., & Maulana, H. (2019). Anthraquinone in Indonesian infusion tea: analysis by HPLC–UV and risk assessment. *Chemical and Biological Technologies in Agriculture*, 6(1), 1–9. <https://doi.org/10.1186/s40538-019-0155-2>
- Zhang, H., Qi, R., & Mine, Y. (2019). The impact of oolong and black tea polyphenols on human health. *Food Bioscience*, 29, 55–61. <https://doi.org/10.1016/j.fbio.2019.03.009>
- Zhang, S., Takano, J., Murayama, N., Morie, T., Abe, T., Park, I., Seol, J., Ishihara, A., Tanaka, Y., Yajima, K., & Tokuyama, K. (2020). Subacute Ingestion of Caffeine and Oolong Tea Increases Fat Oxidation without Affecting Energy Expenditure and Sleep Architecture: A Randomized, Placebo-Controlled,

Double-Blinded Cross-Over Trial. *Nutrients*, 12(3671), 1–13.

Zhao, C. N., Tang, G. Y., Cao, S. Y., Xu, X. Y., Gan, R. Y., Liu, Q., Mao, Q. Q., Shang, A., & Li, H. Bin. (2019). Phenolic profiles and antioxidant activities of 30 tea infusions from green, black, oolong, white, yellow and dark teas. *Antioxidants*, 8(7), 9–13. <https://doi.org/10.3390/antiox8070215>

Zheng, G., Sayama, K., Okubo, T., Juneja, L. R., & Oguni, I. (2004). Anti-obesity effects of three major components of green tea, catechins, caffeine and theanine, in mice. *In Vivo*, 18(1), 55–62.

Zhou, B., Wang, Z., Yin, P., Ma, B., Ma, C., Xu, C., Wang, J., Wang, Z., Yin, D., & Xia, T. (2022). Impact of prolonged withering on phenolic compounds and antioxidant capability in white tea using LC-MS-based metabolomics and HPLC analysis: Comparison with green tea. *Food Chemistry*, 368(March 2021), 130855. <https://doi.org/10.1016/j.foodchem.2021.130855>

