

PAPER NAME

19.-Indra-Adi-Susianto.pdf

WORD COUNT

3267 Words

CHARACTER COUNT

16747 Characters

PAGE COUNT

8 Pages

FILE SIZE

353.3KB

SUBMISSION DATE

Mar 6, 2023 7:35 AM GMT+7

REPORT DATE

Mar 6, 2023 7:36 AM GMT+7

● 17% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

- 14% Internet database
- 8% Publications database
- Crossref database
- Crossref Posted Content database
- 13% Submitted Works database

● Excluded from Similarity Report

- Manually excluded text blocks

PREVALENCE OF TUBERCULOSIS IN PREGNANT WOMEN IN SEMARANG

Indra Adi Susianto¹⁾, Jessica Christanti²⁾,
Nur Dian Rakhmawati³⁾, Ririn Nurmandhani⁴⁾

¹⁾Department of Obstetrics and Gynecology, Faculty of Medicine,
Universitas Katolik Soegijapranata Semarang

²⁾Department of Public Health, Faculty of Medicine,
Universitas Katolik Soegijapranata Semarang

³⁾ Department of Disease Control and Eradication, Semarang Health Office

⁴⁾Study Program of Public Health, Faculty of Health, Universitas Dian Nuswantoro

ABSTRACT

Background: In Indonesia, the prevalence of TB in pregnant women is comparable to the general population. There is a greater risk to the pregnant woman and her baby if TB disease is not treated. Babies born to women with untreated TB disease may have lower birth weight than those babies born to women without TB. This study aimed to describe the prevalence of TB in pregnant women in Semarang.

Subjects and Method: This study used secondary data obtained from drug-susceptible TB (DS TB) record at Semarang City Health Office, from 2020 to 2022. The target population was pregnant women with TB. Pregnant women with TB were described by sociodemographic, clinical, TB program, variables. The data were described in frequency (n) and percent (%).

Results: A total of 23 pregnant women had been diagnosed with DS TB and reported at the Semarang City Health Office between 2020 and 2022. Seven (31.8%) of the TB pregnant women were diagnosed clinically and 15 (68.2%) bacteriologically. Average age was 28.7, minimum= 19, maximum= 37. Fifteen (68.2%) lived in Semarang. Ten (45.5) received TB treatment at the community health center, and 12 (54.5%) received TB treatment at hospital. Four (18.2%) needed referral to hospital. Six (27.3%) completed treatment. One (4.5%) dropped out. Three (13.6%) were cured. Thirty six (36.4%) were under treatment.

Conclusion: Pregnant women with TB in Semarang have been described by sociodemographic, clinical, and TB program, variables.

Keywords: drug-susceptible TB (DS TB), sociodemography, pregnant women.

Correspondence:

Indra Adi Susianto. Department of Obstetrics and Gynecology, Faculty of Medicine, Universitas Katolik Soegijapranata Semarang. Email: indraadisusianto@unika.ac.id. Mobile: 0811297595.

BACKGROUND

Tuberculosis is still one of the top global health problems. The World Health Organization (WHO) reports that there are 1.3 million deaths caused by tuberculosis (TB), of which 300,000 deaths are accompanied by HIV (WHO, 2018). Indonesia is the third

country with the highest TB cases in the world after India and China. Two-thirds of TB cases in the world are dominated by 8 countries including India, China, Indonesia, the Philippines, Pakistan, Nigeria, Bangladesh and South Africa. In Indonesia alone, the highest prevalence of TB is on the

Islands of Sumatra, Java, and Bali (Kristini and Hamidah, 2020).

TB is one of the most terrible killers of women of childbearing age. Pregnancy may increase the risk of developing TB in previously infected women, especially in the last trimester or early postnatal period. The incidence of TB in pregnant women increases significantly. The increased risk of newborns with TB mothers includes the risk of TB infection in infants, premature and low birth weight, perinatal and neonatal death, and becoming orphans (Harahap and Fauzar, 2019).

The prevalence of pregnant women with TB is comparable or almost the same as the incidence of TB in the general population, so it can be assumed that TB treatment in pregnant women is as important as TB treatment in the general population. It is necessary to map and coordinate health service facilities in the city of Semarang so that pregnant women can screen for TB in view of the high risk. This study aimed to describe the prevalence of TB in pregnant women in Semarang.

SUBJECTS AND METHOD

1. Design Study

This research is a quantitative research with a cross-sectional study. This study used a secondary data source record of drug-susceptible TB (DS TB) for 2020-2022 from the Semarang City Health Office.

2. Population and Sample

This study used secondary data sources for DS TB records for 2020-2022 from the Semarang City Health Office with a total of 3,915 DS TB patients. Then the

researchers filtered the data with inclusion criteria, namely pregnant women. The number of pregnant women with DS TB in that period of the year was 23 pregnant women. Existing data is edited, coded, entered, and cleaned. From the records obtained, data were found for 1 pregnant woman aged 69 years, after confirming the data was removed from the data analysis, so that the data analyzed in this study totaled 22 pregnant women with DS TB.

3. Study Variable

Variables in this study were occupation, place of treatment, domicile, referral status, type of TB diagnosis, anatomical classification, history of DM, history of HIV, OAT guidelines, source of drugs, end result of treatment.

4. Definition Operational of Variables

Work was the main activity carried out by the subject and earns income for this activity and is still being carried out at the time of the interview.

The place of treatment was the choice of health service chosen by the subject to check his health condition.

Domicile was the residence of the research subject.

Referral status was the status of delegation of authority and responsibility for TB cases suffered by subjects when they come to health services.

The type of TB diagnosis was the result of examination of the subject seen from the source of confirmation of disease enforcement.

Anatomical classification was the anatomical location of the tuberculosis disease suffered by the subject.

History of DM was a history of DM disease owned by the subject.

History of HIV was a history of HIV disease owned by the subject.

OAT guidelines were categories of drugs given to subjects in the treatment of tuberculosis.

The source of the drug was the source of funds for treating TB patients to get OAT.

The final result of treatment was the subject's treatment status at the time of data collection.

5. Study Instrument

The data for this study were taken from secondary data records of SO TB for 2020-2022 from the Semarang City Health Office.

6. Data Analysis

The data were described in frequency (n) and percent (%).

RESULTS

Table 1 shows that the average age of pregnant women with TB SO is at a productive age of 28.7 years with a range of 19 years to 37 years.

Table 2 shows that 23 pregnant women have been diagnosed with DS TB and reported to the Semarang City Health Office between 2020 and 2022. A total of 7 (31.8%) pregnant women with TB were diagnosed clinically and 15 (68.2%) bacteriologically. The mean age was 28.7, minimum= 19, maximum= 37. A total of 15 (68.2%) pregnant women lived in Semarang. As many as 10 (45.5%) received TB treatment at the Puskesmas, and 12 (54.5%) received TB treatment at the hospital. A total of 14 (18.2%) required referral to hospital. A total of 6 (27.3%) completed treatment. A total of three (13.6%) recovered. A total of 36 (36.4%) were under treatment.

Table 1. Age of pregnant women with TB SO by age in Semarang City Health Office

Variable	Mean	Minimum	Maximum
Age	28.7	19	37

Table 2. Characteristics of pregnant women with TB SO based on secondary data records from the Semarang City Health Office

Characteristic	Category	Frequency (N)	Percentage (%)
Occupation	Student	1	4.5
	Housewife	11	50
	Non governmental laborer	2	9
	Unknown	1	4.5
	No Work	3	13.6
Treatment place	Public Health Center	4	18.4
	Hospital	10	45.5
Domicile	Semarang City	12	54.5
	Outside Semarang	15	68.2
Referral status	Come alone	7	31.8
	Referral	18	81.8
TB diagnostic type	Clinical	4	18.2
	Bacteriological	7	31.8
	Extrapulmonary TB	15	68.2
		4	18.2

Characteristic	Category	Frequency (N)	Percentage (%)
Anatomical Classification	Pulmonary TB	18	81.2
History of DM	No DMs	10	45.5
	Unknown	12	54.5
History of HIV	HIV negative	7	31.8
	Unknown	15	68.2
OAT Guide	Category 1	21	95.5
	Not standard TB	1	4.5
Drug source	Insurance	1	4.5
	TB program	21	95.5
Treatment end result	Complete treatment	6	27.3
	Dropped out of treatment	1	4.5
	Not evaluated	2	9.1
	Healed	3	13.6
	Unknown	2	9.1
	Process of treatment	8	36.4

DISCUSSION

Based on the results of previous studies, it was found that there was a relationship between age and TB sufferers. Age is one of the predisposing factors that influence behavior which can be related to psychological and physical maturity. High mobility and also productive age activity is a demand for community activities and to fulfill life's needs which allows increasing contact with other people (Sikumbang et al., 2022).

Most pregnant women with DS TB are housewives (50%). The type of work does not affect the growth and proliferation of *Mycobacterium tuberculosis* but does affect the risk factors encountered (Siregar, 2021). People who don't work relatively spend more time at home and can increase the intensity of contact with families who live in one house. The results of the study by Rukmini and Chatarina (2011) showed that the biggest risk factor for the incidence of TB in adults was hou-

sehold contacts (Rukmini and Chatarina, 2011). In addition, damp house conditions and lack of ventilation will increase the risk of TB transmission.

Most were treated at the hospital (54.5%) with most coming alone or without referrals (81.8%). TB treatment can be carried out in health care facilities, both primary health care facilities and above. From the results of the study it was found that more TB patients took medication on their own or without referrals, this was also because the research subjects were pregnant women who carried out routine checks at health care facilities. In addition to this, this also shows that the level of knowledge and attitudes of the community is good. So far, screening for TB suspects has been mostly carried out by Health Center City with community-based partnerships because many people still give a negative stigma to TB sufferers. This is necessary to create the need for quality TB services through the involvement of cadres and former TB patients through the provision of adequate information and outreach about

the importance of TB prevention and control (Wijaya et al., 2013).

DM is a disease with the most common risk factors that occur in TB patients. The TB prevalence increases with the increase in the prevalence of DM. The frequency of DM in TB patients is reported to be around 10-15% and the prevalence of this infection is 2-5 times higher in diabetic patients compared to non-diabetic patients (RI Ministry of Health, 2019). DM can reduce the immune system so that a person can be more susceptible to TB, and vice versa TB sufferers can exacerbate DM because it causes the patient's body to be unable to control blood sugar levels so that TB patients can more easily suffer from DM (Lin et al., 2019).

HIV makes patients more susceptible to tuberculosis due to decreased immunity (Mulyadi and Fitrika, 2020). Likewise with HIV TB, it is necessary to know whether there are HIV comorbidities in TB patients, this will affect treatment decision-making as well as monitoring and evaluating program performance (RI Ministry of Health, 2019). Previous studies have shown that co-morbidities can cause TB patients to have higher risk factors for MDR (Multidrug Resistance Tuberculosis) (Workicho et al., 2017). From the results of the study it was found that the majority of pregnant women with Mulyadi had no known DM status (54.5%) and no known HIV status (68.2%). Determining the status of pregnant women with DS TB related to comorbidities is very important to do.

The most important thing in the healing process for tuberculosis is adherence to taking OAT medication

because taking medication regularly and being obedient is the only way to recover from tuberculosis. Compliance in taking medication means that the patient follows the rules for taking medication and administering it properly (Rokhmah, 2010). This study found that not all OAT drug sources were obtained from government programs, there were still 4.5% of drugs sourced from insurance.

Based on the final results of treatment, there were still 9.1% of patients who were not evaluated at the end of treatment and only 13.6% with cured treatment status. The results of Soeparman's research (2011) explained that due to the long treatment process, it causes boredom for sufferers to continue taking the drug. Besides that, patients also sometimes feel healthy after receiving treatment for a while, even though the treatment has not been completed (Soeparman, 2011). This is due to a lack of knowledge about TB. In addition to knowledge, this is also due to the side effects of OAT which are considered uncomfortable for patients, such as tingling and numbness in the hands and feet, nausea, pink urine, loss of appetite, mild stomach pain, and several other discomforts (Seniantara et al., 2018). WHO launched the 'End Tuberculosis' strategy, which is part of the Sustainable Development Goals, with one goal, namely to end the TB epidemic worldwide. Approaches, methods and strategies are still needed for the development and reduction of TB prevalence. In accordance with the 2013 and 2016 Joint External Monitoring Mission (JEMM) recommenda-

tions and the Semarang city government pays high attention to TB problems, especially TB with pregnancy.

It is necessary to strengthen program management and increase access to quality TB services during ANC examinations at health facilities, by providing technical guidance to human resources at health facilities to make them more competent and effective and strengthening a proactive surveillance system (mandatory notification) based on data obtained from SITH and SEMARBETUL so that it can increase the independence of the community in terms of reporting to health facilities through the RT/RW/Village regarding suspected cases of TB in the neighborhood.

AUTHOR CONTRIBUTION

In this study, the conception of the study is the contribution of Indra Adi Susianto. Data collection was carried out by the writer Indra Adi Susianto. Jessica Christanti, Nur Dian Rakhmawati, and Ririn Nurmandhani. Data analysis and interpretation was performed by Ririn Nurmandhani. Manuscript writing, review of results and writing of the final version of the manuscript were carried out by all authors.

FUNDING AND SPONSORSHIP

There is no funding or sponsorship in this research.

CONFLICT OF INTEREST

There is no conflict of interest in writing this article.

ACKNOWLEDGMENT

The researcher would like to thank the Semarang City Health Office for helping provide data and information related to this research.

REFERENCE

- Harahap Y, Fauzar (2019). Tuberculosis diseminata pada kehamilan (Disseminated tuberculosis in pregnancy). *Majalah Kedokteran Andalas*. 42(3S): 75-84. Retrieved from: <https://doi.org/10.25077/mka.v42.i3s.p75-84.2019>.
- Kristini D, Hamidah R (2020). Potensi penularan tuberculosis paru pada anggota keluarga penderita (The potential for transmission of pulmonary tuberculosis to family members of sufferers). *Jurnal Kesehatan Masyarakat Indonesia*, 15(1):24-28. doi: <https://doi.org/10.26714/jkmi.15.1.2020.24-28>
- Kementerian Kesehatan Republik Indonesia (2019). Keputusan menteri kesehatan republik Indonesia Nomor HK.01.07/Menkes/755/-2019 tentang pedoman nasional pelayanan kedokteran tata laksana tuberculosis (Decree of the Minister of Health of the Republic of Indonesia Number HK.01.-07/Menkes/755/2019 concerning national guidelines for medical services for the management of tuberculosis).
- Lin T, Harries AD, Critchley JA, Owiti P, Dejgaard A, Kumar AMV, Crevel R, Dlodlo RA (2019). Management of diabetes mellitus-tuberculosis. *International Union Against Tuberculosis and Lung*

Disease (The Union) 68, Boulevard Saint Michel, 75006 Paris, France

Mulyadi, Fitrika Y (2020). Hubungan tuberkulosis dengan HIV/AIDS (The relationship between tuberculosis and HIV/AIDS). *Jurnal PSIK-FK Unsyiah*, 11(2): 162-166. Doi: <https://doi.org/10.52199/inj.v2i2.6376>

Rokhmah D (2010). Gender and tuberculosis: the implication to low health care access for the poor. *National Public Health Journal*, 7(10):447-452.

Rukmini, Chatarina U (2011). Faktor-faktor yang berpengaruh terhadap kejadian TB paru dewasa Indonesia (analisis data riset kesehatan dasar tahun 2010) (Factors that influence the incidence of pulmonary tuberculosis in Indonesian adults (2010 basic health research data analysis). *Buletin Penelitian Sistem Kesehatan*, 14(4):320-331.

Seniantara IK, Theresia I, Adang YG (2018). Pengaruh efek samping OAT (obat anti tuberkulosis) terhadap kepatuhan minum obat pada pasien TBC di Puskesmas (The effect of anti-tuberculosis drugs (OAT) side effects on medication adherence in TB patients at the health center). *Jurnal Keperawatan Suaka Insan*. 3(2):1-12. Doi: <https://doi.org/10.51143/jk-si.v3i2.98>

Sikumbang RH, Eyaner PC, Siregar NP (2022). Faktor-faktor yang berhubungan dengan kejadian TB paru pada usia produktif di wilayah kerja Puskesmas Tegal Sari Kecamatan Medan Denai

Tahun 2018. Factors related to the incidence of pulmonary TB at productive age in the working area of the Tegal Sari Health Center, Medan Denai District, 2018). *Ibnu Sina: Jurnal Kedokteran dan Kesehatan*. Fakultas Kedokteran Universitas Islam Sumatera Utara. 21(1):32-43. Retrieved from: <https://jurnal.fk.uisu.ac.id/index.php/ibnusina/article/view/196>

Siregar MD (2021). Hubungan kondisi fisik rumah dengan kejadian tuberkulosis paru (TB paru) di wilayah kerja Puskesmas Sibuhuan tahun 2021 (The relationship between the physical condition of the house and the incidence of pulmonary tuberculosis (pulmonary TB) in the working area of the Sibuhuan Health Center in 2021). Universitas Islam Negeri Sumatera Utara.

Soeparman WS (2011). Buku ilmu penyakit dalam. Balai Penerbitan FKUI.

Wijaya IMW, Murti B, Suriyasa P (2013). Hubungan pengetahuan, sikap, dan motivasi kader kesehatan dengan aktivitasnya dalam pengendalian kasus tuberkulosis di Kabupaten Buleleng (Relationship between knowledge, attitude and motivation of health cadres with their activities in controlling tuberculosis cases in Buleleng Regency). *Jurnal Magister Kedokteran Keluarga*, 1(1): 38-48. Doi: <http://jurnal.pasca.uns.ac.id/>

Workicho A, Kassahun W, Alemseged F (2017). Risk factors for multidrug resistant tuberculosis among tu-

berculosis patients: a case-control study. *Infect Drug Resist.* 2017 Mar 13;10:91-96. doi: 10.2147/IDR.S126274.

World Health Organization (2018). *Global tuberculosis report 2018.*

● 17% Overall Similarity

Top sources found in the following databases:

- 14% Internet database
- 8% Publications database
- Crossref database
- Crossref Posted Content database
- 13% Submitted Works database

TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	jurnal.fk.uisu.ac.id	Internet	2%
2	UM Surabaya on 2023-02-10	Submitted works	1%
3	cdc.gov	Internet	1%
4	Universitas Indonesia on 2023-02-15	Submitted works	1%
5	garuda.kemdikbud.go.id	Internet	<1%
6	midwifery.iocspublisher.org	Internet	<1%
7	nardus.mpn.gov.rs	Internet	<1%
8	Baidi Bukhori, Ema Hidayanti, Dominikus David Biondi Situmorang. "Rel...	Crossref	<1%

9	jurnal.syedzasaintika.ac.id	Internet	<1%
10	Lincoln University College on 2023-01-26	Submitted works	<1%
11	id.scribd.com	Internet	<1%
12	Universitas Islam Indonesia on 2018-12-31	Submitted works	<1%
13	edisciplinas.usp.br	Internet	<1%
14	pasca.uns.ac.id	Internet	<1%
15	repository.ub.ac.id	Internet	<1%
16	Mainul Haque, Tariqul Islam, Nor Azlina A Rahman, Judy McKimm, Adn...	Crossref	<1%
17	ojs.fdk.ac.id	Internet	<1%
18	eprints.undip.ac.id	Internet	<1%
19	journal.uinjkt.ac.id	Internet	<1%
20	media.neliti.com	Internet	<1%

21	Citra Dewi, Adryan Fristiohady, Riezki Amalia, Nur Kusaira Khairul Ikra...	<1%
	Crossref	
22	journal.stikessuakainsan.ac.id	<1%
	Internet	
23	Lincoln University College on 2023-01-25	<1%
	Submitted works	
24	Xinyue Xiao, Gula Da, Xianda Xie, Xiaoqing Liu et al. "Tuberculosis in pa...	<1%
	Crossref	
25	amrsjournals.com	<1%
	Internet	
26	iGroup on 2017-04-02	<1%
	Submitted works	
27	journal.stikessuakainsan.ac.id	<1%
	Internet	
28	journal.unnes.ac.id	<1%
	Internet	
29	repositorio.yachaytech.edu.ec	<1%
	Internet	
30	zora.uzh.ch	<1%
	Internet	
31	Pramita Iriana, Nelly Yardes, Ace Sudrajat, Munifah Syamara Mahanani...	<1%
	Crossref	

● Excluded from Similarity Report

- Manually excluded text blocks

EXCLUDED TEXT BLOCKS

Faculty of Medicine, Universitas

jurnal.unimus.ac.id

Department of Obstetrics and Gynecology

M Nishijo. "Effects of maternal exposure to cadmium on pregnancy outcome and breast milk * COMMENTA...

Study Program of Public Health, Faculty of Health, Universitas Dian Nuswantoro

journal.unnes.ac.id

The 9th International Conference on Public Health Solo, Indonesia, November 23-2...

UM Surabaya on 2023-02-10