

7. DAFTAR PUSTAKA

- Amin, M. N. G., Kusnadi, J., Hsu, J-L., Doerksen, R. J., & Huang, T-C. (2020). Identification of a novel umami peptide in tempeh (Indonesian fermented soybean) and its binding mechanism to the umami receptor T1R. *Food Chemistry*. doi: 10.1016/j.foodchem.2020.127411
- Arai, S., Yamashita, M., & Fujimaki, M. (1972). Glutamyl oligopeptides as factors responsible for tastes of a proteinase-modified soybean protein. *Agricultural and Biological Chemistry*, 36(7): 1253-1256. doi: 10.1080/00021369.1972.10860400
- Aristoy, M. –C., & Toldrá, F. (1991). Deproteinization techniques for HPLC amino acid analysis in fresh pork muscle and dry-cured ham. *Journal of Agriculture and Food Chemistry*, 39: 1792-1795. doi: 10.1021/jf00010a020
- Baharuddin, M., Amat, H., Febryanti, A., Sappewati, & Azis, F. (2021). Characterization of sarcoplasmic protein in the meat using SDS-PAGE method. *IOP Conference Series: Earth and Environmental Science*, 762: 1-5. doi: 10.1088/1755-1315/762/1/012071
- Banes, A. J., & Yamauchi, M. (1983). Reversed-phase high-performance liquid chromatography method for separation of collagen tryptic peptides. *Journal of Chromatography*, 272: 366-372. doi: 10.1016/S0378-4347(00)86140-X
- Barbarino, E., & Lourenço, S. O. (2005). An evaluation of methods for extraction and quantification of protein from marine macro-and microalgae. *Journal of Applied Phycology*, 17(5): 447-460. doi: 10.1007/s10811-005-1641-4
- Barea, P., Melgosa, R., Illera, A. E., Alonso-Riaño, P., Díaz de Cerio, E., Benito-Román, O., Beltrán, S., & Sanz, M. T. (2023). Production of small peptides and low molecular weight amino acids by subcritical water from fish meal: Effect of pressurization agent. *Food Chemistry*, 418. doi: 10.1016/j.foodchem.2023.135925
- Belghit, I., Rasinger, J. D., Heesch, S., Biancarosa, I., Liland, N., Torstensen, B., Waagbo, R., Lock, E-J. & Bruckner, C. G. (2017). In-depth metabolic profiling of marine macroalgae confirms strong biochemical differences between brown, red and green algae. *Algal Research*, 26: 240-249. doi: 10.1016/j.algal.2017.08.001
- Cermeño, M., Kleekayai, T., Amigo-Benavent, M., Harnedy-Rothwell, P. & FitzGerald, R. (2020). Current knowledge on the extraction, purification, identification, and validation of bioactive peptides from seaweed. *Electrophoresis*, 41(20): 1694-1717. doi: 10.1002/elps.202000153
- Chan, P. T. & Matanjun, P. (2016). Chemical composition and physicochemical properties of tropical red seaweed, *Gracilaria changii*. *Food Chemistry*, 221: 302-310. doi: 10.1016/j.foodchem.2016.10.066

- Chaudhari, N., Pereira, E., & Roper, S. D. (2009). Taste receptors for umami: the case for multiple receptors. *The American Journal of Clinical Nutrition*, *90*: 738S-742S. doi: 10.3945/ajcn.2009.27462h
- Choudhury, S. & Sekar, N. S. (2017). Algae as source of natural flavor enhancers – a mini review. *Plant Science Today*, *4*: 172-176. doi: 10.14719/pst.2017.4.4.338
- Claeys, E., De Smet, S., Balcaen, A., Raes, K., & Demeyer, D. (2004). Quantification of fresh meat peptides by SDS-PAGE in relation to ageing time and taste intensity. *Meat Science*, *67*: 281-288. doi: 10.1016/j.meatsci.2003.11.001
- Dang, Y., Gao, X., Ma, F. & Wu, X. (2015). Comparison of umami taste peptides in water-soluble extractions of Jinhua and Parma hams. *LWT – Food Science and Technology*, *60*(2): 1179-1186. doi: 10.1016/j.lwt.2014.09.014
- Dashdorj, D., Amna, T., & Hwang, I. (2015). Influence of specific taste-active components on meat flavor as affected by intrinsic and extrinsic factors: An overview. *European Food Research and Technology*, *241*: 157-171. doi: 10.1007/s00217-015-2449-3
- Dunkel, A., Koster, J. & Hofmann, T. (2007). Molecular and sensory characterization of gamma-glutamyl peptides as key contributors to the kokumi taste of edible beans (*Phaseolus vulgaris* L.). *Journal of Agricultural and Food Chemistry*, *55*(16): 6712-6719. doi: 10.1021/jf071276u
- Dziuba, M., Iwaniak, A., & Minkiewicz, P. (2003). Computer-aided characteristics of proteins as potential precursors of bioactive peptides. *Polimery*, *48*, 2003: 50-53. Diakses dari <https://polimery.ichp.vot.pl/index.php/p/article/view/1836>
- Fan, S-H., Liu, T-T., Wan, P., Zhu, Q., Xia, N., Wang, Q-Z., & Chen, D-W. (2021). Enrichment of the umami-taste-active amino acids and peptides from crab sauce using ethanol precipitation and anion-exchange resin. *Journal of Food Processing and Preservation*. 1-8. doi: 10.1111/jfpp.15390
- Feng, T., Wu, Y., Zhang, Z., Song, S., Zhuang, H., Xu, Z., Yao, L. & Sun, M. (2019). Purification, Identification, and Sensory Evaluation of Kokumi Peptides from *Agaricus bisporus* Mushroom. *Foods*, *8*(43): 1-12. doi: 10.3390/foods8020043
- Gallego, M., Mora, L. & Toldra, F. (2019). The relevance of dipeptides and tripeptides in the bioactivity and taste of dry-cured ham. *Food Production, Processing and Nutrition*, *1*(2). doi: 10.1186/s43014-019-0002-7
- Gallego, M., Toldrá, F., & Mora, L. (2022). Quantification and *in silico* analysis of taste dipeptides generated during dry-cured ham processing. *Food Chemistry*, *370*. doi: 10.1016/j.foodchem.2021.130977
- Gasteiger, E., Hoogland, C., Gattiker, A., Duvaud, S., Wilkins, M. R., Appel, R. D., Bairoch, A. (2005). *Protein Identification and Analysis Tools on the ExPASy Server*, The Proteomics Protocols Handbook, Humana Press. 571-607. Diakses melalui <https://web.expasy.org/protparam/> pada Oktober 2023

- Gomez-Ruiz, J. A., Taborda, G., Amigo, L., Ramos, M., & Molina, E. (2007). Sensory and mass spectrometric analysis of the peptidic fraction lower than one thousand daltons in Manchego Cheese. *Journal of Dairy Science*, 90(11): 4966-4973. doi: 10.3168/jds.2007-0350
- Grigorov, M. G., Schlichtherle-Cerny, H., Affolter, M., & Kochhar, S. (2003). Design of virtual libraries of umami-tasting molecules. *Journal of Chemical Information and Computer Sciences*, 43: 1248-1258. doi: 10.1021/ci025652g
- Guiry, M. D. & Guiry, G. M. (2023a). AlgaeBase. World-wide electronic publication, National University of Ireland, Galway (taxonomic information republished from AlgaeBase with permission of M. D. Guiry). *Carpophyllum plumosum* (A. Richard) J. Agardh, 1878. Diakses melalui World Register of Marine Species di <https://www.marinespecies.org/aphia.php?p=taxdetails&id=373349> pada 12 Oktober 2023
- Guiry, M. D. & Guiry, G. M. (2023b). AlgaeBase. World-wide electronic publication, National University of Ireland, Galway (taxonomic information republished from AlgaeBase with permission of M. D. Guiry). *Sargassum (Sargassum) ilicifolium* (Turner) C. Agardh, 1820. Diakses melalui World Register of Marine Species di <https://www.marinespecies.org/aphia.php?p=taxdetails&id=211937> pada 12 Oktober 2023
- Guiry, M. D. & Guiry, G. M. (2023c). AlgaeBase. World-wide electronic publication, National University of Ireland, Galway (taxonomic information republished from AlgaeBase with permission of M. D. Guiry). *Sargassum thunbergii* (Mertens ex Roth) Kuntze, 1880. Diakses melalui World Register of Marine Species di <https://www.marinespecies.org/aphia.php?p=taxdetails&id=494874> pada 12 Oktober 2023
- Gupta, S., Kapoor, P., Chaudhary, K., Gautam, A., Kumar, R., Open Source Drug Discovery Consortium, & Raghava, G. P. S. (2013). *In silico* approach for predicting toxicity of peptides and proteins. *PLOS ONE*, 8(9): e73957. doi: 10.1371/journal.pone.0073957
- Hajeb, P. & Jinap, S. (2015). Umami Taste Components and Their Sources in Asian Foods, *Critical Reviews in Food Science and Nutrition*, 55(6): 778-791. doi: 10.1080/10408398.2012.678422
- Harnedy, P. & FitzGerald, R. (2013). Extraction of protein from the macroalga *Palmaria palmate*. *LWT-Food Science and Technology*, 51(1): 375-382. doi: 10.1016/j.lwt.2012.09.023.
- Hartley, I. E., Liem, D. G. & Keast, R. (2019). Umami as an 'Alimentary' Taste. A New Perspective on Taste Classification. *Nutrients*, 11, 182: 1-18. doi: 10.3390/nu11010182
- Hazmi, G. G. A. & Harijono. (2019). Pengaruh Pengeringan dan Lama Maserasi dengan Pelarut Ganda Etanol dan Heksana terhadap Senyawa Bioaktif Daging Biji Palem

- Putri (*Veitchia merillii*). *Jurnal Pangan dan Agroindustri*, 7(2): 13-23. doi: 10.21776/ub.jpa.2019.007.02.2
- Hillmann, H. & Hofmann, T. (2016). Quantitation of Key Tastants and Reengineering the Taste of Parmesan Cheese. *Journal of Agricultural and Food Chemistry*. doi: 10.1021/acs.jafc.6b00112
- Ikeda, K. (2002). New Seasonings. *Chemical Senses*, 27(9): 847–849. doi: 10.1093/chemse/27.9.847
- Indumathi, P. & Metha, A., J. (2016). A novel anticoagulant peptide from the Nori hydrolysate. *Journal of Functional Foods*, 20, 606-617. doi: 10.1016/j.jff.2015.11.016
- Ishida, B. K., Ma, J. C., Chan, B. G., Bartley, G. E., & Grossman, J. N. (2001). A modified method for simple, rapid HPLC analysis of lycopene isomers. *Acta Horti*, 542: 235-242. doi: 10.17660/ActaHort.2001.542.30
- Iwaniak, A., Minkiewicz, P., Darewicz, M. & Hryniewicz, M. (2016a). Food protein-originating peptides as tastants – Physiological, technological, sensory, and bioinformatics approaches. *Food Research International*, 89: 27-38. doi: 10.1016/j.foodres.2016.08.010
- Iwaniak, A., Minkiewicz, P., Darewicz, M., Sieniawski, K. & Starowicz, P. (2016b). BIOPEP database of sensory peptides and amino acids. *Food Research International*. doi: 10.1016/j.foodres.2016.04.031
- Ji, P., Zou, J., & Feng, W. (2009). Effect of alcohol on the solubility of amino acid in water. *Journal of Molecular Catalysis B: Enzymatic*, 56: 185-188. doi: 10.1016/j.molcatb.2008.06.008
- Jiang, S., Liu, S., Zhao, C., & Wu, C. (2016). Developing protocols of Tricine-SDS-PAGE for separation of polypeptides in the mass range 1-30 kDa with minigel electrophoresis system. *International Journal of Electrochemical Science*, 11: 640-649. doi: 10.1016/S1452-3981(23)15870-6
- Joubert, Y., & Fleurence, J. (2008). Simultaneous extraction of proteins and DNA by an enzymatic treatment of the cell wall of *Palmaria palmata* (Rhodophyta). *Journal of Applied Phycology*, 20(1): 55-61. doi: 10.1007/s10811-007-9180-9
- Kadi, A. (2005). Beberapa Catatan Kehadiran Marga *Sargassum* Diperairan Indonesia. *Oseana*, 3(4): 19-29. Tersedia pada: www.oseanografi.lipi.go.id
- Kadi, A. (2014). *Seaweed* sebagai produk alami dari perairan Indonesia. *Jurnal Oseana*, 32(3): 31-40. Tersedia pada: http://oseanografi.lipi.go.id/dokumen/os_xxxix_3_2014-4.pdf
- Kazir, M., Abuhassira, Y., Robin, A., Nahor, O., Luo, J., Israel, A., Golberg, A. & Livney, Y. D. (2018). Extraction of proteins from two marine macroalgae, *Ulva* sp. and *Gracilaria* sp., for food application, and evaluating digestibility, amino acid

- composition and antioxidant properties of the protein concentrates. *Food Hydrocolloids*. doi: 10.1016/j.foodhyd.2018.07.047
- Kęska, P. & Stadnik, J. (2017). Taste-active peptides and amino acids of pork meat as components of dry-cured meat products: An *in-silico* study. *Journal of Sensory Studies*, 32(6): 1-9. doi: 10.1111/joss.12301
- Kim, H. & Li-Chan, E. C. Y. (2006). Quantitative structure—Activity relationship of bitter peptides. *Journal of Agricultural and Food Chemistry*, 54: 10102-10111. doi: 10.1021/jf062422j
- Kirimura, J., Shimizu, A., Kimizuka, A., Ninomiya, T., & Katsuya, N. (1969). Contribution of peptides and amino acids to the taste of foods. *Journal of Agricultural and Food Chemistry*, 17: 689-695. doi: 10.1021/jf60164a031
- Kong, Y. Zhang, L. L., Zhao, J. Zhang, Y. Y., Sun, B. G. & Chen, H. T. (2019). Isolation and identification of the umami peptides from shiitake mushroom by consecutive chromatography and LC-Q-TOF-MS. *Food Research International*, 121: 463-470. doi: 10.1016/j.foodres.2018.11.060
- Kose, A. (2021). *In silico* bioactive peptide prediction from the enzymatic hydrolysates of edible seaweed rubisco large chain. *Turkish Journal of Fisheries and Aquatic Sciences*, 21(12): 615-626. doi: 10.4194/1303-2712-v21_12_04
- Kurihara, K. (2015). Umami the Fifth Basic Taste: History of Studies on Receptor Mechanisms and Role as a Food Flavor. *BioMed Research International*. doi: 10.1155/2015/189402
- Kuroda, M., Kato, Y., Yamazaki, J., Kai, Y., Mizukoshi, T., Miyano, H. & Eto, Y. (2012). Determination and Quantification of γ -Glutamyl-valyl-glycine in Commercial Fish Sauces. *Journal of Agricultural and Food Chemistry*, 60: 7291-7296. doi: 10.1021/jf3012932
- Kuroda, M., Kato, Y., Yamazaki, J., Kai, Y., Mizukoshi, T., Miyano, H. & Eto, Y. (2013). Determination and quantification of the *Kokumi* peptide, γ -Glutamyl-valyl-glycine, in commercial soy sauces. *Food Chemistry*. doi: 10.1016/j.foodchem.2013.03.070
- Laohakunjit, N., Selamassakul, O., & Kerdchoechuen. (2014). Seafood-like Flavour obtained from the enzymatic hydrolysis of the protein by-products of seaweed (*Gracilaria* sp.). *Food Chemistry*, 158: 162-170. doi: 10.1016/j.foodchem.2014.02.101
- Legler, G., Müller-Platz, C. M., Mentges-Hettkamp, M., Pflieger, G., & Jülich, E. (1985). On the Chemical Basis of the Lowry Protein Determination. *Analytical Biochemistry*, 150, 278-287. doi: 10.1016/0003-2697(85)90511-1
- Li, Q., Zhang, L. & Lametsch, R. (2020). Current progress in kokumi-active peptides, evaluation and preparation methods: a review. *Critical Reviews in Food Science and Nutrition*. doi: 10.1080/10408398.2020.1837726

- Li, X., Xie, X., Wang, J., Xu, Y., Yi, S., Zhu, W., Mi, H., Li, T. & Li, J. (2019). Identification, taste characteristics and molecular docking study of novel umami peptides derived from the aqueous extract of the clam *Meretrix meretrix* Linnaeus. *Food Chemistry*. doi: 10.1016/j.foodchem.2019.126053
- Lim, S. J., Aida, W. M. W., Yusof, M. M., Mamot, S., Ropien, J., & Mohd, D. M. (2014). Isolation and antioxidant capacity of fucoidan from selected Malaysian seaweeds. *Food Hydrocolloids*. doi: 10.1016/j.foodhyd.2014.03.007.
- Lioe, H. N., Takara, K., & Yasuda, M. (2006). Evaluation of peptide contribution to the intense umami taste of Japanese soy sauces. *Journal of Food Science*, 71: S277-S283. doi: 10.1111/j.1365-2621.2006.tb15654.x
- Liu, Z., Zhu, Y., Wang, W., Zhou, X., Chen, G. & Liu, Y. (2020). Seven novel umami peptides from *Takifugu rubripes* and their taste characteristics. *Food Chemistry*. doi: 10.1016/j.foodchem.2020.127204
- Mæhre, H. K., Dalheim, L., Edvinsen, G. K., Elvevoll, E. O., & Jensen, I-J. (2018). Protein Determination – Method Matters. *Foods*, 7(5): 1-11. doi: 10.3390/foods7010005
- Mattio, L. & Payri, C. E. (2010). 190 years of *Sargassum* taxonomy, facing the advent of DNA phylogenies. *The Botanical Review*, 77(1): 31-70. doi: 10.1007/s12229-010-9060-x
- Mikami, M., Sekikawa, M., & Shimada, K. (1999). Changes in peptides during conditioning of three type of beef samples. *Proceedings of the 45th International Congress of Meat Science and Technology*, 288-289. Yokohama, Japan.
- Milinic, J., Campos, B., Diniz, M., & Noronha, J. (2020). Umami free amino acids in edible green, red, and brown seaweeds from the Portuguese seashore. *Journal of Applied Phycology*, 32(5): 3331-3339. doi: 10.1007/s10811-020-02169-2
- Milinic, J., Mata, P., Diniz, M., & Noronha, J. P. (2021). Umami taste in edible seaweeds: The current comprehension and perception. *International Journal of Gastronomy and Food Science*, 23: 1-9. doi: 10.1016/j.ijgfs.2020.100301
- Minkiewicz, P., Dziuba, J., Iwaniak, A., Dziuba, M. & Darewicz, M. (2008). BIOPEP database and other programs for processing bioactive peptide sequences. *Journal of AOAC International*, 91: 965-980. doi:10.1093/jaoac/91.4.965
- Miyamura, N., Kuroda, M., Kato, Y., Yamazaki, J., Mizukoshi, T., Miyano, H. & Eto, Y. (2014). Determination and Quantification of a *Kokumi* Peptide, γ -Glutamyl-Valyl-Glycine, in Fermented Shrimp Paste Condiments. *Food Science and Technology Research*, 20(3): 699-703. doi: 10.3136/fstr.20.699
- Miyamura, N., Lida, Y., Kuroda, M., Kato, Y., Yamazaki, J., Mizukoshi, T. & Miyano, H. (2015). Determination and quantification of *kokumi* peptide, γ -glutamyl-valyl-glycine, in brewed alcoholic beverages. *Journal of Bioscience and Bioengineering*. doi: 10.1016/j.jbiosc.2015.01.018

- Noble, J. E. & Bailey, M. J. A. (2009). Quantitation of protein. *Methods Enzymol.* 463, 73-95. doi: 10.1016/S0076-6879(09)63008-1
- Noguchi, M., Arai, S., Yamashita, M., Kato, H. & Fujimaki, M. (1975). Isolation and identification of acidic oligopeptides occurring in a flavor-potentiating fraction from a fish protein hydrolysate. *Journal of Agricultural and Food Chemistry*, 23(1): 49-53. doi: 10.1021/jf60197a003
- Oka, S. & Nagata, K. (1974). Isolation and characterization of acidic peptides in soy sauce. *Agricultural and Biological Chemistry*, 38(6): 1195-1202. doi: 10.1080/00021369.1974.10861306
- Pakidi, C. S. & Suwoyo, H. S. (2016). Potensi dan Pemanfaatan Bahan Aktif Alga Cokelat *Sargassum* sp. *Octopus: Jurnal Ilmu Perikanan*, 6(1): 551-562. doi: 10.26618/octopus.v5i2.720
- Peng, Y., Xie, E., Zheng, K., Fredimoses, M., Yang, X., Zhou, X., Wang, Y., Yang, B., Lin, X., Liu, J., & Liu, Y. (2013). Nutritional and chemical composition and antiviral activity of cultivated seaweed *Sargassum naozhouense* Tseng et Lu. *Marine Drugs*, 11(1): 20-32. doi: 10.3390/md11010020
- Pereira, L. (2011). *A review of the nutrient composition of selected edible seaweeds. Seaweed: ecology, nutrient composition and medical uses.* Nova Science Publishers, Inc., USA.
- Pereira, L. (2016). *Edible Seaweeds of the World.* CRC Press, Taylor & Francis Group, LLC, USA.
- Phewpan, A., Phuwaprisirisan, P., Takahashi, H., Ohshima, C., Ngamchuachit, P., Techaruvichit, P., Dirndorfer, S., Dawid, C., Hofmann, T. & Keeratipibul, S. (2020). Investigation of Kokumi Substances and Bacteria in Thai Fermented Freshwater Fish (Pla-ra). *Journal of Agricultural and Food Chemistry*. doi: 10.1021/acs.jafc.9b06107
- Pinsorn, P., Oikawa, A., Watanabe, M., Sasaki, R., Ngamchuachit, P., Hoefgen, R., Saito, K. & Sirikantaramas, S. (2018). Metabolic Variation in the Pulps of Two Durian Cultivars: Unraveling the Metabolites that Contribute to the Flavor, *Food Chemistry*. doi: 10.1016/j.foodchem.2018.06.066
- Poojary, M. M., Orlie, V., Passamonti, P., & Olsen, K. (2017). Improved extraction methods for simultaneous recovery of umami compounds from six different mushrooms. *Journal of Food Composition and Analysis*, 63: 171-183. doi: 10.1016/j.jfca.2017.08.004
- Pratiwi, A. R., Fadlilah, I., Ananingsih, V. K. & Meiliana. (2021). Protein dan asam amino pada edible *Sargassum aquifolium*, *Ulva lactuca*, dan *Gracilariopsis longissima*. *Jurnal Pengolahan Hasil Perikanan Indonesia*, 24(3): 337-346. doi: 10.17844/jphpi.v24i3.37085

- Pratiwi, A. R., Yonata, D., Meiliana & Ananingsih, V. K. (2020). Metode Pembuatan Penyedap Rasa Berbahan Ganggang *Sargassum aquafolium*, Paten Indonesia. S00202009231
- Rahmadhani, R., Ganda Putra, G. P., & Suhendra, L. (2020). Karakteristik Ekstrak Kulit Biji Kakao (*Theobroma cacao* L.) sebagai Sumber Antioksidan pada Perlakuan Ukuran Partikel dan Waktu Maserasi. *Jurnal Rekayasa dan Manajemen Agroindustri*, 8(2): 246. doi: 10.24843/jrma.2020.v08.i02.p09
- Ramos, M. V., Monteiro, A. C. O., Moreira, R. A., & Carvalho, A. D. F. A. F. U. (2000). Amino Acid Composition of Some Brazilian Seaweed Species. *Journal of Food Biochemistry*, 24, 33–39. doi: 10.1111/j.1745-4514.2000.tb00041.x.
- Rhyu, M. R. & Kim, E. Y. (2011). Umami taste characteristics of water extract of *Doenjang*, a Korean soybean paste: Low-molecular acidic peptides may be possible clue to the taste. *Food Chemistry*, 127:1210-1215. doi: 10.1016/j.foodchem.2011.01.128
- Saepudin, E., Sinurat, E., & Suryabrata, I. A. (2018). Depigmentation and Characterization of Fucoidan from Brown Seaweed *Sargassum binderi* Sonder. *IOP Conference Series: Materials Science and Engineering*, 299: 1-5. doi: 10.1088/1757-899X/299/1/012027
- Salger, M., Stark, T. D. & Hofmann, T. (2019). Taste modulating peptides from overfermented cocoa beans. *Journal of Agricultural and Food Chemistry*, 67(15): 4311-4320. doi: 10.1021/acs.jafc.9b00905.
- Schägger, H. (2006). Protocol: Tricine-SDS-PAGE. *Nature Protocols*, 1(1): 16-22. doi: 10.1038/nprot.2006.4
- Schägger, H., & Jagow, G. V. (1987). Tricine-Sodium Dodecyl Sulfate-Polyacrylamide Gel Electrophoresis for the separation of proteins in the range from 1 to 100 kDa. *Analytical Biochemistry*, 166 (2), 368-379. doi: 10.1016/0003-2697(87)90587-2
- Schaich, K. M. (2016). Analysis of Lipid and Protein Oxidation in Fats, Oils, and Foods. *Oxidative Stability and Shelf Life of Foods Containing Oils and Fats*, 1-131. doi: 10.1016/B978-1-63067-056-6.00001-X
- Sharmin, K. N., Amiza, M. A., Ahmad, F., Razali, S. A., & Hashim, F. (2022). *In silico* analysis of *Gracilaria changii* proteins for potential bioactive peptides. *IOP Conference Series: Earth and Environmental Science*, 967. doi: 10.1088/1755-1315/967/1/012017
- Shibata, M., Hirotsuka, M., Mizutani, Y., Takahashi, H., Kawada, T., Matsumiya, K., Hayashi, Y. & Matsumura, Y. (2017). Isolation and characterization of key contributors to the “kokumi” taste in soybean seeds. *Bioscience, Biotechnology, and Biochemistry*. doi: 10.1080/09168451.2017.1372179
- Shuuluka, D., Bolton, J. J., & Anderson, R. J. (2013). Protein content, amino acid composition and nitrogen-to-protein conversion factors of *Ulva rigida* and *Ulva*

capensis from natural populations and *Ulva lactuca* from an aquaculture system, in South Africa. *Journal of Applied Phycology*, 25: 677-685. doi: 10.1007/s10811-012-9902-5

- Sinurat, E., & Marraskuranto, E. (2011). Fucoidan from brown seaweed and its bioactivity. *SQUALEN Bulletin of Marine and Fisheries Postharvest and Biotechnology*, 7(3): 131-137. doi: 10.15578/squalen.v7i3.8
- Sinurat, E., Peranginangin, R., & Saepudin, E. (2015). Purification and characterization of fucoidan from the brown seaweed *Sargassum binderi* sonder. *SQUALEN Bulletin of Marine & Fisheries Postharvest & Biotechnology*, 10: 79-87. doi: 10.15578/squalen.v10i2.133
- Solms, J. (1969). Taste of amino acids, peptides, and proteins. *Journal of Agricultural and Food Chemistry*, 17: 686-688. doi: 10.1021/jf60164a016
- Spreitzer, R. J., & Salvucci, M. E. (2002). RUBISCO: Structure, regulatory interactions, and possibilities for a better enzyme. *Annual Review of Plant Biology*, 53: 449-475. doi: 10.1146/annurev.arplant.53.100301.135233
- Su, G., Chi, C., Zheng, L., Yang, B., Ren, J. & Zhao, M. (2012). Isolation and identification of two novel umami and umami-enhancing peptides from peanut hydrolysate by consecutive chromatography and MALDI-TOF/TOF MS. *Food Chemistry*, 135(2): 479-485. doi: 10.1016/j.foodchem.2012.04.130
- Su, G., Ren, J., Yang, B., Cui, C., & Zhao, M. (2011). Comparison of hydrolysis characteristics on defatted peanut meal proteins between a protease extract from *Aspergillus oryzae* and commercial proteases. *Food Chemistry*, 126: 1306-1311. doi: 10.1016/j.foodchem.2010.11.083
- Suzuki, H., Nakafuji, Y. & Tamura, T. (2017). New Method to Produce Kokumi Seasoning from Protein Hydrolysates Using Bacterial Enzymes. *Journal of Agricultural and Food Chemistry*, 65(48): 10514-10519. doi: 10.1021/acs.jafc.7b03690
- Tajima, M. & Ito, T. (1999). Change in polypeptides in beef soup stock during simmering. *Proceedings of the 45th International Congress of Meat Science and Technology*, 186-187, Yokohama, Japan.
- Tambun, R., Limbong, H. R., Pinem, C., & Manurung, E. (2017). Pengaruh ukuran partikel, waktu dan suhu pada ekstraksi fenol dari lengkuas merah. *Jurnal Teknik Kimia USU*, 5(4). doi: 10.32734/jtk.v5i4.1555
- Tamura, M., Nakatsuka, T., Tada, M., Kawasaki, Y., Kikuchi, E. & Okai, H. (1989). The relationship between taste and primary structure of "Delicious peptide" (Lys-Gly-Asp-Glu-Glu-Ser-Leu-Ala) from beef soup. *Agricultural and Biological Chemistry*, 53(2): 319-325. doi: 10.1080/00021369.1989.10869317
- Temussi, P. A. (2011). The good taste of peptides. *Journal of Peptide Science*, 18: 73-82. doi: 10.1002/psc.1428

- The UniProt Consortium. (2015). UniProt: a hub for protein information. *Nucleic Acids Research*, 43: D204-D212. doi: 10.1093/nar/gku989
- Toelstede, S. & Hofmann, T. (2009). Kokumi-Active Glutamyl Peptides in Cheeses and Their Biogenesis by *Penicillium roquefortii*. *Journal of Agricultural and Food Chemistry*, 57: 3733-3748. doi: 10.1021/jf900280j
- Tu, M., Cheng, S., Lu, W. & Du, M. (2018). Advancement and prospects of bioinformatics analysis for studying bioactive peptides from food-derived protein: sequence, structure, and functions. *Trends in Analytical Chemistry*. doi: 10.1016/j.trac.2018.04.005
- Udenigwe, C. C. (2014). Bioinformatics approaches, prospects and challenges of food bioactive peptide research. *Trends in Food Science & Technology*, 36: 137-143. doi: 10.1016/j.tifs.2014.02.004
- Udenigwe, C. C., Okolie, C. L., Qian, H., Ohanenye, I. C., Agyei, D., & Aluko, R. E. (2017). Ribulose-1,5-bisphosphate carboxylase as a sustainable and promising plant source of bioactive peptides for food applications. *Trends on Food Science and Technology*, 69: 74-82. doi: 10.1016/j.tifs.2017.09.001
- Ueda, Y., Sakaguchi, M., Hirayama, K., Miyajima, R. & Kimizuka, A. (1990). Characteristic Flavor Constituents in Water Extract of Garlic. *Agriculture and Biological Chemistry*, 54(1): 163-169. doi: 10.1080/00021369.1990.10869909
- Wang, R., Yang, C., & Song, H. (2012). Key meat flavour compounds formation mechanism in a glutathione-xylose Maillard reaction. *Food Chemistry*, 131: 280-285. doi: 10.1016/j.foodchem.2011.08.079
- Wang, W., Cui, Z., Ning, M., Zhou, T., & Liu, Y. (2022). *In-silico* investigation of umami peptides with receptor T1R1/T1R3 for the discovering potential targets: A combined modeling approach. *Biomaterials*, 281. doi: 10.1016/j.biomaterials.2021.121338
- Warkoyo & Saati, E. A. (2011). The solvent effectiveness on extraction process of seaweed pigment. *Makara Journal of Technology*, 15(1): 5-8. doi: 10.7454/mst.v15i1.850
- Weber, K., & Osborn, M. (1969). The reliability of molecular weight determinations by dodecyl sulfate-polyacrylamide gel electrophoresis. *The Journal of Biological Chemistry*, 244(16): 4406-4412. doi: 10.1016/S0021-9258(18)94333-4
- Wu, Q., Cai, Q. F., Yoshida, A., Sun, L. C., Liu, Y. X., Liu, G. M., Su, W. J., & Cao, M. J. (2017). Purification and characterization of two novel angiotensin I-converting enzyme inhibitory peptides derived from R-phycoerythrin of red algae (*Bangia fusco-purpurea*). *European Food Research and Technology*, 243(5), 779-789. doi: 10.1007/s00217-016-2792-z
- Xu, X. D., Xu, R., Song, Z., Jia, Q., Feng, T., Huang, M. G. & Song, S. Q. (2019). Identification of umami-tasting peptides from *Volvariella volvacea* using ultra

- performance liquid chromatography quadrupole time-of-flight mass spectrometry and sensory-guided separation technique. *Journal of Chromatography A*, 1596: 96-103. doi: 10.1016/j.chroma.2019.03.003
- Xu, X., You, M., Song, H., Gong, L. & Pan, W. (2018). Investigation of umami and kokumi taste-active components in bovine bone marrow extract produced during enzymatic hydrolysis and Maillard reaction. *International Journal of Food Science and Technology*. 1-17. doi: 10.1111/ijfs.13893
- Yamasaki, Y. & Maekawa, K. (1978). A peptide with delicious taste. *Agricultural and Biological Chemistry*, 42: 1761-1765. doi: 10.1080/00021369.1978.10863242
- Yan, F., Cui, H., Zhang, Q., Hayat, K., Yu, J., Hussain, S., Tahir, M. U., Zhang, X. & Ho, C. T. (2021). Small Peptides Hydrolyzed from Pea Protein and Their Maillard Reaction Products as Taste Modifiers: Saltiness, Umami, and Kokumi Enhancement. *Food and Bioprocess Technology*, 14: 1132-1141. doi: 10.1007/s11947-021-02630-1
- Yip, W. H., Joe, L. S., Mustapha, W. A. W., Maskat, M. Y., & Said, M. (2014). Characterisation and stability of pigments extracted from *Sargassum binderi* obtained from Semporna, Sabah. *Sains Malaysiana*, 43(9): 134-1354.
- Yu, X., Zhang, L., Miao, X., Li, Y., & Liu, Y. (2017). The structure features of umami hexapeptides for the T1R1/T1R3 receptor. *Food Chemistry*, 221(15): 599-605. doi: 10.1016/j.foodchem.2016.11.133
- Yudiati, E. & Isnansetyo, A. (2017). Characterizing the three different alginate type of *Sargassum siliquosum*. *Ilmu Kelautan*, 22(1): 7-14. doi: 10.14710/ik.ijms.22.1.7-14.
- Zhang, M., Wang, X., Liu, Y., Xu, X., & Zhou, G. (2012). Isolation and identification of flavour peptides from Puffer fish (*Takifugu obscurus*) muscle using an electronic tongue and MALDI-TOF/TOF MS/MS. *Food Chemistry*, 135(3), 1463-1470. doi: 10.1016/j.foodchem.2012.06.026
- Zhang, Y., Gao, X., Pan, D., Zhang, Z., Zhou, T., & Dang, Y. (2021). Isolation, characterization and molecular docking of novel umami dan umami-enhancing peptides from *Ruditapes philippinarum*. 343(1). doi: 10.1016/j.foodchem.2020.128522
- Zhang, Y., Ma, Y., Ahmed, Z., Geng, W., Tang, W., Liu, Y., Jin, H., Jiang, F., Wang, J. & Wang, Y. (2019a). Purification and identification of kokumi-enhancing peptides from chicken protein hydrolysate. *International Journal of Food Science and Technology*. doi: 10.1111/ijfs.14122
- Zhang, Y., Venkitasamy, C., Pan, Z., Liu, W. & Zhao, L. (2017). Novel Umami Ingredients: Umami Peptides and Their Taste. *Journal of Food Science*, 82(1): 16-23. doi: 10.1111/1750-3841.13576

- Zhang, J., Zhao, M., Su, G., & Lin, L. (2019b). Identification and taste characteristics of novel umami and umami-enhancing peptides separated from peanut protein isolate hydrolysate consecutive chromatography and UPLC-ESI-QTOF-MS/MS. *Food Chemistry*, 278, 674-682. doi: 10.1016/j.foodchem.2018.11.114
- Zhao, W., He, J., Yu, Z., Wu, S., Li, J., Liu, J., & Liao, X. (2022). *In silico* identification of novel small molecule umami peptide from ovotransferin. *International Journal of Food Science Technology*. doi: 10.1111/IJFS.15166
- Zhao, Y., Zhang, M., Devahastin, S., & Liu, Y. (2019). Progresses on processing methods of umami substances: A review. *Trends in Food Science and Technology*, 93: 125–135. doi: 10.1016/j.tifs.2019.09.012.
- Zhu, X., Sun-Waterhouse, D., Chen, J., Cui, C. & Wang, W. (2020). Comparative study on the novel umami-active peptides of the whole soybeans and the defatted soybeans fermented soy sauce. *Journal of the Science of Food and Agriculture*. doi: 10.1002/jsfa.10626
- Zhuang, M., Lin, L., Zhao, M., Dong, Y., Sun-Waterhouse, D., Chen, H., Qiu, C., & Su, G. (2016). Sequence, taste and umami-enhancing effect of the peptides separated from soy sauce. *Food Chemistry*, 206, 174-181. doi: 10.1016/j.foodchem.2016.03.058