

2014 Bag disertasi_Sayur asin_EFRT_Oct 2014 Q1.pdf

 Unika Soegijapranata1

Document Details

Submission ID

trn:oid:::28973:124071425

23 Pages

Submission Date

Dec 9, 2025, 12:01 PM GMT+7

7,294 Words

Download Date

Dec 9, 2025, 12:13 PM GMT+7

38,116 Characters

File Name

2014 Bag disertasi_Sayur asin_EFRT_Oct 2014 Q1.pdf

File Size

967.5 KB

14% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

Filtered from the Report

- ▶ Bibliography
- ▶ Quoted Text
- ▶ Cited Text
- ▶ Small Matches (less than 10 words)

Exclusions

- ▶ 84 Excluded Sources
- ▶ 5 Excluded Matches

Match Groups

-  **57** Not Cited or Quoted 14%
Matches with neither in-text citation nor quotation marks
-  **0** Missing Quotations 0%
Matches that are still very similar to source material
-  **0** Missing Citation 0%
Matches that have quotation marks, but no in-text citation
-  **0** Cited and Quoted 0%
Matches with in-text citation present, but no quotation marks

Top Sources

11%	 Internet sources
13%	 Publications
6%	 Submitted works (Student Papers)

Integrity Flags

0 Integrity Flags for Review

No suspicious text manipulations found.

Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

Match Groups

-  57 Not Cited or Quoted 14%
Matches with neither in-text citation nor quotation marks
-  0 Missing Quotations 0%
Matches that are still very similar to source material
-  0 Missing Citation 0%
Matches that have quotation marks, but no in-text citation
-  0 Cited and Quoted 0%
Matches with in-text citation present, but no quotation marks

Top Sources

- 11%  Internet sources
- 13%  Publications
- 6%  Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

Rank	Source Type	Source	Percentage
1	Publication	Sumathy Babitha, Carlos Ricardo Soccol, Ashok Pandey. "Effect of stress on growth and yield of mustard seed (Sinapis alba L.) under different sowing dates and irrigation regimes." <i>Journal of Agricultural Science</i> 145, no. 1 (2007): 1-10.	<1%
2	Internet	www.ddsis.lv	<1%
3	Internet	www.freepatentsonline.com	<1%
4	Publication	Franck Doussot, Bernard De Jéso, Stéphane Quideau, Patrick Pardon. "Extractives and phenolic compounds in mustard seed (Sinapis alba L.) under different sowing dates and irrigation regimes." <i>Journal of Agricultural Science</i> 145, no. 1 (2007): 1-10.	<1%
5	Internet	ndl.ethernet.edu.et	<1%
6	Publication	Kelly A Meckling, Melanie Gauthier, Rebecca Grubb, Jennifer Sanford. "Effects of a high protein diet on the performance of broiler chickens." <i>Journal of Animal Science</i> 63, no. 1 (1995): 1-10.	<1%
7	Submitted works	National University of Singapore on 2023-02-25	<1%
8	Publication	Paweł Górnawski, Inga Miśina, Silvija Ruisa, Edgars Rubauskis, Gunārs Lācis, Dalija Sečka, Agnese Čeplāne, Jānis Čeplāns. "Effect of different sowing dates and irrigation regimes on the yield and quality of mustard seed (Sinapis alba L.)." <i>Journal of Agricultural Science</i> 145, no. 1 (2007): 1-10.	<1%
9	Submitted works	Canterbury Christ Church University on 2022-05-04	<1%
10	Publication	Van Eylen, D.. "Temperature and pressure stability of mustard seed (Sinapis alba L.)." <i>Journal of Agricultural Science</i> 145, no. 1 (2007): 1-10.	<1%

11 Publication

Theresa Walter, Herbert Wieser, Peter Koehler. "Degradation of gluten in rye sou... <1%

12 Publication

Hanne Christine Bertram, Lotte Bach Larsen, Xiaoping Chen, Per Bendix Jeppesen... <1%

13 Submitted works

Nottingham Trent University on 2021-08-30 <1%

14 Publication

Vesna Katavic, Ganesh Kumar Agrawal, Martin Hajduch, Stefan L. Harris, Jay J. Th... <1%

15 Internet

ds.inflibnet.ac.in <1%

16 Internet

www2.mdpi.com <1%

17 Publication

Timo D. Stark, Sofie Lösch, Oliver Frank, Onesmo B. Balemba, Thomas Hofmann. "... <1%

18 Internet

portalcientifico.ucavila.es <1%

19 Internet

ruidera.uclm.es <1%

20 Internet

www.cambridge.org <1%

21 Publication

Oliviero, T.. "Effect of water content and temperature on glucosinolate degradati... <1%

22 Submitted works

Texas A&M University, College Station on 2020-09-09 <1%

23 Internet

coek.info <1%

24 Publication

de Oliveira, Maria João Simões Cardoso. "A Comprehensive Study on Surveillance ... <1%

25	Internet	
harvest.usask.ca		<1%
26	Publication	
Materska, Małgorzata. "Flavone C-glycosides from Capsicum annuum L.: relations...		<1%
27	Internet	
www.agriculturejournals.cz		<1%
28	Internet	
www.awri.com.au		<1%
29	Internet	
zaguan.unizar.es		<1%
30	Publication	
H. Hajjaj, P. J. Blanc, E. Groussac, G. Goma, J. L. Uribelarrea, P. Loubiere. "Improve...		<1%
31	Internet	
d-nb.info		<1%
32	Internet	
journals.uni-lj.si		<1%
33	Internet	
www.coursehero.com		<1%
34	Publication	
Mutamed Ayyash, Abdelmoneim Abdalla, Basim Abu-Jdayil, Thom Huppertz et al. ...		<1%
35	Internet	
amsdottorato.unibo.it		<1%
36	Internet	
efsa.onlinelibrary.wiley.com		<1%
37	Internet	
ftb.com.hr		<1%
38	Internet	
ithuteng.ub.bw		<1%

39	Internet	
orbi.uliege.be	<1%	
40	Internet	
repositorio.uchile.cl	<1%	
41	Internet	
theses.hal.science	<1%	
42	Internet	
core-cms.prod.aop.cambridge.org	<1%	
43	Internet	
www.nature.com	<1%	
44	Internet	
www.trinityuniversity.edu.ng	<1%	
45	Internet	
datapdf.com	<1%	
46	Internet	
digitalcommons.lsu.edu	<1%	
47	Internet	
www.ots.ac.cr	<1%	
48	Publication	
Ahmed, Umbreen, and Phillip S. Oates. "Dietary fat level affects tissue iron levels ...	<1%	
49	Publication	
Free-Radical-Induced DNA Damage and Its Repair, 2006.	<1%	
50	Publication	
Julia Szutowska, Daniela Gwiazdowska, Iga Rybicka, Katarzyna Pawlak-Lemańska ...	<1%	
51	Publication	
Luigi Michele Sena, Claudia Zappelli, Fabio Apone, Ani Barbulova et al. " hairy roo...	<1%	
52	Publication	
Shah, N.N., and R.S. Singhal. "Fermented Fruits and Vegetables", Current Develop...	<1%	

53

Submitted works

Wageningen University on 2019-01-25 <1%

54

Publication

X. Jin, T. Oliviero, R.G.M. van der Sman, R. Verkerk, M. Dekker, A.J.B. van Boxtel. "I... <1%

55

Internet

eis.hu.edu.jo <1%

56

Internet

eprints.undip.ac.id <1%



European Food Research & Technology

Editor-in-Chief: Werner Tatzusch



Springer

[Skip to main content](#) Search 


- [Log in](#)

[Search SpringerLink](#) Search 

You're seeing our new journal sites and we'd like your opinion, please [send feedback](#)

[European Food Research and Technology](#)

- [Journal home](#) >
- [Volumes and issues](#) >
- [Volume 240, issue 3](#)

[Search within journal](#) Search 



Volume 240, issue 3, March 2015


23 articles in this issue

27 1. Effects of environmental factors on functional properties of Chinese chestnut (*Castanea mollissima*) protein isolates

Authors (first, second and last of 5)

- o Jiao Lv

- Fengjun Wang
- Content type: Review Paper
- Published: 26 October 2014



2. Nutritional properties of different composite flours from maize (*Zea mays*) variety ica v109 and pearl millet (*Pennisetum glaucum*) malted with calcium chloride and gibberellic acid


Authors

- Guillermo Balseiro
- Arnulfo Taron
- Luis A. Garcia-Zapateiro
- Content type: Original Paper
- Published: 10 October 2014

3. About the origin of asclepic acid derived from crude homo- and heterolipids during successive solvent extraction of rapeseeds

Authors (first, second and last of 9)


- Jan Kyselka
- Frantisek Kreps
- Vladimir Filip
- Content type: Original Paper
- Published: 04 October 2014



4. Nuclear magnetic resonance-based metabolomics reveals that dairy protein fractions affect urinary urea excretion differently in overweight adolescents

Authors (first, second and last of 8)

- Hanne C. Bertram
- Content type: Original Paper
- Published: 01 October 2014




29

5. Accelerate and enhance the release of haze-protective polysaccharides after alcoholic fermentation in winemaking

Authors (first, second and last of 5)

- Cristiane Moriwaki
- Gracielle Matioli
- Ana Briones
- Content type: Original Paper
- Published: 04 October 2014



16

6. Identification and characterization of new allergen troponin C (Pen m 6.0101) from Indian black tiger shrimp *Penaeus monodon*

Authors


- Aravindan Kalyanasundaram
- Thairiyam Chinnappan Santiago
- Content type: Original Paper
- Published: 12 October 2014

11

7. Degradation of gluten in rye sourdough products by means of a proline-specific peptidase

Authors

- Theresa Walter
- Herbert Wieser
- Peter Koehler
- Content type: Original Paper
- Published: 04 October 2014



18. **Analysis of grapes and the first stages of the vinification process in wine contamination with *Brettanomyces bruxellensis***


Authors (first, second and last of 7)

- P. Garijo
- L. González-Arenzana
- A. R. Gutiérrez
- Content type: Original Paper
- Published: 02 October 2014

9. **Chemical and sensory evaluation of wine matured in oak barrel: effect of oak species involved and toasting process**

Authors


- Kleopatra Chira
- Pierre-Louis Teissedre
- Content type: Original Paper
- Published: 16 October 2014



26. **Flavone C-glycosides from *Capsicum annuum* L.: relationships between antioxidant activity and lipophilicity**

Authors

- Małgorzata Materska
- Content type: Original Paper
- Open Access



20 11. **Retention of glucosinolates during fermentation of *Brassica juncea*: a case study on production of *sayur asin***


Authors (first, second and last of 5)

- Probo Y. Nugrahedi
- Budi Widianarko
- Teresa Oliviero
- Content type: Original Paper
- Published: 17 October 2014

36 12. **Spectroscopic detection of aspartame in soft drinks by surface-enhanced Raman spectroscopy**

Authors (first, second and last of 5)


- Guluzar Gorkem Buyukgoz
- Akif Goktug Bozkurt
- Ismail Hakki Boyaci
- Content type: Orginal Paper
- Published: 14 October 2014



30 13. **Effect of low-frequency magnetic field on formation of pigments of *Monascus purpureus***

Authors (first, second and last of 4)


- Jialan Zhang
- Dongjie Zeng
- Mengxiang Gao
- Content type: Orginal Paper
- Published: 17 October 2014



6 14. **Impact of a 6-week very low-calorie diet and weight reduction on the serum and fecal metabolome of overweight subjects**

Authors (first, second and last of 6)


- Mette S. Schmedes
- Christian C. Yde
- Hanne C. Bertram
- Content type: Original Paper
- Published: 26 October 2014



19 15. **Phenolic characterization of minor red grape varieties grown in Castilla-La Mancha region in different vinification stages**

Authors (first, second and last of 5)

- María Jesús Cejudo-Bastante
- Almudena Vicario
- María Soledad Pérez-Coello
- Content type: Original Paper
- Published: 31 October 2014



28 16. **Aromatic compounds released from natural precursors by selected *Oenococcus oeni* strains during malolactic fermentation**

Authors (first, second and last of 5)

- Fátima Pérez-Martín
- Pedro Miguel Izquierdo-Cañas
- María Llanos Palop

Content type: Original Paper


o Published: 31 October 2014

23

17. **A new sensitive method for the detection of chloramphenicol in food using time-resolved fluoroimmunoassay**

Authors (first, second and last of 7)

- o Bin Zhou
- o Jue Zhang
- o Biao Huang
- o Content type: Original Paper
- o Published: 07 November 2014




32

18. **Effects of different substrates and oils on aflatoxin B₁ production by *Aspergillus parasiticus***

Authors (first, second and last of 4)

- o Huihui Ma
- o Niya Zhang
- o Desheng Qi
- o Content type: Original Paper
- o Published: 29 October 2014




1

19. **Effects of glycerol on pigments and monacolin K production by the high-monacolin K-producing but citrinin-free strain, *Monascus pilosus* MS-1**

Authors (first, second and last of 4)


- o Yanli Feng
- o Yanchun Shao
- o Fusheng Chen



40 20. **Reactivity of pure and commercial grape skin tannins with cell wall material**

Authors (first, second and last of 5)


- Ana Belén Bautista-Ortín
- Noelia Molero
- Encarna Gómez-Plaza
- Content type: Orginal Paper
- Published: 26 October 2014



31 21. **Influence of gel from ground chia (*Salvia hispanica L.*) for wheat bread production**

Authors (first, second and last of 4)

- V. Zettel
- A. Krämer
- B. Hitzmann
- Content type: Orginal Paper
- Published: 28 October 2014



22. **Composition of tocopherol in kernels recovered from different sweet cherry (*Prunus avium L.*) cultivars: RP-HPLC/FLD and RP-UPLC-ESI/MSⁿ study**

Authors (first, second and last of 6)

- Dalija Seglića
- Content type: Short Communication
- Published: 12 November 2014

Vortex \times 10 sec, after 10 min of boiling
 Cool immediately in an ice-water bath
 Hold there for the whole time of extraction
 Add 2.5 ml NaCl (10 g/L) and vortex \times 5 sec.
 Add 2.5 ml of *n*-hexane:ethyl acetate (9:1; v/v)
 Vortex \times 15 sec, and centrifuge \times 5 min (1000 \times g, at 4 °C)
 Remove organic layer to the round bottom flask
 Repeat *n*-hexane:ethyl acetate extraction \times 3

7 23. **Erratum to: Effect of different halophilic microbial fermentation patterns on the volatile compound profiles and sensory properties of soy sauce moromi**

Authors (first, second and last of 4)

- Rui-Ying Cui
- Jia Zheng
- Rong-Qing Zhou
- Content type: Erratum
- Published: 20 January 2015

13 **For authors**

[Submission guidelines](#) [Ethics & disclosures](#) [Contact the journal](#) [Submit manuscript](#)

Explore

[Online first articles](#) [Volumes and issues](#) [Sign up for alerts](#)

22 Over 10 million scientific documents at your fingertips

Switch Edition

- [Academic Edition](#)
- [Corporate Edition](#)
- [Home](#)
- [Impressum](#)
- [Legal information](#)
- [Privacy statement](#)
- [California Privacy Statement](#)
- [How we use cookies](#)
- [Manage cookies/Do not sell my data](#)

- [Contact us](#)

Not logged in - 36.80.117.169

Universitas Katolik Soegijapranata (3991461827)

[Springer Nature](#) SPRINGER NATURE

© 2020 Springer Nature Switzerland AG. Part of [Springer Nature](#).

33

[Skip to main content](#)[Search](#)

- [Authors & Editors](#)
- [My account](#)

[Menu](#)

- [Authors & Editors](#)
- [My account](#)

[European Food Research and Technology](#)

- [Journal home](#) >
- Editors

Editors

Editors-in-Chief

T. Henle

Technische Universität Dresden
Institut für Lebensmittelchemie
01062 Dresden Germany

17

T. Hofmann

Lehrstuhl für Lebensmittelchemie und Molekulare Sensorik

Technische Universität München

Lise-Meitner-Str. 34

85354 Freising

Germany



E. Anklam, Geel, Belgium
A. Arnoldi, Milan, Italy
P. S. Belton, Norwich, UK
M. Defernez, Norwich, UK
R. Einspanier, Berlin, Germany
J. Empis, Lisbon, Portugal
R.J. Fritsch, Glenview, USA
T. Haertle, Nantes, France
H. Kallio, Turku, Finland
M. Kovac, Bratislava, Slovak Republic
J. Löliger, Kemptthal, Switzerland
M. Petz, Wuppertal, Germany
J.S. Rader, Washington DC, USA
M. Ramos, Madrid, Spain
S. Saguy, Rehovot, Israel
P. Sandra, Ghent, Belgium
P. Schieberle, Garching, Germany
P. Schreier, Würzburg, Germany
L. Simon-Sarkady, Budapest, Hungary
L. Skibsted, Frederiksberg C, Denmark
A.J. Taylor, Loughborough, UK
F.A. Tomás-Barberán, Murcia, Spain
J. Velisek, Prague, Czech Republic
A. Visconti, Bari, Italy
J. Wilska-Jeszka, Lodz, Poland

For authors

9 [Submission guidelines](#) [Ethics & disclosures](#) [Contact the journal](#) [Submit manuscript](#)

Explore

[Online first articles](#) [Volumes and issues](#) [Sign up for alerts](#)



Publish with us

[Authors & Editors](#)
[Journal authors](#)
[Publishing ethics](#)
[Open Access & Springer](#)

[Books A-Z](#)
[Journals A-Z](#)
[Video](#)

Other services

[Instructors](#)
[Librarians \(Springer Nature\)](#)
[Societies and Publishing Partners](#)
[Advertisers](#)
[Shop on Springer.com](#)

About Springer

[About us](#)
[Help & Support](#)
[Contact us](#)
[Press releases](#)
[Impressum](#)

Legal

[General term & conditions](#)
[California Privacy Statement](#)
[Rights & permissions](#)
[Privacy](#)
[How we use cookies](#)
[Manage cookies/Do not sell my data](#)
[Accessibility](#)

Not logged in - 36.80.117.169

Not affiliated

SPRINGER NATURE

Retention of glucosinolates during fermentation of *Brassica juncea*: a case study on production of *sayur asin*

Probo Y. Nugrahedi · Budi Widianarko ·
Matthijs Dekker · Ruud Verkerk · Teresa Oliviero

Received: 18 June 2014 / Revised: 9 September 2014 / Accepted: 5 October 2014
© Springer-Verlag Berlin Heidelberg 2014

Abstract Fermentation can reduce the concentration of health-promoting glucosinolates in *Brassica* vegetables. The endogenous enzyme myrosinase is hypothesised to mainly responsible for the degradation of glucosinolates during fermentation. In order to retain glucosinolates in the final fermented product, the role of myrosinase activity during the production of *sayur asin* was investigated. *Sayur asin* is a traditionally fermented product of Indian mustard (*Brassica juncea*) commonly consumed in Indonesia. It is prepared by a spontaneous fermentation of withered (sundried) *B. juncea* leaves. The leaves of *B. juncea* contain a substantial amount of the aliphatic glucosinolate sinigrin. Three withering methods were investigated to obtain *B. juncea* leaves with different myrosinase activities prior to fermentation. Results show that withering by oven at 35 °C for 2.5 h and by microwave at 180 W for 4.5 min reduced myrosinase activity by 84 and 74 %, respectively. Subsequently, sinigrin was not detectable in the leaves after 24 h of incubation in the fermentation medium. However, withering by microwave for 2 min at 900 W inactivated myrosinase completely and produced *sayur asin* with a sinigrin concentration of 11.4 µmol/10 g dry matter after 7 days of fermentation. This high power-short time pretreatment

of *B. juncea* leaves contributes to the production of *sayur asin* containing significant levels of health-promoting glucosinolate. In this study, the effect of myrosinase activity during *Brassica* fermentation was quantified, and optimised production methods were investigated to retain glucosinolate in the final product.

Keywords Glucosinolate · Myrosinase activity · Fermentation · *Brassica juncea*

Introduction

Glucosinolates (GLSs) are a group of β -thioglucoside *N*-hydroxysulphates with a sulphur-linked β -D-glucopyranose moiety and side-chain group. GLSs are secondary metabolites derived mainly from one of certain amino acids, such as methionine, tryptophan, and phenylalanine. Depending on side-chain group, glucosinolate (GLS) can be classified either as aliphatic, aromatic, or indolyl [1–3].

Reviews of epidemiological studies reported inverse associations between intake of *Brassica* vegetables and the risk of certain cancers [4, 5]. Isothiocyanates, one of the GLS breakdown products, are assumed to lower the risk of cancer by inhibiting phase 1 and inducing phase 2 enzymes during carcinogen metabolism [6]. Upon damage of plant tissue, GLSs are highly prone to hydrolysis catalysed by myrosinase (β -thioglucosidase EC 3.2.1.147), an enzyme that occurs in *Brassica* vegetables [2, 7].

Fermentation is a processing method commonly performed on milks, meats, and vegetables [8]. Fermentation of *Brassica* vegetables involves the growth and metabolic activity of lactic acid bacteria (LAB), either spontaneously or starter-induced to produce fermented products [9]. Previous studies reported that fermentation considerably reduces

P. Y. Nugrahedi · M. Dekker · R. Verkerk (✉) · T. Oliviero
Food Quality and Design Group, Department of Agrotechnology
and Food Science, Wageningen University, PO Box 17,
6700 AA Wageningen, The Netherlands
e-mail: ruud.verkerk@wur.nl

P. Y. Nugrahedi
e-mail: probo.nugrahedi@wur.nl

P. Y. Nugrahedi · B. Widianarko
Department of Food Technology, Soegijapranata Catholic
University, Jl. Prawiyatno Luhur IV/1, Semarang 50234, Indonesia

the concentration of GLSs in *sauerkraut* [10, 11] and radish *kimchi* [12].

The mechanisms underlying GLS changes in *Brassica* vegetables during fermentation are less investigated than other processing methods, such as boiling, steaming, and microwave (MW) processing [13, 14]. Several factors can influence the loss of GLS during fermentation, such as the myrosinase activity, LAB growth and activity, low pH, and sodium chloride concentration [9, 11, 15]. *Sayur asin* is a traditional fermented product of Indian mustard (*Brassica juncea*) commonly produced in some regions in Indonesia [16] and other Asian countries, in different local names [17]. The product is prepared by a spontaneous fermentation for 7 days of withered (sun-dried) Indian mustard leaves in the addition of salt followed by rubbing [16]. Either starch water of boiling rice or coconut water can be used as the fermentation medium. In the preliminary study, it was shown that during the production of *sayur asin* sinigrin concentration was reduced to 1.3 % after 3 days of fermentation [18]. The loss of GLS is probably attributed to hydrolytic activity of myrosinase. The myrosinase activity is mainly ascribed to the endogenous myrosinase present in the leaves, although microorganisms can also show a myrosinase-like activity [19, 20].

This study aims to retain the GLS in Indian mustard during the *sayur asin* production and to study how the myrosinase activity affects the GLS concentration during the fermentation. For this purpose, three withering methods were applied to produce Indian mustard leaves with different myrosinase activities prior to fermentation, namely by oven at 35 °C and by MW at 180 and 900 W. Upon withering, the samples were fermented for 7 days, and the GLS concentration and the myrosinase activity were monitored. Previous studies reported the diversity of microbes in traditionally fermented *sayur asin* [16] and mustard products [21]. Since the present study modifies the production method of *sayur asin*, the microbiological profile of *sayur asin* was also investigated.

Materials and methods

Preparation of inoculum and fermentation medium for *sayur asin*

Since the withering treatments will greatly reduce the number of natural LAB on the leaves, a standard inoculum was prepared for all three treatments, by taking an aliquot of coconut water of a 5 days fermented *sayur asin* previously withered in the oven at 35 °C. The aliquot was inoculated in the coconut water (Omega & More, Witsenburg Natural Products BV) at 1 % (v/v) to achieve about similar number of total LAB that is reached at the beginning

of fermentation of a traditionally withered *sayur asin* (i.e. approximately 6 log CFU/ml).

Preparation of *sayur asin*

Raw Indian mustard was obtained from a local supplier (Oriental Webshop, Duiven, The Netherlands). Indian mustard leaves were separated, washed, drained and randomly divided into portions of 200 g (3–4 leaves). There were three withering methods performed, i.e. by an oven at 35 °C for 2.5 h to mimic the traditional process, by MW at a power of 180 W for 4.5 min to reduce the myrosinase activity, and by MW at 900 W for 2 min to inactivate the myrosinase activity. Hence, the withered leaves containing different levels of myrosinase activity were obtained. Then, salt was added to the withered vegetables (2.5 %, w/w) and rubbed gently by hand, similar to the traditional process. Subsequently, the leaves were immersed in coconut water containing 1 % (v/v) of the standard inoculum, at a ratio of 1:5 (w/v). Each product was packed in 360-mL glass jar and incubated in the dark at a room temperature of 22 °C for 7 days. Each treatment was carried out in triplicate.

Sample preparation

Half of the amount of the vegetable and of the medium in the jar were taken aseptically followed by a quick freezing in liquid nitrogen, while the remaining vegetable and medium of the sample was mixed thoroughly with a stomacher for further microbiological analysis. The raw, withered, and fermented vegetables as well as the fermentation medium were frozen in liquid nitrogen followed by lyophilisation. Subsequently, the vegetables were grinded and stored at –22 °C until further analyses of GLS and myrosinase activity. These analyses were also performed to the fermentation medium.

Myrosinase activity analysis

The myrosinase activity was determined by a coupled enzymatic procedure followed by spectrophotometry measurement [22], in which the produced glucose, from the reaction of native myrosinase and a standard GLS, was measured. A 50 mg lyophilised sample was mixed overnight in 140 mL of 50 mM potassium phosphate buffer (pH 7.0) at 15 °C. On the second day, the plant matrix was removed from the myrosinase solution by centrifugation at 2,670g for 10 min followed by clarification (folded filters Grade 595 1/2–4–7 µm, Whatman). Subsequently, 2 mL of this solution was filtered (cut-off 30 kD Amicon Ultra-4 Millipore) from the molecules smaller than myrosinase with the aid of centrifugation at 4,000g for 10 min. The concentrated enzyme was dissolved in 500 µL of potassium phosphate buffer. Eventually, the myrosinase activity was measured by a D-glucose enzyme

42
54
10

43

25

37

51

50

53

56

kit (Enzyplus EZS 781+). The reaction mixture consisted of a water solution containing 0.05 g/L of magnesium chloride and 1 g/L ascorbic acid, a ATP/NADP⁺ solution, buffer solution, hexokinase/glucose-6-P-dehydrogenase solution and 50 µL sample solution containing the myrosinase, and sinigrin 30 mg/mL solution. The formation of NADPH was followed by a spectrophotometer (Cary UV 50) at 340 nm for 7 min. To quantify myrosinase activity, an external calibration was determined by following the same procedure for the samples analysis. In the reaction mixture of 50 µL, a standard myrosinase solution instead of the myrosinase sample extract was added. Activity was expressed as unit/g dry material (DM) where one unit produces 1.0 µmol glucose per minute from sinigrin at pH 6.0 at 25 °C.

Glucosinolate analysis

Extraction of GLSs with hot methanol (70 %), containing an internal standard of glucotropaeolin, followed by on-column desulphation and analysed by high-performance liquid chromatography was done in duplicate according to the procedure described by Sarvan et al. [15]. The de-sulpho-GLSs were separated by using the Lichrospher100 column (Merck RP-18, 5 µm) with an attached LiChroCART guard column (RP-18, 4 × 4 mm) (Merck, Darmstadt) at a flow rate of 1 mL/min. The elution was performed by the gradient system of water and acetonitrile. Detection was performed with a DAD detector (Spectra System UV 6000 LP) at a wavelength of 229 nm. The GLSs were identified by the spectra as well as comparing with the external standard of sinigrin and internal standard of glucotropaeolin.

Microbiological analyses

Samples of the fermenting liquid and vegetables were collected from the jars for counting of total bacteria, LAB, yeasts, and *Bacillus cereus* by the standard spread plate method. Total colony counts, LAB, *B. cereus*, and yeasts were counted on Plate Count agar (PCA, Oxoid), deMan-Rogosa-Sharpe agar (MRSA, Merck), Mannitol Yolk Polymyxin agar base (MYP, Merck) plus egg yolk emulsion and *B. cereus* selective supplement, and Oxytetracycline Glucose Yeast Extract agar (OGYE, Oxoid), respectively. Plates of PCA, MYP, and OGYE were aerobically incubated at 35 °C for 2 days, at 32 °C for 1 day, and at 25 °C for 5 days, respectively. Plates of MRS were anaerobically incubated at 35 °C for 3 days. The colonies that developed on each plate were counted.

Statistical analysis

Effects of withering methods, namely the oven, MW 180, and MW 900, on GLS concentration, total LAB, and total colony count during the production of *sayur asin* were

analysed by one-way ANOVA and Duncan's multiple range test at the 5 % significance level (IBM SPSS Statistics version 20, U.S.). Each sample was produced in triplicate, and the GLS analysis was performed in duplicate.

Results and discussion

Myrosinase activity

The effect of withering methods on myrosinase activity

Raw leaves of Indian mustard showed a myrosinase activity of 162 ± 56 units/g DM. The effect of the three withering methods on myrosinase activity in the leaves can be seen in the Table 1. Withering by oven at 35 °C and by MW at 180 W retained the myrosinase activity at 16 and 26 %, respectively. While, after a MW treatment at 900 W no myrosinase activity was detected in the leaves. Accordingly, various studies reported that microwaving is an efficient treatment for partial or complete inactivation of myrosinase in *Brassicas*, by varying the time–temperature and power [23–25]. The MW conditions and amount and type of vegetables determine largely the degree of enzyme inactivation. Moreover, thermal inactivation of myrosinase depends also on the plant source, with reported inactivation temperatures in the range from 30 to 60 °C [26, 27].

Myrosinase activity during the fermentation

The present study shows that the remaining myrosinase activity in the oven and MW at 180 W treated samples was almost completely lost after one day of fermentation (Table 1). Further fermentation led to complete loss of myrosinase activity in all samples. Myrosinase activity was also not detected in the fermentation medium (data not shown). In accordance, Kim and Rhee [12] reported a decrease of myrosinase activity from the radish tissue after 3 days of fermentation until less than 3 %. They explained this by reduced stability due

Table 1 Myrosinase activity (units/g DM) of withered and fermented Indian mustard leaves (*n* = 3)

	Withering treatment		
	Oven	MW 180	MW 900
After withering	26.5 ± 1.9	41.9 ± 4.7	n.d.
Fermentation day-1	0.043 ± 0.004	0.109 ± 0.018	n.d.
Fermentation day-3–7	n.d.	n.d.	n.d.

Oven, MW 180, and MW 900 = oven at 35 °C for 2.5 h, MW at 180 W for 4.5 min, and at 900 W for 2 min, respectively

n.d. not detectable

Values are presented as mean ± SD

Table 2 Glucosinolate concentration ($\mu\text{mol}/10\text{ g DM}$) in Indian mustard leaves treated by three withering methods ($n = 6$)

Glucosinolate	Raw leaves	Withering treatment		
		Oven	MW 180	MW 900
Sinigrin	178.9 ± 51.6^c	53.3 ± 19.0^a	85.1 ± 8.2^b	86.0 ± 21.2^b
4-Hydroxy-glucobrassicin	1.1 ± 0.6^a	$1.0 \pm 0.4^a (n = 5)$	1.3 ± 0.5^a	1.5 ± 1.3^a
Glucobrassicin	1.4 ± 0.6^b	$0.5 \pm 0.4^a (n = 5)$	$0.4 \pm 0.08^a (n = 5)$	0.5 ± 0.3^a
4-Methoxy-glucobrassicin	3.3 ± 0.9^c	1.1 ± 0.4^a	2.3 ± 0.4^b	1.9 ± 0.9^b
Total glucosinolates	184.7	55.9	89.1	89.9

Oven, MW 180, and MW 900 = oven at 35 °C for 2.5 h, MW at 180 W for 4.5 min, and at 900 W for 2 min, respectively

Mean reported of duplicate analyses from 3 treatment replications

Values are presented as mean \pm SD

Values with superscript letters on the same row, which are different are significantly different ($P < 0.05$)

to the decrease of pH and the possible production of microbial proteolytic enzyme during fermentation. The initial pH of the fermentation medium in the present study was 4.6 and decreased after 1 day and during fermentation of *sayur asin* of all treatments to the range of 3.3–3.7. Moreover, the high water activity of the *sayur asin* may affect the myrosinase stability. A previous study reported that the inactivation rate of myrosinase in broccoli increases with the increase of water activity of the sample [28].

Glucosinolates concentration

Glucosinolates in raw Indian mustard

Sinigrin was found to be the predominant aliphatic GLS in Indian mustard accounting for more than 95 % of the total GLSs with a concentration of $179 \pm 52 \mu\text{mol}/10\text{ g DM}$ (Table 2). Many studies on the corresponding isothiocyanate of the sinigrin, the allyl isothiocyanate, show that the isothiocyanate inhibits proliferation of human prostate and bladder cancer cells [29, 30].

Three indolyl GLSs, i.e. 4-hydroxy-glucobrassicin, glucobrassicin, and 4-methoxy-glucobrassicin, were detected in the Indian mustard leaves in total accounting for about $5.8 \mu\text{mol}/10\text{ g DM}$. In general, the profile of the main GLSs is in accordance with previous studies on Indian mustard [31–33]. However, other minor GLSs, such as progoitrin, glucobrassicinapin, gluconapin, neoglucoibrassicin, and gluconasturtiin, have been reported [33, 34]. Differences in the GLS concentration and profile could be due to the variations in cultivars, growing conditions, environmental factors, and the analytical methods [2, 11, 33–35].

The effect of withering methods on glucosinolates

Withering Indian mustard leaves caused a considerable reduction of total GLSs ranging from about 50 % by MW

treatments to 70 % by oven treatment (Table 2). Decrease of GLSs during withering could be caused by thermal degradation and myrosinase hydrolysis. Previous studies reported a lower GLS loss upon MW treatment on cabbage [23, 25] than the present study. The different size and shape as well as structural matrix between Indian mustard and cabbage could have led to a different loss of GLSs in Indian mustard. Moreover, the damage of the cell membranes in the leaves containing active myrosinase, followed by diffusion, may have allowed myrosinase to hydrolyse the GLSs, which can occur at a different rate for different vegetables.

Glucosinolates during fermentation

After the first day of fermentation, there was a marked drop of total GLSs in all samples (Table 3). The MW at 180 W treated sample led to the greatest loss of up to 95 %, followed by up to 90 % for the oven-treated sample, and up to 70 % for the MW at 900 W treated sample, relative to the total GLS concentration upon withering. Furthermore, GLSs were not detectable after the third day of fermentation in the *sayur asin* previously withered in the oven and in the MW at 180 W. The withering by MW at 900 W resulted in a retention of sinigrin of 13 % upon fermentation when compared to the concentration after withering (Table 3), while negligible amount of sinigrin was found in the fermentation medium (data not shown).

Accordingly, previous studies reported that fermentation can significantly reduce the amount of GLSs in vegetables [9–11, 36]. During *sauerkraut* production, Tolonen et al. [9] observed a very low quantity of 4-methoxy-glucobrassicin, while others reported traces or even no detectable GLSs in the fermented product [10, 11]. Kim and Rhee [12] also reported a 25 % loss of total GLSs in radish (*Raphanus sativus* L.) compared to the total GLSs in raw radish on the third day of radish *kimchi* fermentation. Moreover, Suzuki et al. [36] reported smaller quantities of the GLSs in

Table 3 Glucosinolate concentration ($\mu\text{mol}/10\text{ g DM}$) in Indian mustard leaves during the production of *sayur asin* ($n = 6$)

Glucosinolate	Withering treatment	Fermentation (days)			
		1	3	5	7
Sinigrin	Oven	4.5 \pm 2.6 ^a	n.d.	n.d.	n.d.
	MW 180	3.3 \pm 0.7 ^a	n.d.	n.d.	n.d.
	MW 900	22.6 \pm 5.8 ^{b,2}	12.3 \pm 2.9 ¹	11.3 \pm 1.8 ¹	11.4 \pm 2.5 ¹
4-Hydroxy-glucobrassicin	Oven	0.6 \pm 0.3 ^{ab}	n.d.	n.d.	n.d.
	MW 180	0.3 \pm 0.1 ^a	n.d.	n.d.	n.d.
	MW 900	1.2 \pm 0.7 ^{b,2}	1.4 \pm 0.5 ²	1.7 \pm 1.1 ²	0.4 \pm 0.2 ¹
Glucobrassicin	Oven	n.d.	n.d.	n.d.	n.d.
	MW 180	n.d.	n.d.	n.d.	n.d.
	MW 900	n.d.	n.d.	n.d.	n.d.
4-Methoxy-glucobrassicin	Oven	0.2 \pm 0.1 ^{ab}	n.d.	n.d.	n.d.
	MW 180	0.2 \pm 0.05 ^a	n.d.	n.d.	n.d.
	MW 900	0.4 \pm 0.1 ^b	0.2 \pm 0.06	n.d.	n.d.

Oven, MW 180, and MW 900 = oven at 35 °C for 2.5 h, MW at 180 W for 4.5 min, and at 900 W for 2 min, respectively

n.d. not detectable

Mean reported of duplicate analyses from 3 treatment replications

Values are presented as mean \pm SD

Values with superscript letters on the same column and superscript number on the same row, which are different are significantly different ($P < 0.05$)

nozawana-zuke, a fermented product of *Brassica rapa* L. in Japan, relative to the raw *nozawana*. It could be that the cell and vacuole walls were damaged during fermentation due to the increase of osmotic pressure as a result of difference concentration of salt and water activity. Consequently, hydrolysis of GLSs by active myrosinase could have occurred.

The sample that was withered by MW at 900 W does not contain active myrosinase showed a much lower reduction of GLSs during fermentation. Accordingly, Sarvan et al. [15] observed that GLSs were better retained during fermentation in *sauerkraut* where myrosinase was inactivated (by blanching) prior to fermentation. However, the GLSs were reduced (about 50 % reduction for sinigrin) in the pre-blanching upon 71 h fermentation samples [15]. In the present study, about 80 % of GLSs were lost during the fermentation even when the myrosinase was inactivated upon withering. Therefore, it might be that during fermentation of the *sayur asin* some myrosinase-like activity was produced by bacteria that catalysed the GLS hydrolysis [19, 20].

The present study shows that inactivation of myrosinase in Indian mustard by using a MW heat treatment can lead to a partial retention of GLS upon fermentation in the product. Ingestion of GLS-containing products without active plant myrosinase can still result in the formation and bioavailability of bioactive breakdown products by enzymes from the gut flora, although the catalytic efficiency of the plant endogenous myrosinase is higher [6, 19, 37–39].

The GLSs breakdown product formation upon withering and fermentation should be also analysed to better

understand the mechanisms that lead to different breakdown profiles in the final product. Among many other factors, the pH plays an important role to influence the breakdown product profile [2]. Moreover, the activity of epithiospecifier proteins may also affects the further conversions upon enzymatic GLS hydrolysis, i.e. the nitriles formation is higher when such proteins are active [2].

LAB and colony count during *sayur asin* fermentation

The natural LAB on the leaves were not detected after MW withering. Therefore, a standard inoculum was added into coconut water prior to fermentation. The total LAB and total colony count in *sayur asin* during fermentation are shown in Table 4. Total LAB increased only until the first to third day of fermentation followed by a steady decline into 7.1–7.3 log CFU/mL. There is no difference in effect of withering methods on the total LAB in the samples. It is likely that in each sample, there were similar conditions that led to similar LAB growth. A previous study on microbial ecology of *sayur asin* reported a successional development of species within the LAB group during *sayur asin* fermentation. Fermentation was initiated by *Leuconostoc mesenteroides* and *Lactobacillus confusus* and completed by *Lac. curvatus*, *Pediococcus pentosaceus* and *Lac. plantarum* [16].

Another study on microbial community of *fu-tsai* and *suan-tsai*, traditional fermented mustard (*B. juncea* Coss.) products of Taiwan, reported a high level of diversity in

Table 4 Total LAB and total colony count (log CFU/mL) during *sayur asin* fermentation ($n = 3$)

	LAB	COL	Withering treatment					
			Oven		MW 180		MW 900	
			LAB	COL	LAB	COL	LAB	COL
Raw leaves	4.4 ± 0.0	7.2 ± 0.2						
After withering		n.d.	3.4 ± 0.1 ^a	n.d.	5.7 ± 0.5 ^b	n.d.	6.4 ± 0.5 ^b	
Fermentation day-1		8.9 ± 0.2 ¹	8.8 ± 0.1 ^a	9.0 ± 0.3 ¹	9.2 ± 0.1 ^b	9.1 ± 0.1 ¹	9.1 ± 0.1 ^b	
Fermentation day-3		9.1 ± 0.2 ¹	9.0 ± 0.1 ^a	9.3 ± 0.1 ¹	8.9 ± 0.1 ^a	9.0 ± 0.2 ¹	9.0 ± 0.2 ^a	
Fermentation day-5		8.0 ± 0.3 ¹	7.9 ± 0.4 ^a	7.9 ± 0.2 ¹	7.9 ± 0.1 ^a	7.8 ± 0.2 ¹	7.7 ± 0.2 ^a	
Fermentation day-7		7.2 ± 0.2 ¹	6.7 ± 0.2 ^a	7.1 ± 0.3 ¹	6.9 ± 0.3 ^{ab}	7.3 ± 0.2 ¹	7.3 ± 0.2 ^b	

Oven, MW 180, and MW 900 = oven at 35 °C for 2.5 h, MW at 180 W for 4.5 min, and at 900 W for 2 min, respectively

LAB, COL total lactic acid bacteria, total colony count

n.d. not detectable

Values are presented as mean ± SD

Values with superscript letters or numbers on the same row, which are different are significantly different ($P < 0.05$)

LAB at the different stages of fermentation, including *Enterococcus faecalis*, *Lac. alimentarius*, *Lac. brevis*, *Lac. plantarum*, *Leu. citreum*, *Leu. mesenteroides*, *Ped. pentosaceus*, *Weissella cibaria*, and others [21]. The diversity of LAB was also reported during fermentation of cabbage *kimchi* [17] and *sauerkraut* [40].

The present study also found that total colony in *sayur asin* increased to 8.8–9.2 log CFU/mL after the first day of fermentation as compared to the ones in the withered leaves, followed by a decline into about 7.7–7.9 and 6.7–7.3 log CFU/mL after the fifth and seventh day of fermentation, respectively. Accordingly, Puspito and Fleet [16] reported the total bacterial count is similar with the total LAB count; therefore, LAB are mainly responsible for the fermentation of *sayur asin* fermentation. Additionally, total yeast was detected at 2.6 ± 0.4 log CFU/mL at day 0, but no yeast colonies were detected after the seventh day of fermentation (data not shown). At day 0, total *B. cereus* was also detected, but uncounted due to interference of other microbes. While at the seventh day of fermentation, the total *B. cereus* was not detected (data not shown).

Conclusions

Sinigrin is the most dominant GLS in raw Indian mustard, accounting for about 95 % of the total GLSs with a concentration of $179 \pm 52 \mu\text{mol}/10 \text{ g DM}$. Both withering and fermentation substantially reduced the GLS concentration in the final *sayur asin* product. The myrosinase activity contributed strongly to the remaining GLS in the vegetable upon fermentation. A withering treatment that fully inactivates myrosinase results in a fermented product containing substantial amounts of GLS, whereas treatments with

residual myrosinase activity result in products void of GLS. The total LAB and total colony counts were not affected by the preparation methods. Lactic acid bacteria are mainly responsible for the fermentation of *sayur asin*.

The present study revealed the importance of myrosinase activity on GLS concentration of Indian mustard during fermentation. This finding can contribute to design the *sayur asin* production in order to obtain a final product that has higher GLS concentration plus the probiotics of a fermented product. Nevertheless, further studies are needed to identify the factors, i.e. pH, salinity, substrate, and LAB, and the breakdown products of the GLS hydrolysis to better understand the mechanisms underlying the GLS degradation during fermentation. Moreover, other withering methods can be investigated such as blanching and steaming, to better retain GLSs upon *sayur asin* production.

Acknowledgments The project entitled *Glucosinolates Behaviour throughout the Production of "Sayur Asin"* was funded by the International Foundation for Science (IFS, Sweden), Grant No. E/5360-1. The authors thank Xandra Bakker-de Haan for the microbiological analysis assistance.

Conflict of interest None.

Compliance with Ethics Requirements This article does not contain any studies with human or animal subjects.

References

- Clarke DB (2010) Glucosinolates, structures and analysis in food. *Anal Method* 2:310–325
- Verkerk R, Schreiner M, Krumbein A, Ciska E, Holst B, Rowland I, De Schrijver R, Hansen M, Gerhäuser C, Mithen R (2009) Glucosinolates in *Brassica* vegetables: the influence of the food

supply chain on intake, bioavailability and human health. *Mol Nutr Food Res* 53:S219–S265

- Halkier BA, Gershenson J (2006) Biology and biochemistry of glucosinolates. *Annu Rev Plant Biol* 57:303–333
- Cartea ME, Velasco P (2008) Glucosinolates in *Brassica* foods: bioavailability in food and significance for human health. *Phytochem Rev* 7:213–229
- Herr I, Büchler MW (2010) Dietary constituents of broccoli and other cruciferous vegetables: implications for prevention and therapy of cancer. *Cancer Treat Rev* 36:377–383
- Traka M, Mithen R (2009) Glucosinolates, isothiocyanates and human health. *Phytochem Rev* 8:269–282
- Bones AM, Rossiter JT (2006) The enzymic and chemically induced decomposition of glucosinolates. *Phytochemistry* 67:1053–1067
- Caplice E, Fitzgerald GF (1999) Food fermentations: role of microorganisms in food production and preservation. *Int J Food Microbiol* 50:131–149
- Tolonen M, Taipale M, Viander B, Pihlava JM, Korhonen H, Ryhänen EL (2002) Plant-derived biomolecules in fermented cabbage. *J Agric Food Chem* 50:6798–6803
- Ciska E, Pathak DR (2004) Glucosinolate derivatives in stored fermented cabbage. *J Agric Food Chem* 52:7938–7943
- Martinez-Villaluenga C, Peñas E, Frias J, Ciska E, Honke J, Piskula M, Kozlowska H, Vidal-Valverde C (2009) Influence of fermentation conditions on glucosinolates, ascorbigen, and ascorbic acid content in white cabbage (*Brassica oleracea* var. *capitata* cv. Taler) cultivated in different seasons. *J Food Sci* 74:C62–C67
- Kim MR, Rhee HS (1993) Decrease of pungency in “radish kimchi” during fermentation. *J Food Sci* 58:128–131
- Nugraheni PY, Verkerk R, Widanarko B, Dekker M (2013) A mechanistic perspective on process induced changes in glucosinolate content in *Brassica* vegetables: a review. *Crit Rev Food Sci Nutr* doi:10.1080/10408398.2012.688076
- Ruiz-Rodriguez A, Marín FR, Ocaña A, Soler-Rivas C (2008) Effect of domestic processing on bioactive compounds. *Phytochem Rev* 7:345–384
- Sarvan I, Valerio F, Lonigro SL, de Candia S, Verkerk R, Dekker M, Lavermicocca P (2013) Glucosinolate content of blanched cabbage (*Brassica oleracea* var. *capitata*) fermented by the probiotic strain *Lactobacillus paracasei* LMG-P22043. *Food Res Int* 54:706–710
- Puspito H, Fleet GH (1985) Microbiology of *sayur asin* fermentation. *Appl Microbiol Biotechnol* 22:442–445
- Lee C-H (1997) Lactic acid fermented foods and their benefits in Asia. *Food Control* 8:259–269
- Nugraheni PY, Priatko CA, Verkerk R, Dekker M, Widanarko B (2013) Reduction of glucosinolates content during *sayur asin* production. *Jurnal Teknologi dan Industri Pangan (Journal of Food Technology and Industry)* 24:236–240
- Oliviero T, Verkerk R, Vermeulen M, Dekker M (2014) *In vivo* formation and bioavailability of isothiocyanates from glucosinolates in broccoli as affected by processing conditions. *Mol Nutr Food Res* 58:1447–1456
- Shapiro TA, Fahey JW, Wade KL, Stephenson KK, Talalay P (1998) Human metabolism and excretion of cancer chemoprotective glucosinolates and isothiocyanates of cruciferous vegetables. *Cancer Epidemiol Biomark Prev* 7:1091–1100
- Chao S-H, Wu R-J, Watanabe K, Tsai Y-C (2009) Diversity of lactic acid bacteria in *suan-tsai* and *fu-tsai*, traditional fermented mustard products of Taiwan. *Int J Food Microbiol* 135:203–210
- Van Eylen D, Indrawati O, Hendrickx M, Van Loey A (2006) Temperature and pressure stability of mustard seed (*Sinapis alba* L.) myrosinase. *Food Chem* 97:263–271
- Verkerk R, Dekker M (2004) Glucosinolates and myrosinase activity in red cabbage (*Brassica oleracea* L. var. *Capitata f. rubra* DC.) after various microwave treatments. *J Agric Food Chem* 52:7318–7323
- Fuller Z, Louis P, Mihajlovski A, Rungapamestry V, Ratcliffe B, Duncan AJ (2007) Influence of cabbage processing methods and prebiotic manipulation of colonic microflora on glucosinolate breakdown in man. *Brit J Nutr* 98:364–372
- Rungapamestry V, Duncan AJ, Fuller Z, Ratcliffe B (2006) Changes in glucosinolate concentrations, myrosinase activity, and production of metabolites of glucosinolates in cabbage (*Brassica oleracea* var. *capitata*) cooked for different durations. *J Agric Food Chem* 54:7628–7634
- Ludikhuyze L, Ooms V, Weemaes C, Hendrickx M (1999) Kinetic study of the irreversible thermal and pressure inactivation of myrosinase from broccoli (*Brassica oleracea* L. cv. *Italica*). *J Agric Food Chem* 47:1794–1800
- Yen GC, Wei QK (1993) Myrosinase activity and total glucosinolate content of cruciferous vegetables, and some properties of cabbage myrosinase in Taiwan. *J Sci Food Agric* 61:471–475
- Oliviero T, Verkerk R, Van Boekel M, Dekker M (2014) Effect of water content and temperature on inactivation kinetics of myrosinase in broccoli (*Brassica oleracea* var. *italica*). *Food Chem* 163:197–201
- Xiao D, Srivastava SK, Lew KL, Zeng Y, Hershberger P, Johnson CS, Tramp DL, Singh SV (2003) Allyl isothiocyanate, a constituent of cruciferous vegetables, inhibits proliferation of human prostate cancer cells by causing G2/M arrest and inducing apoptosis. *Carcinogenesis* 24:891–897
- Bhattacharya A, Tang L, Li Y, Geng F, Paonessa JD, Chen SC, Wong MK, Zhang Y (2010) Inhibition of bladder cancer development by allyl isothiocyanate. *Carcinogenesis* 31:281–286
- Font R, del Río M, Fernández-Martínez JM, de Haro-Bailón A (2004) Use of near-infrared spectroscopy for screening the individual and total glucosinolate contents in Indian mustard seed (*Brassica juncea* L. Czern. & Coss.). *J Agric Food Chem* 52:3563–3569
- He H, Fingerling G, Schnitzler W (2003) Changes in glucosinolate concentrations during growing stages of tai tsai (*Brassica campestris* L. ssp. *chinensis* var. *tai-tsai* Hort.) and pootherb mustard (*Brassica juncea* Coss.). *Acta Hort* 620:77–84
- Krumbein A, Schonhof I, Schreiner M (2005) Composition and contents of phytochemicals (glucosinolates, carotenoids and chlorophylls) and ascorbic acid in selected *Brassica* species (*B. juncea*, *B. rapa* subsp. *nipposinica* var. *chinoleifera*, *B. rapa* subsp. *chinensis* and *B. rapa* subsp. *rapa*). *J Appl Bot Food Qual* 79:168–174
- Palmer MV, Yeung SP, Sang JP (1987) Glucosinolate content of seedlings, tissue cultures, and regenerant plants of *Brassica juncea* (Indian mustard). *J Agric Food Chem* 35:262–265
- He H, Liu L, Song S, Tang X, Wang Y (2003) Evaluation of glucosinolate composition and contents in Chinese *Brassica* vegetables. *Acta Hort* 620:85–92
- Suzuki C, Ohnishi-Kameyama M, Sasaki K, Murata T, Yoshida M (2006) Behavior of glucosinolates in pickling cruciferous vegetables. *J Agric Food Chem* 54:9430–9436
- Higdon JV, Delage B, Williams DE, Dashwood RH (2007) Cruciferous vegetables and human cancer risk: epidemiologic evidence and mechanistic basis. *Pharmacol Res* 55:224–236
- Krul C, Humblot C, Philippe C, Vermeulen M, van Nuenen M, Havenga R, Rabot S (2002) Metabolism of sinigrin (2-propenyl glucosinolate) by the human colonic microflora in a dynamic in vitro large-intestinal model. *Carcinogenesis* 23:1009–1016
- Rungapamestry V, Duncan AJ, Fuller Z, Ratcliffe B (2007) Effect of cooking *Brassica* vegetables on the subsequent hydrolysis and metabolic fate of glucosinolates. *P Nutr Soc* 66:69–81
- Plengvidhya V, Breidt F, Lu Z, Fleming HP (2007) DNA fingerprinting of lactic acid bacteria in sauerkraut fermentations. *Appl Environ Microbiol* 73:7697–7702