

III.B.1.c. 2018 Co author_Proceeding icsaf 2018_banana_repository.pdf

 Unika Soegijapranata1

Document Details

Submission ID

trn:oid::28973:127360378

Submission Date

Feb 2, 2026, 2:53 PM GMT+7

Download Date

Feb 2, 2026, 2:54 PM GMT+7

File Name

III.B.1.c. 2018 Co author_Proceeding icsaf 2018_banana_repository.pdf

File Size

1.2 MB

19 Pages

4,185 Words

22,225 Characters

13% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

Filtered from the Report

- ▶ Bibliography
- ▶ Quoted Text
- ▶ Cited Text
- ▶ Small Matches (less than 10 words)

Exclusions

- ▶ 4 Excluded Sources

Match Groups

- 34 Not Cited or Quoted 13%**
Matches with neither in-text citation nor quotation marks
- 0 Missing Quotations 0%**
Matches that are still very similar to source material
- 0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
- 0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 13% Internet sources
- 5% Publications
- 6% Submitted works (Student Papers)

Integrity Flags

0 Integrity Flags for Review

No suspicious text manipulations found.

Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

Match Groups

- **34 Not Cited or Quoted 13%**
Matches with neither in-text citation nor quotation marks
- **0 Missing Quotations 0%**
Matches that are still very similar to source material
- **0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
- **0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 13% Internet sources
- 5% Publications
- 6% Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

| | | | |
|----|-----------------|--|-----|
| 1 | Internet | | |
| | | coa.ctu.edu.vn | 2% |
| 2 | Submitted works | | |
| | | Universiti Teknologi Malaysia on 2019-06-30 | 2% |
| 3 | Internet | | |
| | | foodbiotech.au.edu | 1% |
| 4 | Internet | | |
| | | hillpublisher.com | <1% |
| 5 | Internet | | |
| | | d-nb.info | <1% |
| 6 | Internet | | |
| | | repository.au.edu | <1% |
| 7 | Internet | | |
| | | anyflip.com | <1% |
| 8 | Internet | | |
| | | dost.hochiminhcity.gov.vn | <1% |
| 9 | Publication | | |
| | | Tran Quang Hieu, Doan Thi Thanh Thao. "Enhancing the Solubility of Curcumin M... | <1% |
| 10 | Internet | | |
| | | lppm.ub.ac.id | <1% |

| | | | |
|----|-----------------|------------------------------|-----|
| 11 | Internet | www.researchgate.net | <1% |
| 12 | Internet | qmfs.hust.edu.vn | <1% |
| 13 | Internet | www.hindawi.com | <1% |
| 14 | Internet | ourarchive.otago.ac.nz | <1% |
| 15 | Internet | stu.edu.vn | <1% |
| 16 | Internet | vdocuments.mx | <1% |
| 17 | Internet | www.au.edu | <1% |
| 18 | Submitted works | University of Debrecen / DEA | <1% |
| 19 | Internet | journals.scicell.org | <1% |
| 20 | Internet | www.myfoodresearch.com | <1% |

Organized by:

SUCOR
TECHNOLOGY
UNIVERSITY



Co-organized by:



turnitin

miss Ranges of 20 idln 20978y 12 10 00357 on

ICSA
20



turnitin

miss Ranges of 20 idln 20978y 12 10 00357 on

FOOD INNOVATION

FOR ASIAN COMMUNITY DEVELOPMENT

PROCEEDINGS OF THE 3rd INTERNATIONAL CONFERENCE
ON SUSTAINABLE GLOBAL AGRICULTURE AND FOOD SECURITY
09 - 10 November 2018, Ho Chi Minh City, Vietnam



SCIENCE AND TECHNICS PUBLISHING HOUSE

FOOD INNOVATION FOR ASIAN COMMUNITY DEVELOPMENT

Organized by:



Co-organized by:



FOOD INNOVATION

FOR ASIAN COMMUNITY DEVELOPMENT

1
PROCEEDINGS OF THE 3rd
INTERNATIONAL CONFERENCE
ON SUSTAINABLE GLOBAL
AGRICULTURE AND FOOD

09 - 10 November 2018, Ho Chi Minh City, Vietnam



SCIENCE AND TECHNICS PUBLISHING HOUSE

PREFACE

15 The International Conference on Sustainable Global Agriculture and Food (ICSAF) is an important event organized every two years. The 3rd ICSAF was held in Ho Chi Minh City, Vietnam, under the auspices of Vietnam Association of Food Science and Technology (VAFOST). Saigon Technology University, Vietnam (STU) was Organizer, and Assumption University, Thailand (AU), Fu Jen Catholic University, Taiwan (FJCU), Soegijapranata Catholic University, Indonesia (SCU) were Co-Organizers of this conference. The theme was “**Food Innovation for Asian Community Development**”.

16 The purpose of the Conference was to highlight significant research and developments for sustainable global agriculture and food production with an emphasis on product innovation. The Conference also aimed to advance regional cooperation in order to promote and apply the research in food science and technology. Aside from food technology development and food innovation, the conference also discussed issues related to Nutrition, Sensory Science, Food Science, Food Engineering and Technology, Food Safety and Quality. This Proceedings will cover all of these discussions.

The Proceedings is divided into 4 sessions covering a wide range of topics of food science and technology. Furthermore, issues related to agriculture, biotechnology, and the environment are also addressed in line with the theme of the Conference.

The papers have been peer-reviewed and edited by an editorial board of the Organizing Committee to be as readable as possible.


The Organizing Committee wishes to thank all authors and delegates, and all who have contributed to the success of the Conference.



12 **Assoc. Prof. Ngo Tien Hien**
President of Vietnam Association
of Food Science and Technology
(VAFoST)



Assoc. Prof. Cao Hao Thi
Principle of Saigon
Technology University
(STU)



Prof. Dr. Luu Dzuan
Dean of Food Technology Faculty
(STU)

FOOD INNOVATION FOR ASIAN COMMUNITY DEVELOPMENT

PROCEEDINGS OF THE 3rd INTERNATIONAL CONFERENCE ON SUSTAINABLE GLOBAL AGRICULTURE AND FOOD

9-10 November 2018, Ho Chi Minh City, Vietnam



Conference Organizers

Saigon Technology University, Vietnam (STU)

Assumption University, Thailand (AU)

Fu Jen Catholic University, Taiwan (FJCU)

Soegijapranata Catholic University, Indonesia (SCU)



Local Organizing Committee

Assoc. Prof. Dr. Cao Hao Thi
Saigon Technology University (Chairman)

MSc. Luu Mai Huong
Saigon Technology University

Assoc. Prof. Dr. Hoang Kim Anh
Saigon Technology University



Scientific Committee

Prof. Dr. Luu Dzuan
Saigon Technology University (Chairman)

Dr. Churdchai Cheowtirakul
Assumption University, Thailand

Prof. Dr. Bing-Huei Chen
Fu Jen Catholic University, Taiwan

Assoc. Prof. Dr. Hoang Kim Anh
Saigon Technology University

Dr. Probo Y. Nugrahedhi
Soegijapranata Catholic University,
Indonesia

Prof. Dr. Kwan Hwa Park
Seoul National University, Korea

Prof. Dr. Jenshinn Lin
National Pingtung University of
Science and Technology, Taiwan

Assoc. Prof. Dr. Ly Nguyen Binh
Can Tho University

7

Assoc. Prof. Dr. Le Nguyen Doan Duy
Ho Chi Minh City University of Technology

Dr. Phan The Dong
Hoa Sen University

8

Assoc. Prof. Dr. Phan Tai Huan
Nong Lam University of Hochiminh City

Assoc. Prof. Dr. Nguyen Minh Thuy
Can Tho University

18

19

Assoc. Prof. Dr. Pham Van Hung
International University, Vietnam National
University in Hochiminh City

Prof. Dr. Le Van Viet Man
Hochiminh City University of
Technology

Dr. Le Minh Hung
Sub-Institute of Agricultural Engineering
and Post-harvest Technology

Dr. Pham Kim Phuong
Saigon Hitech Analytical Center



Secretariat

Ms. Tran Ngoc Hieu
Saigon Technology University

Mr. Ngo Trinh Tat Dat
Saigon Technology University

Ms. Doan Nguyen Thuy Quynh
Saigon Technology University

Mr. Nguyen Thanh Sang
Saigon Technology University

INDEX

PREFACE

SESSION 1

NUTRITION, SENSORY, PACKAGING AND FOOD DEVELOPMENT

| NAME OF ARTICLE | AUTHOR | PAGE |
|--|---|------|
| Gearing towards community service success cases: from laboratory bench to society | Patchanee Yasurin | 17 |
| Effect of packaging materials on quality changes during storage of nonsee-peel cured jellyfish | Atittaya Tandhanskul; Porranee Hempanpairoh | 24 |
| <i>In vivo</i> nutritive values of protein extracted from brackish gutweed <i>Enteromorpha</i> sp. | Tran Ngoc Hieu; Le Diem My; Dao Phat Lam; Le Van Viet Man; Hoang Kim Anh | 32 |
| Consumers' drivers for acceptance of Vietnamese coffees. Application of CATA question and preference mapping | Pham Tuong Thuy Ung; Luu M. Huong; Ly Ngan; Luu Thuy; Pham Thanh; Luu Dzuan | 43 |
| Addition of vegetables in the extrusion of rice puffed snacks to increase vegetables consumption in children | Novita Ika Putri; Jeanny Citrananda Sanusi; Sally Dwi Purnamasari; Probo Yulianto Nugrahedhi | 52 |
| Product development of rice-pork snack using three Thai rice with corn flour and potato starch | Pornpen Panjapiyakul; Pawares Theepsawang | 60 |
| Development of vegetable-rich kaeng liang cube | Roungdao Klinjapo; Kitiporn Tongprasong; Pornpong Sutthirak | 75 |

17

| | NAME OF ARTICLE | AUTHOR | PAGE |
|---|---|---|------|
| | Determination of biochemical composition and <i>in vitro</i> digestibility of protein isolate from brackish algae <i>Chaetomorpha sp.</i> | Bach Ngoc Minh; Nguyen Thanh Sang; Tran Ngoc Hieu; Le Thi Hong Anh; Hoang Kim Anh | 88 |
| 8 | Utilization of gac fruit (<i>Momordica cochinchinensis spreng.</i>) in making Japanese waxy rice cake (mochi) | Tuan Q. Dang; Minh L. N. Ha | 98 |

SESSION 2

FOOD CHEMISTRY, MICROBIOLOGY, QUALITY AND SAFETY

| | NAME OF ARTICLE | AUTHOR | PAGE |
|---|---|---|------|
| 7 | Antibacterial and antioxidant activity of extract from some green vegetable leaf in Vietnam and its application in food product | Le Nguyen Doan Duy; Lam Hoang Quan | 113 |
| 6 | The development of <i>Centella asiatica</i> extract-loaded plga nanoparticles to improve bioavailability | Ming-Shan Huang; Suksun Amornraksa; Malinee Sriariyanun; Suvaluk Asavasanti; Churdchai Cheowtirakul; Patchanee Yasurin | 132 |
| 9 | Synthesis complexes of curcumin with Fe (III), Cu (II) and Zn (II) and initially applied in the treatment of burns in mice | Tran Quang Hieu; Nguyen Van Hai; Luu Mai Huong; Doan Thi Thanh Thao; Le Quang Tri; Nguyen Thanh Sang | 143 |
| | Usage of isomalt as binding agent for improving tablecity of yogurt tablet | Anchidta Tangsuk; Tatsawan Tipvarakarnkoon | 154 |
| | Study of effect of mixed emulsifiers on stability of virgin coconut oil (VCO) emulsion enriched with hyaluronic acid | Sireerat Laodheerasiri; Pattarawadee Vongcharoen | 166 |

| NAME OF ARTICLE | AUTHOR | PAGE |
|---|---|------|
| Evaluation of histamine content in tuna fish and mackerel fish before and after cooking with pineapple and ginger by HPLC-PDA method | Tran Quang Hieu; Nguyen Quynh; Pham Kim Phuong | 177 |
| Bioaccumulation, depuration of heavy metals (As, Cd, Pb) and metabolism of these metals in body of mussels (<i>Maretrix lyrata</i>) during 20 days in artificial media of culture | Pham Kim Phuong; Nguyen Thi Dung; Chu Pham Ngoc Son; Luu Dzuan | 185 |
| Detection and quantification of auramine O contamination level in market foods | Tuan Q. Dang; Trang N. T. Nguyen | 195 |

SESSION 3

FOOD PROCESSING AND ENGINEERING

| NAME OF ARTICLE | AUTHOR | PAGE |
|---|--|------|
| Quality characteristics of banana flour and steamed cake produced from three banana varieties | Rainier R. Zunggaval; Katharina Ardanawari; Probo Y. Nugraedi | 209 |
| Anthocyanin retention improvement of microencapsulated butterfly pea flower crude extract by using freeze drying and β -cyclodextrin | Wunwisa Krasaekoopt; Natnicha Veerathummanoon | 217 |
| Processing of beverages made from Dangshen root (<i>Codonopsis Pilosula</i>) by-products and wastes | Le Minh Hung; Tran Thi Kim Oanh; Nguyen Vinh Phuc | 232 |
| Study of ultrasound-assisted extraction of phenolic compounds from Xao Tam Phan (<i>Paramignya trimera</i>) | Nguyen Thanh Sang; Hoang Kim Anh | 243 |
| Screening and optimization of ultrasound-assisted extraction of betacyanin and total phenolic compounds from dragon fruit peel (<i>Hylocereus undantus</i>) | Ngo Trinh Tac Dat; Pham Quoc Thanh; Tran Pham Xuan Huong; Nguyen Minh Xuan Hong | 253 |
| Effect of phosphates and treatment conditions on quality and safety of frozen fillet tra fish (<i>Pangasianodon hypophthalmus</i>) | Nguyen Van Muoi; Tran Van Nghi; Tran Thanh Truc | 266 |

QUALITY CHARACTERISTICS OF BANANA FLOUR AND STEAMED CAKE PRODUCED FROM THREE BANANA VARIETIES

Rainier R. Zunggaval; Katharina Ardanareswari; *Probo Y. Nugrahedi

Department of Food Technology, Soegijapranata Catholic University, Jl. Pawiyatan Luhur IV/1,
Semarang, 50234, Semarang, Indonesia

*Email: probo@unika.ac.id

ABSTRACT

Processing banana into flour is a way to increase its shelf life and functionality. In food industry, banana flour can be used as wheat flour substitute. In this study, banana flour and, subsequently, the banana steamed cake, were produced from three local banana varieties namely *ambon*, *kepok*, and *nangka*. Physicochemical quality attributes measured on the flour included moisture content, yield, color, water holding capacity, starch, amylose, amylopectin, fiber, and potassium content. The measurement on quality attributes of the steamed cake was texture (hardness and ease of swallowing), color, volume expansion, and potassium content. Hedonic sensory analysis was also performed for the cake. Results showed that different banana varieties gave significant differences in some parameters of the flour namely moisture, yield, color and starch content. When compared to wheat flour, banana flour had higher starch, potassium, and fiber content. Meanwhile, steamed cake quality attributes showed significant differences compared to wheat flour in terms of color, texture, and volume expansion. Hedonic test result showed that wheat flour steamed cake had the highest score but did not differ significantly from the *ambon* banana steamed cake.

Keywords: banana, flour, steamed-cake, variety, quality

1 INTRODUCTION

Banana is one of the most widely found and consumed tropical fruits in Indonesia. Banana production in Indonesia is the highest compared to other fruits such as mangoes and oranges, at up to 6.8 million tons in 2014 [1]. However, postharvest loss of banana is considered high. Therefore, there is a need to increase the shelf life as well as the commercial value of banana. One option is by converting the fruit into flour. Banana flour has low water content and water activity; hence, it will increase the shelf life [2]. Processing banana into flour also gives other benefits, such as its flexibility to be fortified with other nutrients as well as to be main ingredient of final food products [3]. Flour also make banana easier to be handled along the chains of packaging, distribution, and storage [4].

Banana flour was made from unripe banana (yellowish-green color) because ripened banana will disrupt the drying process due to its sugar content that can trigger caramelisation and contributes to the unexpected color, aroma and taste. Nevertheless, banana flour has high nutritional content. In food industries, banana flour is often used to make weaning food and bakery products [4]. Starch content in banana flour contributes to the food texture characteristics.

Many varieties of banana can be processed into flour. It was hypothesized that different banana varieties will affect to different flour characteristics. Studies by Elvis [5] and Abbas *et al* [6] reported that different banana varieties resulted in significant differences of water holding capacity, yield, and color of the flour. These can be due to the differences in physical shape, water content, and starch content of the banana [5].

Moreover, application of banana flour into food production was also reported to have different characteristics than employing wheat flour. Silfia [7] investigated brownies production applying banana or wheat flour and observed that brownies of banana flour had lower water and protein content compared to wheat flour brownies. Sensorially, taste, texture, and aroma of banana flour brownies was more preferred than wheat flour brownies. Our study applies banana flour to produce steamed cake to be compared with cake made from wheat flour.

2 MATERIALS AND METHODS

2.1 Banana flour production

Three local unripe (yellowish green) banana varieties namely *ambon*, *kepok*, and *nangka* were obtained from Gunungpati traditional market in Semarang. Each banana was peeled and sliced using *robot-coupe CL 50* slicer. Then, the slices were soaked in 0.2% NaS₂O₅ solution (w/v) for 15 min. Drying was performed by a cabinet dryer at 60°C for 8 h. Dried banana slices were grinded (*Philips dry-mill blender*) and sieved at 80 mesh.

2.2 Steamed cake production

Steamed cake was made by firstly mixing the dry ingredients such as flour, milk powder, and baking powder. In a different mixing bowl, eggs, sugar and emulsifier were mixed for three min. The dry mixture was added with the batter and mixed. Subsequently, melted margarine was added to the batter and mixed evenly, followed by pouring into baking tray and steaming for 25 min.

2.3 Physicochemical analysis

Analyses of water content, color, and water activity were performed by using moisture balance (*Ohaus MB45*), chromameter (Minolta CR-400), and Aw meter, subsequently. Amylose content and starch content were measured by spectrophotometry method. Water holding capacity (WHC)

was measured by centrifugation method. Crude fiber content was analysed by thermogravimetric method after extraction with 0.25 N H₂SO₄ and 0.25 N NaOH solutions. Potassium content was analysed by atomic absorption spectroscopy. Texture (hardness) was measured by a texture analyzer (*Lloyd instrument TA plus*), and volume expansion was measured by seed displacement method.

2.4 Hedonic test

Sensory analysis was performed by a rating hedonic test at a five-scale, by 40 SCU Food Technology Department students. Some attributes were evaluated, i.e. aroma, color, texture (in terms of hardness and ease of swallowing), taste, and overall perception.

2.5 Data analysis

Data obtained from the physicochemical measurements was analysed by one way ANOVA followed by Duncan test. Meanwhile, data of hedonic test was analysed by Kruskal Wallis followed by Mann Whitney test.

3 RESULTS AND DISCUSSION

3.1 Banana flour

Prior to drying, moisture content of *ambon* banana is the highest ($66.29 \pm 2.93\%$) and significantly different than *kepok* banana ($57.76 \pm 0.76\%$) and *nangka* banana ($59.13 \pm 0.89\%$). This study used mature but unripe banana, at yellowish green color of the peel. In general, ripe banana can have higher moisture content than the unripe one. Putri [8] reported that the moisture contents of *ambon*, *nangka*, and *kepok* bananas are 73.8%, 68%, and 62%, respectively. Low moisture content and high amount of starch of unripe banana can affect to the yield of flour [4]. This study found that *ambon* banana, which contains the highest moisture content, produced the lowest yield (25%) and the *nangka* banana produced the highest yield of flour (29%).

Quality attributes of banana flour can be seen at Table 1. Moisture content of the flour is below 10% and as expected, the highest is found at *ambon* banana flour. *Nangka* banana flour has the highest yield and significantly different as compared to the lowest one. Result on water holding capacity, water activity, fiber, and starch content show that there are no significant difference between banana varieties. Meanwhile, for potassium content, the *nangka* banana flour has the highest among others.

Table 1: Physicochemical quality attributes of banana flour

| Parameter | Banana Variety | | | Ref. (Wheat) | |
|-----------------------------|---------------------------|----------------------------|---------------------------|----------------------------|----------------------------|
| | <i>Kepok</i> | <i>Ambon</i> | <i>Nangka</i> | | |
| Water (%) | 8.35 ± 1.23 ^{ab} | 9.08 ± 1.13 ^a | 7.90 ± 0.59 ^b | | |
| Yield (%) | 29.18 ± 1.42 ^b | 25.71 ± 0.26 ^a | 31.65 ± 2.49 ^b | | |
| Color | L | 88.95 ± 1.54 ^a | 90.35 ± 0.41 ^b | 86.74 ± 1.92 ^c | 95,54 ± 0,24 ^d |
| | a* | -0.90 ± 1.01 ^{bc} | -1.25 ± 0.55 ^a | -0.43 ± 0.75 ^c | -0,85 ± 0,07 ^{ab} |
| | b* | 17.98 ± 2.54 ^b | 16.28 ± 1.31 ^a | 16.14 ± 2.76 ^a | 9,48 ± 0,11 ^c |
| WHC (g H ₂ O/ g) | 40°C | 2.77 ± 0.28 ^a | 3.03 ± 0.38 ^a | 2.97 ± 0.41 ^a | 7,55 ± 0,11 ^b |
| | 60°C | 2.73 ± 0.24 ^a | 2.83 ± 0.43 ^a | 2.75 ± 0.20 ^a | 5,67 ± 0,06 ^b |
| Aw | | 0.46 ± 0.01 ^a | 0.44 ± 0.04 ^a | 0.45 ± 0.06 ^a | |
| Fiber (%) | | 11.6 ± 2.4 ^a | 12.2 ± 2.6 ^a | 11.4 ± 2.8 ^a | 5,3 ± 1,2 ^b |
| Starch (%) | | 65.71 ± 2.68 ^a | 64.60 ± 2.66 ^a | 67.21 ± 4.88 ^a | 58,92 ± 0,81 ^b |
| Amylose (%) | | 35.27 ± 1.20 ^a | 36.22 ± 2.94 ^a | 34.88 ± 0.90 ^a | 22,89 ± 0,44 ^b |
| Amylopectin (%) | | 64.73 ± 2.56 ^{ab} | 63.78 ± 4.65 ^a | 65.12 ± 5.16 ^{ab} | 77,11 ± 0,91 ^b |
| Potassium (mg/100g) | | 783 ± 114 ^a | 967 ± 154 ^b | 988 ± 193 ^b | |

*) Mean ± SD of duplicate determinations (n=9). Values with different superscripts differ significantly (p < 0.05).

Water holding capacity (WHC) is the interaction between components in flour such as protein and starch with water in food. Our study found that the WHC of banana flour is in the range of the WHC of flour needed in the manufacturing of viscous and chewy food products such as soups, sauces, and bakery products [9]. This indicates that banana flour has good WHC to be applied in bakery products. Different temperatures were used to measure WHC, which aimed to check the gelatinization ability of each flour [10]. All banana flour show no significant changes of WHC at 40°C and 60°C, but slightly decrease for the wheat flour at 60°C. This could be due to the heating process did not reach the gelatinization temperature of banana flour, which is 68.8-77.5°C, while the gelatinization temperature of wheat flour is 53-63°C [7]. Another possible explanation is due to different protein content. One of the factors affecting WHC is protein

because it can absorb and bind water, higher protein content will impact to higher binding water ability [11].

The low water activity of the flour (0.44-0.46) could prevent growth of microorganisms and reduce the rate of rancidity of flour [12]. Banana flour also contains higher fiber content than wheat flour. Referring that fiber requirement for adults per day is ± 30 g, depending on total calorie intake [13], intake of 100 g of these banana flour can provide 30% of fiber. Moreover, potassium is one of the essential minerals to control total volume of body fluids, acids and electrolytes, as well as maintain cell function. Potassium contents in the banana flour are 0.783-0.988%. The daily nutritional intake recommended by WHO for potassium is 3510 mg/day for adults [14]. This indicates that 100 grams of banana flour can fulfilled 22-28% potassium needs per day.

3.2 Steamed cake

Quality attributes of steamed cake produced from banana flour and wheat flour are shown in Table 2. Wheat cake has the brightest appearance while *ambon* banana cake is the darkest. The wheat cake is also more yellow than others. Apparently, this is due to the color of the flour; whereas wheat flour is white and banana flour is light brown. All banana steamed cakes show no significant difference in volume expansion but the wheat cake is significantly the highest. This could be due the role of protein in wheat in entrapping gas formed during mixing and contribute to dough viscosity and extensibility [15].

In addition, hardness level of wheat cake is the highest. Texture level of the cake is correlated with the WHC of the flour, i.e. higher WHC is followed by harder texture. This is probably due to the smaller size of starch granule in wheat flour (2-35 μ m) than banana flour (60-105 μ m). Large starch granules can cause bonded water molecules to be easily released during heating or cooking process. The more water is released during heating, the softened the product could be obtained [7].

Table 2: Physicochemical quality attributes of steamed cake

| Parameter | | Type of Flour* | | | |
|---------------|----|--------------------------------|---------------------------------|--------------------------------|--------------------------------|
| | | <i>Kepok</i> | <i>Ambon</i> | <i>Nangka</i> | Wheat |
| Color | L | 63,08 \pm 2,27 ^{bc} | 61,01 \pm 1,60 ^b | 66,53 \pm 2,84 ^c | 82,01 \pm 2,81 ^a |
| | a* | 1,15 \pm 0,09 ^c | 1,86 \pm 0,11 ^b | 0,39 \pm 0,20 ^d | -4,59 \pm 0,16 ^a |
| | b* | 17,49 \pm 0,3 ^b | 17,06 \pm 1,54 ^b | 18,92 \pm 0,84 ^b | 31,36 \pm 1,51 ^a |
| Hardness (gf) | | 158,18 \pm 3,39 ^b | 167,45 \pm 7,17 ^{bc} | 196,53 \pm 4,83 ^c | 319,53 \pm 8,88 ^a |

| Parameter | Type of Flour* | | | |
|----------------------|---------------------------|---------------------------|---------------------------|---------------------------|
| | <i>Kepok</i> | <i>Ambon</i> | <i>Nangka</i> | Wheat |
| Volume Expansion (%) | 48,51 ± 0,20 ^a | 48,78 ± 0,58 ^a | 48,85 ± 0,33 ^a | 53,95 ± 0,99 ^b |
| Potassium (mg/100g) | 44,9 ± 3,8 ^a | 53,8 ± 2,4 ^b | 58,5 ± 0,8 ^c | - |

5 *) Mean ± SD of duplicate determinations (n=9). Values with different superscripts differ significantly (p < 0.05).

Hedonic test that was performed by 40 respondents shows that kepok flour cake is the least liked than others (Table 3). The most liked cake is the wheat flour cake; but, it has no significant difference with *nangka* banana cake.

Table 3: Hedonic rate on the steamed cakes

| Type of Flour | Parameter* | | | | |
|---------------|------------------------|-------------------------|------------------------|-------------------------|------------------------|
| | Aroma | Color | Texture | Taste | Overall |
| <i>Kepok</i> | 2,48±1,24 ^b | 2,78±0,95 ^{bc} | 3,25±0,90 ^a | 2,95±0,93 ^c | 2,80±0,82 ^b |
| <i>Ambon</i> | 3,55±0,85 ^a | 2,88±0,94 ^b | 3,63±0,84 ^a | 3,50±0,91 ^{ab} | 3,45±0,68 ^a |
| <i>Nangka</i> | 3,20±1,18 ^a | 2,35±0,98 ^c | 3,05±0,96 ^a | 3,20±0,97 ^{bc} | 3,03±0,83 ^b |
| Wheat | 3,55±1,11 ^a | 3,68±1,19 ^a | 3,40±1,03 ^s | 3,73±0,75 ^a | 3,73±0,78 ^a |

5 *) Mean ± SD of duplicate determinations (n=9). Values with different superscripts differ significantly (p < 0.05).

For color, panelists preferred the color of wheat flour cake the most. Meanwhile, for texture (hardness and ease of swallowing), there is no significant difference of preference between samples. For aroma and taste, wheat flour cake achieves the highest score compared to *kepok* and *nangka* banana cakes but has no significant difference with *ambon* banana cake. When judging the overall attributes, the steamed cakes made of *ambon* banana and wheat flour are the most preferred by respondents.

4 CONCLUSIONS

Among three banana varieties, *nangka* banana flour produces higher yield, starch, and potassium content, while *ambon* banana flour has brighter color and higher fiber content. For steamed cakes, the wheat flour cake has brighter appearance, harder texture and higher volume expansion. For overall attribute of sensory, there is no significant difference between wheat and *ambon* banana cakes in term of hedonic rating.

REFERENCES

- [1] Martiansyah I. - Petunjuk teknis: budidaya pisang asal kultur *in vitro* dengan teknologi PPBBI. Pusat Penelitian Bioteknologi dan Bioindustri Indonesia, (2014).
- [2] Hui Y. H. - Handbook of food products manufacturing. John Wiley & Sons, Inc. (2007).
- [3] Histifarina D., Rachman A., Rahadian D., and Sukmaya. - Teknologi pengolahan tepung dari berbagai jenis pisang menggunakan cara pengeringan matahari dan mesin pengering, Agrin: Jurnal Penelitian Pertanian **16** (2) (2012) 125-133
- [4] Widowati S. - Petunjuk teknis proses pembuatan aneka tepung dari bahan pangan sumber karbohidrat lokal. Balai Penelitian Pascapanen Pertanian, Jakarta, (2002).
- [5] Elvis O. - Properties of starch extracted from three different plantains (*Musa paradisiaca*) cultivars in Nigeria. Advances in Food Science and Technology **2** (11) (2014) 275-280.
- [6] Abbas F., Saifullah R., and Azhar M. E. - Assessment of physical properties of ripe banana flour prepared from two varieties: cavendish and dream banana. International Food Research Journal **16** (2009) 183-189.
- [7] Silfia - Pengaruh substitusi tepung pisang pada pembuatan brownies terhadap sifat kimia dan penerimaan organoleptik. Jurnal Litbang Industri **2** (2) (2012) 71-78.
- [8] Putri A. R. - Pengaruh kadar air terhadap tekstur dan warna keripik pisang kepok (*Musa parasidiaca formatypica*). Fakultas Pertanian Universitas Hasanuddin. Makassar, (2012).
- [9] Shad M.A., Nawaz H., Noor M., Ahmad H.B., Hussain M., and Choudry M.A. - Functional properties of maize flour and its blend with wheat flour: optimization of preparation conditions by response surface methodology. Pakistan Journal of Botany **45** (6) (2013) 2027-2035.
- [10] Rodriguez-Ambriz S.L., Islas-Hernandez, Agama-Acevedo, Tovar, Bello-Perez. - Characterization of a fiber-rich powder prepared by liquefaction of unripe banana flour. Food Chemistry **107** (2007) 1515-1521.
- [11] Traynham T., Myers D., Carriquiry A.L., and Johnson L.A. - Evaluation of water holding capacity for wheat-soy flour blends. Journal of the American Oil Chemists' Society **84** (2007) 151-155.
- [12] Carter B. P., Galloway M. T., Morris C. F., Weaver G. L., and Carter A. H. - The case for water activity as a specification for wheat tempering and flour production. Cereal Foods World **60** (2015) 166-170.

- [13] Badrialaily - Studi tentang pola konsumsi serat pada mahasiswa. Institut Pertanian Bogor, (2004).
- [14] WHO - Potassium intake for adults and children. WHO Document Production Service. Geneva, (2012).
- [15] Damodaran S. and Paraf A - Food proteins and their application. Marcel Dekker, Inc. New York, (1997).

SAIGON TECHNOLOGY UNIVERSITY, VIETNAM
ASSUMPTION UNIVERSITY, THAILAND
SEOGIJAPRANATA CATHOLIC UNIVERSITY, INDONESIA
FU JEN UNIVERSITY, TAIWAN

FOOD INNOVATION

FOR ASIAN COMMUNITY DEVELOPMENT

PROCEEDINGS OF THE 3rd INTERNATIONAL CONFERENCE ON SUSTAINABLE GLOBAL AGRICULTURE AND FOOD

9-10 November 2018, Ho Chi Minh City, Vietnam

Responsible for publication

VO TUAN HAI

Editor : TRUONG THANH SON
: PHAM THI MAI
Layout : NGUYEN PHAM HAI TRAN
Cover design : HOANG VIET
Copy Editor : HOANG THIEN MINH

SCIENCE AND TECHNICS PUBLISHING HOUSE

70 Tran Hung Dao St., Hoan Kiem Dist., Ha Noi, Viet Nam

Tel: (84)(24) 3942 2443 Fax: (84)(24) 3822 0658

Website: <http://www.nxbkhkt.com.vn> Email: nxbkhkt@hn.vnn.vn

BRANCH IN HO CHI MINH CITY

28 Dong Khoi St., Dist. 1, HCMC, Viet Nam

Tel: (84)(28) 38225062 Fax: (84)(28) 38296628

Email: chinhanhnxbkhkt@yahoo.com.vn

Quantity: 200 copies, size: 21 cm x 29 cm at Nhat Nam Printing and Trading Joint Stock Company

Address: 007 Lo I, Tan Binh Industrial, Tay Thanh Ward, Tan Phu District, HCMC, Viet Nam

Publishing license No: 3977-2018/CXBIPH/6-127/KHKT

Publishing decision No: 130/QĐ-NXBKHK, issued on 05/11/2018

ISBN: 978-604-67-1160-5

Printing completed and copyright deposited in the 4th quarter of 2018