

6. DAFTAR PUSTAKA

Adila, Raisa. (2015). Produksi Rumput Laut Ditargetkan Capai 22,17 Juta Ton. Okezone 29 Januari 2015. <http://economy.okezone.com> 6 April 2015.

Alam, A. (2011). Kualitas Karaginan Rumput Laut Jenis *Eucheuma spinosum* di Perairan Desa Punaga Kabupaten Tahalar. Konsentrasi Eksplorasi Sumber Daya Hayati Laut, Jurusan Ilmu Kelautan dan Perikanan. Universitas Hasanudin. Makassar.

Banerjee, Soumya and Suvendu Bhattacharya. (2014). Food Gels: Gelling Process and New Applications. Critical Reviews in Food Science and Nutrition, 52(4), 334-336, DOI: 10.1080.10408398.2010.500234

Barbeyron, T, Michel G, Potin P, Henrissat B, and Kloareg B. (2000). ι -Carrageenases constitute a novel family of glycoside hydrolases, unrelated to that of κ -carrageenases. Journal of Biological Chemistry, 275, 35499–35505.

Belton, P. S.; Chilvers, G. R.; Morris, V. J.; and Tanner, S. F. (1984). Effects of Group I Cations on the Gelation of Iota Carrageenan. Int. J. Biol. Macromol., 6, 303–308.

Bono, Awang; S.M. Anusuzzaman; and Ong Wan Ding. (2014). Effect of Process Conditions on the Gel Viscosity and Gel Strength of Semi-refined Carrageenan (SRC) Produced from Seaweed (*Kappaphycus alvarezii*). Journal of King Saud University-Engineering Sciences, 2014, 26, 3-9. DOI: 10.1016/j.jksues.2012.06.001

Bunga, Soelvanda M.; Roike I. Montolalu; Johanna W. Harikedua; Lita A.D.Y. Montolalu; Alexander H. Watung; dan Nurmeilita Taher. (2013). Karakteristik Sifat Fisik Kimia Karaginan Rumput Laut *Kappaphycus alvarezii* pada Berbagai Umur Panen yang diambil dari Daerah Perairan Desa Arakan Kabupaten Minahasa Selatan. Jurnal Media Teknologi Hasil Perikanan, Agustus 2013, 1(2), 54-58.

Burey, P., B. R. Bhandari , T. Howes and M. J. Gidley. (2014). Hydrocolloid Gel Particles: Formation, Characterization, and Application. Critical Reviews in Food Science and Nutrition, 48(5), 361-377, DOI:10.1080/10408390701347801

CPKelco. (2001). GENU Carrageenan Book. Rev 8/01. IFT 2002 SAP# 2000791.

Ditjen PEN. (2013). Rumput Laut Indonesia. Warta Ekspor, Ditjen PEN/MJL/004/9/2013 September. Kementerian Perdagangan Republik Indonesia.

Funami, T., M. Hiroe, S. Noda, I. Asai, S. Ikeda and K. Nishimari. (2007). Influence of molecular structure imaged with atomic force microscopy on the rheological behavior

of carrageenan aqueous systems in the presence or absence of cations. *Food Hydrocolloids*, 21, 617-629.

Gulrez, Syed K. H.; Saphwan Al-Assaf; and Glyn O Phillips. (2011). Hydrogels: Methods of Preparation, Characterisation and Applications, Progress in Molecular and Environmental Bioengineering - From Analysis and Modeling to Technology Applications, Prof. Angelo Carpi (Ed.), ISBN: 978-953-307-268-5.

Hambali, E; A. Suryani; dan Wadli. (2004). Membuat Aneka Olahan Rumput Laut. Penebar Swadaya. Jakarta.

Hoffmann, R.A.; A.R. Russell; and M.J. Gidley. (1996). Molecular Weight Distribution of Carrageenans. In: Gums and Stabilisers for the Food Industry 8, G.O. Phillips, P.J. Williams and D.J. Wedlock (eds). IRL Press at the Oxford University Press, Oxford, pp. 137-148.

Holstein, Barry R. (2000). The van der Waals Interaction. American Journal of Physics 2001, 69(4), 441-449. DOI: 10.1119/1.1341251

Imeson, A. P. (2000). Carrageenan. In: Handbook of Hydrocolloids. pp. 87–102. Phillips, G. O. and Williams, P. A., Eds., CRC Press, Boca Raton.

Jaspars, Marcel and Florence Folmer. (2013). Sea Vegetables for Health. Food & Health Innovation Service, Department of Chemistry, School of Natural and Computing Science, University of Aberdeen.

Kusuma, Hendra. (2015). Rumput Laut Indonesia Tak Laku di Negeri Sendiri. Okezone 16 Januari 2015. <http://economy.okezone.com> 6 April 2015.

Lai, V.M.F.; L. Wong P.A.; and Y. Lii C. (2000). Effects of Cation Properties on Sol-gel Transition and Gel Properties of κ -carrageenan. Journal of Food Science, 2000, 65(8), 1332-1337.

Lobban, Christopher S. and Paul J. Harrison. (1994). Seaweed Ecology and Physiology. Cambridge University Press. UK.

López-Pérez, Paula M., Ricardo M. P. da Silva, Iva Pashkuleva, Francisco Parra, Rui L. Reis, and Julio san Roman. (2009). Hydrophobic-Electrostatic Balance Driving the LCST Offset Aggregation-Redissolution Behavior of *N*-Alkylacrylamide-Based Ionic Terpolymers. *Langmuir* 2010, 26(8), 5934-5941. DOI: 10.1021/la903904t

Mangione, M.R., D. Giacomazza, D. Bulone, V. Martorana, G. Cavallaro, and P.L. San Biagio. (2005). K⁺ dan Na⁺ Effects on The Gelation Properties of κ-Carrageenan. *Biophysical Chemistry* 2005, 113, 129-135. DOI: 10.1016/j.bpc.1004.08.005

Marcus Y. (1997). Ion properties. New York: Marcel Dekker, Inc.

Morris, E. R., Rees, D. A., and Robinson, G. (1980). Cation-specific Aggregation of Carrageenan Helices: Domain Model of Polymer Gel Structure. *J. Mol. Biol.*, 138, 349–362.

Michel, A-S.; M.M. Mestdagh; and M.A.V. Axelos. (1997). Physicochemical Properties of Carrageenan Gels in the Presence of Various Cations. *International Journal of Biological Macromolecules*, 21(1-2), 195-200, ISSN 0141-8130.

Necas, J. and L. Bartosicova. (2013). Carrageenan: a review. *Veterinarni Medicina*, 2013, 58(4), 187-205.

Prasetyo, Arief. (2014). Indonesia Berpeluang menjadi Produsen Rumput Laut Dunia. *GATRAnews* 12 Desember 2014. <http://www.gatra.com> 6 April 2015.

Phillips, G.O. and P.A. Williams. (2009). *Handbook of Hydrocolloids* 2nd Edition. Woodhead Publishing.

Piculell, L. (1991). Effects of Ions on the Disorder-Order Transitions of Gel-forming Polysaccharides. *Food Hydrocoll*, 5, 57-69.

Rochas, C. (1982). Étude de la transition sol-gel du kappa-carraghénane. Thèse Docteur ès Sciences Physiques. Université Scientifique et Médicale et Institute National Polytechnique de Grenoble. Grenoble, France.

Rochas, C. and M. Rinaudo. (1980). Activity Coefficients of Counterions and Conformation in Kappa-Carrageenan Systems. *Biopolymers* 19:1675.

Saha, Dipjyoti and Suvendu Bhattacharya. (2010). Hydrocolloids as Thickening and Gelling Agents in Food: A Critical Review. *J Food Sci Technol*, 2010, 47(6), 587–597. DOI 10.1007/s13197-010-0162-6

Schuessler, Phillip wh and William J. Dobbin. (1981). Gel Time Measurement of Resins. *Anal. Chem*, 1981, 53, 2380-2381.

Sediadi, A. dan U. Budihardjo. (2000). Rumput Laut. Kantor Deputi Menteri Negara Riset dan Teknologi. www.ristek.go.id

Snoeren, THM; Payens TAJ; Jeunink J; and Both P. (1975). Electrostatic Interaction between κ -carrageenan and κ -casein. *Milchwissenschaft* 30, 393–396.

Sukri, N. (2006). Karakteristik *Alkali Tread Cottonii* (ATC) dan Karaginan dari Rumput Laut *Eucheuma cotonii* pada Umur Panen yang Berbeda. Program Studi Teknologi Hasul Perikanan Fakultas Perikanan dan Ilmu Kelautan. Institut Petanian Bogor.

Sunardi. (2006). 116 Unsur Kimia Deskripsi dan Pemanfaatannya. Yrama Widya. Bandung.s

Syamsuar. (2006). Karakteristik Karaginan Rumput Laut *Eucheuma cottonii* pada Berbagai Umur Panen, Konsentrasi KOH dan Lama Ekstraksi. Pasca Sarjana Institut Pertanian Bogor.

Tecante, Alberto and María del Carmen Núñez Santiago. (2012). Solution Properties of κ -Carrageenan and Its Interaction with Other Polysaccharides in Aqueous Media, Rheology, Dr. Juan De Vicente (Ed.), ISBN: 978-953-51-0187-1.

Van de Velde F., Lourenco ND, Pinheiro HM, and Bakkerd M. (2002). Carrageenan: a food-grade and Biocompatible Support for Immobilisation Techniques. Advanced Synthesis and Catalysis 344, 815-835.

Warkoyo. (2007). Studi Ekstraksi Karaginan dari Rumput Laut *Eucheuma cottonii* (Kajian Jenis Larutan Perendam dan Lama Perendaman). Jurnal Protein, 2007, 14(1), 49-55.

Webber, Vanessa; Sabrina Matos de Carvalho; Paulo José Ogliari; Leila Hayashi; and Pedro Luiz Manique Barreto. (2012). Optimization of The Extraction of Carrageenan from *Kappaphycus alvarezii* using Response Surface Methodology. Ciênc. Tecnol. Aliment., Campinas, 2012, 32(4), 812-818. DOI: 10.1590/S0101-20612012005000111

Zavitsas, Andreas A. (2005). Aqueous Solution of Calcium Ions: Hydration Number and the Effect of Temperature. *Journal of Physical Chemistry B*, 2005, 109(43), 20636-20640. DOI: 10.1021/jp053909i.