

6. DAFTAR PUSTAKA

- Adam, M.A., M. Bolger., C.D. Carrington., C.E. Coker., G.M. Cramer., M.J. Dinovi dan S. Dolan. (1993). *Guidance Document for Cadmium in Shellfish*. Guidance Document for Trace Element in Seafood. United States Foods and Drugs Adminstration Washington DC (National Academy Press).
- _____ (1993^b). Guidance for Lead in Shellfish. *Guidance for Trace Element in Seafood*. United States Food and Drug Administration , Washington DC (National Academy Press).
- Afiati, N. (2000). Peran Toksikologi dalam Pengelolaan Wilayah Pesisir. A Paper to be presented at The Contribution of Toxicology in Environmental Management, Semarang, Maret 15. (in Bahasa Indonesia) (*to appear*).
- Ahmed, F.E (1991). Seafod Safety. National Academy Press, Washington DC.
- Ahsanullah, M.; M.C. Mobley and D.S. Neglinski. (1984). Accumulation of Cadmium from Cotaminated Water and Sediment by the Shrimp *Callianassa australiensis*. *Marine Biology* 82: 191 –1 197.
- Almatsier, S. (1997). Prinsip Dasar Ilmu Gizi. Pt. Gramedia Pustaka Utama. Jakarta.
- Anonim (1996). Trace Element in Human Nutrition and Health. WHO. Macmillan. Geneva.
- Anonim (1998). International Program on Chemical Safety. Environmental Health Criteria 135 : Cadmium – Environmental Effect. WHO. Macmillan. Geneva.
- Arbai, M. (2002). Kupang, Manfaat dan Kesehatannya. Kompas. Semarang.
- Barnes, R.D. (1974). Invertebrate Zoology. Third Edition. W.B. Saunders Company. Tokyo.
- Broom, M.J. (1985). The Biology and Culture of Marine Bivalve Mollusc of The Genus *Anadara*. ICLARM. Manila.

Chan, H.M.; M. Trifonopoulos; A. Ing; O. Receveur & E. Johnson (1999). Consumption of freshwater Fish in Kahnawake : Riak and Benefits. *Environmental Research Section A* 80 : S213 – S222.

Day, R.A. & A.L. Underwood. (1992). Analisis Kimia Kuantitatif. Erlangga. Jakarta.

Dewi, I.C. (2001). Concentration of Trace Metal Element in Rice and Their Dietary Exposure by Population of The Volcanic Ecosystem of Asembagus, Jawa timur. A Comparison Between irrigation water with unpolluted water and natural acid surface intake. Skripsi. Unika Soegijapranata. Semarang.

Gaman, P.M. & K.B. Sherington (1994). Ilmu Pangan, Pengantar Ilmu Pangan, Nutrisi dan Mikrobiologi, 2nd ed. Gadjah Mada University Press. Yogyakarta.

Gintings, P. (1992). Mencegah dan Mengendalikan Pencemaran industri. CV Muliasari. Jakarta.

Han , B.C. & T.C. Hung (1990). Green oyster caused by copper pollution the Taiwan Coast. *Environmental Pollution*. 65: 347 – 362.

Han, B.C.; W.L. Jeng; Y.N. Tsai & M.S. Jeng. (1993). Depuration of Copper and Zinc By Green Oysters and Blue Mussels of Taiwan. *J.Environment Pollution* 82 : 93-97.

Han, B.C.; W.L. Jeng; T.C. Hung & M.S. Jeng (1994). Copper intake and Health threat by consuming seafood from copper – contaminated coastal enviroment in Taiwan. *Journal of Environmental Toxicology and Chemistry* 13 (5) : 775 – 780.

Hantoro, I (2000). Metal Contents, Microbial Composition and Biomass Reduction of Tissue of Commercially Important Cockle (*Anadara granosa*) from Coastal Areas in Semarang and Kendal. Skripsi. Unika Soegijapranata Semarang.

Hicks, D & D. Kramer. (1999). Seafood Safety : what consumer need to know. Using HACCP to keep Seafood Safety. University of Delaware Sea Grant Collge Program and the National Seafood HACCP Alliance.U.S.

Ibrahim, N. (1995). Trace Element Content of Malaysian Cockles (*Anadara granosa*) *Journal of Food Chemistry*. 54 : 133-135.

Ibrahim, N. & I. Mat. (1995). Trace Element Content In Relation To The Body Weight of The Marine Bivalve, *Anadara granosa* with Special Reference To The Application of

NAA and ICP-AES AS Analytical Techniques. *Journal of Radio Analytical and Nuclear Chemistry*. 195(1) : 203 –208.

Jorhem, L., J. Engman., B.Sundstrom., A.M. Thim. (1994). Trace Element in Crayfish : regional Differences ang Changes Induced by Cooking. *Archives of Environmental Contamination and Toxicology*. Sweden. 26, 137-142.

Leliveld, M.J. (1999). Tissue Distribution of Metals in The Edible Crab (*Scylla serrata*) : A Human Consumption Risk ?. VU Amsterdam and Unika Soegijapranata Semarang.

Mat, I. (1994). Arsenic and Trace metals in commercially important bivalves, *Anadara granosa* and *Paphia undulata*. *Bulletin of Environmental Contamination Toxicology*, 52: 833 – 839.

Mat, I., M.I., Maah and A.Johari. (1994). Trace metals in Sediments and Potensial availability to *Anadara granosa*. *Archieves of Environmental Contaminant Toxicology*, 27 : 54 – 59.

Metcalf & Eddy. (1990). Waste Water Engineering Treatment Disposal Reuse. Mc. Graw Hill. Publishing Company. Ltd. New Delhi.

Moffat, C.F. & Whittle, K.J. (1999). Environmental Contaminants in Food. Sheffield Academic Press. Sheffield.

Muthiah, P., K.A. Narasimhan, C.P. Gopinathan & D. Sundararajan (1992). Larval rearing, spat Production and Juvenile Growth of the blood Clam *Anadara granosa*. *Journal of Marine Biology Association India*. 34 (1&2): 138 – 143.

Nelson , W.G.; B.J. bergen & D.J. Cobb. (1995). Comparison of PCB and Trace Metal Bioaccumulation in the blue mussel, *Mytilus edulis* and ribbed mussel, *Modiolus demissus*, in New Bedford Harbor, Massachusetts. *Environmental Toxicology and Chemistry* 14 (3); 513-521.

Potter, N.N. & J.H. Hotchkiss. (1996). Food Science. Chapman & Hall, Inc. New York.

Rainbow, P.S. (1995). Physiology, Physiochemistry and Metal Uptake – a Crustacean perspective. *Marine Pollution Bulletin* 31 (1-3) : 55 – 59.

Rainbow, P.S. (1990). The Significance of Accumulated Heavy Metal Concentration in Marine Organism. *Marine Pollution Bulletin* 21 : 321 – 324.

Rainbow, P.S. & S.L. White (1989). Comparative Strategies of Heavy Metal Accumulation by Crustaceans : Zinc, Copper and Cadmium in a Decapod, an Amphipod and a Barnacle. *Hidrobiologia*. 174 : 245-262.

Reilly, C. (1980). Metal Contamination of Food. Applied Science Publishers, Ltd. London.

Ryadi, S. (1990). Pencemaran Air (Dasar-Dasar Pokok Penanggulangannya). Karya Anda. Surabaya.

Shahidi, F. & J.R. Botta. (1994). Seafoods : Chemistry, Processing, Technology and Quality. Blackie Academic . London.

Spiegel, M.R.; I Nyoman S. & Ellen G. (1996). Statistika. 2nd.ed. Erlangga. Jakarta.

Steel, W.E. (1960). Water Supply and Sewerage. Mc. Graw-Hill. Tokyo.

Steinhart, C.E. ; M. E. Doyle & B.A. Cochrane (1996). Food Safety. Marcell Dekker, Inc. New York.

Sudarmadji, S.; B. Haryono & Suhardi. (1996). Analisa Bahan Makanan dan Pertanian. Liberty. Yogyakarta.

Sugianto, D.N. (1998). Bioaccumulated level of Lead (pb) in soft tissue of " Kerang Bulu (*Anadara Inflata reeve*) as Bioindicator of Pollutant under Laboratories. *Environmental Journal* 8-12. ISSN 1410 – 6884.

Sugiharto. (1990). Dasar-Dasar Pengelolaan Air Limbah. UI- Press. Jakarta.

Suhardi. (1991). Petunjuk laboratorium Analisa Air dan Penanganan Limbah. UI-Press. Jakarta.

Sulistyawati, I. (2001). Trace metal and Nutritional Contents of Cockle *Anadara granosa* from Several Location at Nothern Coastal of Central Java. Thesis. Universitas Katolik soegijapranata. Semarang.

Supranto, J. (1994). Statistik, Teori dan Aplikasi. 5th ed. Erlangga Jakarta

Tahvonen, R. & J. Kumpulainen. (1996). Content of Lead and Cadmium in Selected Fish species Consumed in Finland in 1993-1994. *Food Additives and Contaminants*. 13(6) : 647-654.

Takarina, N. D. (1998). Heavy metals levels in surficial river sediment from Semarang, Central Java, Indonesia. Environmental Journal 35 –40. ISSN 1410 – 6884.

Valkily, J. M. (1992). Determination and Comparison of Bivalve Growth, with Emphasis on Thailand and Other Tropical Areas. ICLARM Tech. Rep. 36, 125p.

Widianarko, B. (1997). Urban Ecotoxicology : Spatial and Temporal Heterogeneity of Pollution. Faculty of Biology. PhD Thesis. VU Amsterdam

Widianarko, B (2000). Bioindication of urban metal pollution using the Guppy, *Poecilia reticula*. *Lingkungan & Pembangunan* 20 (2) : 202 – 217.

Widianarko, B.I., Sulistyawati & N.M. Van Straalen (2001). Dietary Exposure to Logam Berat in Cockles, *Anadara granosa* from Four Sites Along the North Coastal of Central Java, Indonesia. (*submitted*).

Widianarko, B.; I. Hantoro. & Soedarini (2000). Metals Content, Microbial Composition and Biomass Reduction of Tissue of Commercially Important Cockle (*Anadara granosa*) from Coastal Areas In Semarang and Kendal. Skripsi. Unika Soegijapranata. Semarang.

Widianarko, B., M., Lelived, S., Pujilestari & N.M., Van Straalen (2001). Food Safety risk of trace metals in the Crab *Scylla serrata* from the coastal area of Semarang, Indonesia. (*submitted*).

Widianarko, B & N. M. Van Straalen., (2000). Spatial distribution of logam berat in sediment from urban streams of Semarang, Central Java, Indonesia. *Ecotoxicology & Environmental Safety* 46 : 101 –107.

Winarno, F.G. (1997). Kimia Pangan dan Gizi. Gramedia Pustaka Utama. Jakarta.

Winarno, F.G. (1990). Pengantar Teknologi Pangan. Gramedia. Jakarta.

Whitten, T.; R.E. Soeriaatmadja & S.A. Afiff. (1996). The Ecology of Java and Bali. Periplus Editions (HK) Ltd. Hongkong.

WHO (1996). Trace elements in human nutrition. WHO. Geneva.

Yoseph, F.H. (2002). Trace Metals, Pathogenic bacteria and Nutritional Contents of Cockles from Coasts of Semarang and Demak. *Thesis*. Universitas Katolik Soegijapranata Semarang.

