

6. DAFTAR PUSTAKA

Adams, M.A., M., Bolger, C.D., Carrington, C.E., Coker, G.M., Cramet, M.J., DiNOvi, and S., Dolan,(1993^a). Guidance Document for Cadmium in Shellfish. Guidance Documents for Trace Elements in Seafood. United States Food and Drug Administration, Washington DC. <http://vm.cfsan.fda.gov/~frf/guid-cd.html> (retreived 3 Juli 2002)

_____ (1993^b). Guidance Document for Lead in Shellfish. Guidance Documents for Trace Elements in Seafood United States Food and Drug Administration, Washington DC. <http://vm.cfsan.fda.gov/~frf/guid-Pb.html> (retreived 3 Juli 2002)

Ahmed, F.E. (editor) (1991). Seafood Safety. National Academy Press, Washington DC.

Ahsanullah, M.;M.C. Mobley & D.S Negilski (1984). Accumulation of cadmium from contaminated water and sediment by the shrimp Callianassa australiensis. Marine Biology 82:191-197.

Aviati, N.A., Sumarno, A., Hadiyarto, H.R., Sunoko and B.T., Basuki,(1987). Kandungan logam berat kerang Anadara granosa bisenensis (Bivalvia) di Perairan Pantai Kotamadya Semarang. Lembaga Penelitian Universitas Diponegoro, Semarang , (*unpublished*).

Anonim (1996). Trace Element in Human Nutrition and Health. WHO. Macmillan. Geneva .

Anonim (1998). International Programme on Chemical Safety. Environmetal Health Criteria 200: Copper. WHO. Macmillan. Geneva.

Atlas, R.M. & R. Barthes (1981). Microbial Ecology Fundametal & Application. Addison –Wesley Publ. Co. Philippines.

Babukutty, Y. & J. Chacko (1995). Chemical partitioning and bioavailability of lead and nickel in an estuarine system. *Environmetal Toxicology and Chemistry* 14(3): 369-373.

Baudrimont, M., J. Metivaud, R.M. Brachet, F. Ribeyre & A. Boudou (1997). Bioaccumulation and metallothionein response in the Asiatic clam (*Corbicula fluminea*) after experimental exposure to cadmium and inorganic mercury. *Environmental Toxicology and Chemistry* 16(10):2096-2105.

Browne,D.R., A., Husni, M.J., Risk and J., Eyles (1999). Exposure Assessment: Heavy Metals in Marine Biota of Coastal Indonesia. Research from the UNDIP-McMaster Collaborative Project in Coastal Zone Eco-Development CIDA UPCD agreement No.098/S47074-(0-99).15 p. <http://www.mcmaster.ca/mieh/indo.htm>.

Broom, M.J. (1985). The Biology and Culture of Marine Bivalve Molluscs of The Genus *Anadara*. ICLARM. Manila.

Bu-Olayan, A.H. & Al-Yakoob, S. (1998). Lead, nickel and vanadium in seafood: an exposure assessment for Kuwaiti consumers. *The Science of Total Environment*, 223, 81-86.

Chan, H.M., M. Trifonopoulos; A. Ing; O. Receveur & E. Johnson (1999). Consumption of freshwater fish in Kahnawake: risk and benefits. *Environmetal Research Section A* 80:S213-S222.

Chan, H.M. & P.S. Rainbow (1993). The accumulation of dissolved zinc by the shore crab *Carcinus maenas* (L). *Ophelia* 38(1):13-30.

Cohen, T., S.S., Que Hee & R.F., Ambrose (2001). Trace metals in fish and invertebrates of three California coastal wetlands. *Marine Pollution Bulletin*, 42, 224, 232.

Cope, W.G.; J.G. Wiener & G.J. Atchison (1994). Hepatic cadmium, metal-binding proteins and bioaccumulation in bluegills exposed to aqueous cadmium. *Environmental Toxicology and Chemistry* 13 (4):553-562.

Decker, M.(1996). Food Safety. *Food Research Institute*, New York.

Depledge, M.H. & P.S. Rainbow (1990). Models of regulation and accumulation of trace metals in marine invertebrates. *Compiler of Biochemical Physiology* 97C(1):1-7.

Farley, D. (1998). Dangers of Lead Still Linger. FDA Cosumer January – February. U.S. FDA.

Grandjean, P., P., Weihe, L.L., Needham, V.W., Burse, D.G., Patterson, E.J., Sampson, P.J., Jorgensen and M., Vahter (1995). Relation of a seafood diet to mercury, selenium, arsenic and polychlorinated biphenyl and other organochlorine concentrations in human milk. *Environmental Pollution*, 71, 29-38.

Groten, J.P. and P.J., Van Bladeren (1994). Cadmium bioavailability and health risk in food. *Trends in Food Science and Technology*, 5, 50-55.

Han, B.C. & Hung, T.C. (1990). Green oysters caused by copper pollution on the Taiwan coast. *Environmental Pollution*, 65, 347-362.

Han, B-C., W-L., Jeng, T-C., Hung, M-S., Jeng (1994). Copper intake and health threat by consuming seafood from copper-contaminated coastal environments in Taiwan. *Environmental Toxicology and Chemistry*, 13, 775-780.

Han, B-C., W-L., Jeng, R-Y., Chen, G-T., Fang, T-C., Hung and R-J., Tseng (1998). Estimation of target hazard quotients and potential health risks for metals by consumption of seafood in Taiwan. *Archives of Environmental Contamination and Toxicology*, 35, 711-720.

Han, B-C., W-L., Jeng, Y-C., Hung, Y-C., Ling, M-J., Shieh and L-C., Chien (2000). Estimation of metal and organochlorine pesticide exposures and potential health threat by consumption of oysters in Taiwan. *Environmental Pollution*, 109, 147-156.

Hantoro , I. (2000). Metals Contents, Microbial Composition and Biomass Reduction of Tissue of Commercially Important Cockle (*Anadara granosa*) from Coastal Areas In Semarang and Kendal. Skripsi. Unika Soegijapranata. Semarang.

Hicks, D. & D. Kramer. (1999). Seafood Safety: what consumers need to know. Using HACCP to keep seafood safety. *University of Delaware Sea Grant College Program and the National Seafood HACCP Alliance*. U.S.

<http://www.ocean.udel.edu/mas/seafood/HACCP>.

Jewett, S.C. & Naidu, A.S. (2000). Assessment of heavy metals in red king crabs following offshore placer gold mining. *Marine Pollution Bulletin*, 40, 478-490.

Johansen, P., T., Pars & P., Bjerregaard (2000). Lead,cadmium,mercury and selenium intake by Greenlanders from local marine food. *The Science of Total Environment*, 245, 187-194.

Jones, G.B., P., Mercurio & F., Olivier (2000). Zinc in fish, crabs, oysters and mangrove flora and fauna from Cleveland Bay. *Marine Pollution Bulletin*, 40, 345-352.

Kang, S-G., D.A., Wright & C-H., Koh (2000). Baseline metal concentration in the Asian periwinkle *Littorina brevicula* employed as a biomonitor to assess metal pollution in Korean coastal water. *The Science of Total Environment*, 263, 143-153.

Mat, I.(1994). Arsenic and trace metals in commercially important bivalves, *Anadara granosa* and *Paphia undulata*. *Bulletin of Environmental Contamination Toxicology*, 52, 833-839.

Mati, I., M.I., Maah and A., Johari (1994). Trace metals in sediments and potential availability to *Anadara granosa*. *Archives of Environmental Contamination Toxicology*, 27, 54-59.

Mortimer, M.R.(2000). Pesticide and trace metal concentrations in Queensland estuarine crabs. *Mar. Pollut. Bull.* 41, 359-366.

Renzoni, A., F., Zino & E., Franchi (1998). Mercury levels along the food chain and risk for exposed population. *Environmental Research*, 77, 68-72.

Sarmani, S.& A.A. Majid. (1994). Dietary intake of toxic trace elements from Seafood Consumption in Malaysia. In B. Widanarko, K. Vink & N.M. Van Straalen (eds): *Environmental Toxicology in South East Asia*. VU University Press. Amsterdam. Pp 209-213

Spiegel, M.R.; S., Nyoman & G., Ellen (1996). Statistika.2nd.ed. Erlangga. Jakarta .

Sulistyawaty, I. (2001). Trace metal and nutritional contents of cockle *Anadara granosa* from several locations at northern coastal of central Java. Program Studi Teknologi Pertanian Universitas Katolik Soegijapranata. Semarang.

Takarina, N.D. (1998). Heavy metal levels in surficial river sediment from semarang, Central Java, Indonesia. *Environmental Journal* 35-40. ISSN 1410-6884.

Whitten, T.; R.E. Soeriaatmadja & S.A. Afiff. (1996). The Ecology of Java and Bali. *Periplus Editions (HK) Ltd.* Hongkong.

WHO (1996). Trace elements in human nutrition. WHO, Geneva.

Widianarko, B. (1997). Urban Ecotoxicology: Spatial and Temporal Heterogeneity of Pollution. Faculty of Biology. *PhD Thesis*. VU Amsterdam.

Widianarko, B. (2000). Biindication of urban metal pollution using the Guppy, *Poecilia reticula*. *Lingkungan & Pembangunan* 20 (2): 202-217.

Widianarko, B., I. Sulistyawati & N.M. Van Straalen (2001a). Dietary exposure to logam berat in cockle *Anadara granosa* From Four sites along The North Coast of Central Java, Indonesia. (*submitted*).

Widianarko, B., M., Leliveld, S., Pujilestari & N.M., Van Straalen (2001). Food Safety risk of trace metals in the crab *Scylla serrata* from the coastal area of Semarang, Indonesia. (*submitted*).

Widianarko, B. & N.M. Van Straalen. (2000). Spatial distribution of logam berat insediment from urban streams of Semarang, Central Java, Indonesia. *Ecotoxicology & Environmental Safety* 46: 101- 107.

Yoseph, H. (2002). Trace Metals, Pathogenic Bacteria And Nutritional Contents of Cockle From The Coast of Semarang and Demak. Program Studi Teknologi Pertanian Universitas Katolik Soegijapranata. Semarang.