

6. DAFTAR PUSTAKA

- Anderson, K. and Naujoks, B. (1994). *The SSB Viscometer System: Starch Application*, Cargill, Inc.
- Angela, L. M. S. (2001). *The Molecular Organization in Starch Based Products. The Influence of Polyol Used a Plasticizer*. <http://igistut-archive-library-uu.nl/dissertation/1979557>. Akses tanggal 20 Desember 2015.
- Angka SL, Suhartono TS. (2000). *Bioteknologi Hasil Laut*. Bogor: Pusat Kajian Sumber Daya Pesisir dan Lautan. Institut Pertanian Bogor. hlm 49-56.
- Carlsen, M. H., B. L Halvorsen, K. Holte, S. K. Bøhn, S. Dragland, L. Sampson, C. Willey, H. Senoo, Y. Umezono, C. Sanada, I. Barikmo, N. Berhe, W. C Willett, K. M Phillips, D. R Jacobs Jr, R. Blomhoff. (2010). *The total antioxidant content of more than 3100 foods, beverages, spices, herbs and supplements used worldwide*. *Nutrition Journal* 2010, 9:3.
- Carlson, R.E. and J. Simpson. (1996). *A Coordinator's Guide to Volunteer Lake Monitoring Methods*. North American Lake Management Society. 96 pp.
- Chueamchaitrakun, P.; P. Chompreeda, V. Haruthaithanasan, T. Suwonsichon and S. Kasemsamran. (2011). *Physical Properties of Butter Cake Made from Mixed Hom-Mali and Glutinous Rice Flours*. *Kasetsart J. (Nat. Sci.)* 45 : 295 - 304 (2011).
- Corke, H. and Bejosano, F. (1999) *Effects of Amaranthus and Buckwheat proteins on the rheological properties of maize starch*. *Food Chemistry*. 1999; 65: 493-501.
- D'Appolonia, B. L. (1977). *Effect Of Bread Ingredient On Starch Gelatinization Properties As Measured By The Amylgraph*. *J. Cereal Chem.* 9:532-543.
- Fennena, O. R. (1985). *Food Chemistry, 2nd edition*. Marcel Dekker Inc. New York.
- Feri, M. and K. Becker. (2004). *On Rice, Biodiversity & Nutrients*. University of Hohenheim, Germany.
- Harijono, Zubaidah, E dan Aryani, F.N., (2000). *Pengaruh Proporsi Tepung Beras Ketan Dengan Tepung Tapioka dan Penambahan Telur Terhadap Sifat Fisik dan Organoleptik Kue Semprong*. *Jurnal Makanan Tradisional Indonesia*. No.3. Vol.2: 39-45.


- Henrikson, R. (2009). Earth food Spirulina. How this remarkable blue-greenalgae can transform your health and our planet. Ronore Enterprises, Inc. Hawaii USA. 175.
- Henrikson, R. (2010). Sprulina Wold Food Updated and Revised. Ronore Enterprises. American.
- Hoseney, R. C. (1994). Principle of Cereal Science and Technology, 2nd edition. The American Association of Cereal Chemistry, inc. USA.
- Kabinawa, N. (2006). Ganggang Penggempur Aneka Penyakit. Agro Media Pustaka, Jakarta.
- Kusnandar, Ferri. (2010). Mengenal Sifat Fungsional Protein. Departemen Ilmu Teknologi Pangan – IPB.
- Knight, J.W. (1969). *The Strach Industry*. Edisi 1, Pergramon Press.
- Laren, Mc., (1986). The Colour Science of Dyes and pigments. second edition. Adam HilgerLtd, Bristol.
- Lebesi, Dimitra M. and Constantina Tzia. (2009). Effect of the Addition of Different Dietary Fiber and Edible Cereal Bran Source on the Baking and Sensory Characteristic of Cupcakes. *Journal Food Bioprocess Technology*.
- Lindriati, T. (2011). Pengembangan Teknologi Pembuatan Edible Film dengan Ekstruder Ulir Tunggal Sebagai Unit Pengaduk dan Compression Molding Sebagai Unit Pencetak. Disertasi.
- Marliyati, S.A; Sulaiman, A. & Faizal, A. (1999). Pengolahan Pangan Tingkat Rumah Tangga. IPB. Bogor.
- Matz, S.A. (1992). Bakery Technology and Eginering. 3th edition. Van Nostrand Reinhold. Texas.
- Moorhead, K.; B. Capelli; and G.R. Cysewski. (2006). Spirulina Nature’S Superfood. Cyanotech Corporation. Hawaii.
- Nelis, Imanningsih. (2012). Profil Gelatinisasi Beberapa Formulasi tepung-Tepungan untuk Pendugaan Sifat Pemasakan. *Penel Gizi Makan* 2012, 35(1): 13-22



- Pirenantyo, P. dan Limantara, L. (2008). Pigmen *Spirulina* Sebagai Senyawa Anti Kanker. Indonesian Journal of Cancer 4, 155-163
- Pomeranz, Y. (1991). Functional Properties of Food Components. Academic Press, Inc. New York.
- Purnamasari, Indah dan Happy, Januarti (2010). Pengaruh Hidrolisa Asam-Alkohol dan Waktu Hidrolisa Asam terhadap Sifat Tepung Tapioka. Jurusan teknik kimia, fakultas teknik, Universitas Diponegoro.
- Ravi, M., S. Lata De, S. Azharudin and S. F. D. Paul. (2010). The Beneficial Effects of *Spirulina* Focusing on its Immunomodulatory and antioxidant Properties Nutrition and Dietary Supplements.
- Ravichandran, K.; N. M. M. T. Saw; A. A.A. Mohdaly; A. M.M. Gabr; A. Kastell; H. Riedel; Z. Cai; D. Knorr; & I. Smetanska. (2013). Impact of Processing of Red Beet on Betalain Content and Antioxidant Activity. Food Research International 50: 670–675.
- Richmond, A., (1988). *Spirulina*, in: Borowitzka, M.A., Borowitzka, L.J.(eds), Microalga Biotechnology. Cambridge University Press, Cambridge, pp. 85/121.
- Roy, P., Daisuke, N., Takahiro, O., Hiroshi, O., Mmanasikan, T., Nobutaka, N., and Takeo, S. (2010). Cooking Properties of Different From of Rice Cooked With an Automatic Induction Heating System Rice Cooker. As. Food Ag-Ind 3(04). 373-388.
- Schmidt, R.H. (1981). Gelation and coagulation. In Protein Functionality in Foods, ed. J.P. Cherry. American Chemical Society Symposium Serries no.147. American Chemical Society, Washington DC, pp. 131-47.
- Scheer, H., (2006). An Overview of Chlorophylls and Bachteriochlorophylls, Biochemistry, Biophy-sics, Functions and Applications.
- Sudarmadji, S.; B. Haryono & E. Suhardi. (1989). Analisa Bahan Makanan. Liberty. Yogyakarta.
- Swinkels, J.J.M. (1985). Source of starch, its chemistry and physics. Di dalam: G.M.A.V. Beynum dan J.A Roels (eds.). Starch Conversion Technology. Marcel Dekker, Inc., New York.

- Tabita, Aurelya. (2010). Karakteristik Fungsional Protein Spirulina Platensis. Skripsi. Universitas Katolik Soegijapranata Semarang.
- Taggart, P. (2004). Starch as an ingredients : manufacture and applications. Di dalam: Ann Charlotte Eliasson (ed). Starch in Food: Structure, Function, and Application. CRC Press, Baco Raton, Florida.
- Tester, R.F. (1997) "Starch: the polisaccharide fractions". In: Frazier, PJ, Donald. A.M. Richmond, P, editors. Starch Structure and Functionality. London: The Royal Society of Chemistry, 1997. p.163-169.
- Tietze, H.W. (2004). Spirulina Micro Food Macro Blessing. Ed ke-4. Australia: Haralz W Tietze Publishing.
- Tomaselli L. (1997). Morphology, Ultrastucture and Taxonomy of *Arthrospira (Spirulina) maxima* and *Arthrospira (Spirulina) platensis*. Di dalam: Vonshak A, editor. *Spirulina platensis (Arthrospira): Physiology, CellbiologyandBiotechnology*. Taylor&Francis., Bristol, USA. hlm.2
- Tongsiri, S. Mang-Amphan, K and Y. Peerapornpisal. (2010). *Effect of Replacing Fishmeal with Spirulina on Growth, Carcass, Composition and Pigment of the Mekong Giant Catfish*. Asian Journal of Agricultural Science 2(3):106-110.
- Whistler, R. L., J. N. Be Miller dan E. F. Paschall. (1984). Starch : Chemistry and Technology. Academic Press Inc., New York.
- Winarno, F. G., (2002). Kimia Pangan dan Gizi. PT Gramedia Pustaka Utama. Jakarta.

7. LAMPIRAN

Lampiran 1. Diagram Warna



Lampiran 2. Kurva Standar Amilosa

Lampiran 3. Hasil Olah Data dengan SPSS versi 16.0

Uji Tekstur

Uji Normalitas

Tests of Normality

perlakuan	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
Texture konsentersasi 0% Spirulina	,155	16	,200 [*]	,917	16	,153
Texture konsentersasi 1% Spirulina	,218	16	,041	,862	16	,021
Texture Konsentersasi 5% Spirulina	,171	16	,200 [*]	,918	16	,157
Texture Konsentersasi 10% Spirulina	,123	16	,200 [*]	,974	16	,892

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

Post Hoc One Way Anova

Texture

Duncan^a

perlakuan	N	Subset for alpha = 0.05	
		1	2
Konsentersasi 10% Spirulina	16	491,3519	
Konsentersasi 5% Spirulina	16	507,2869	
konsentersasi 1% Spirulina	16		717,9994
konsentersasi 0% Spirulina	16		736,8544
Sig.		,193	,124

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 16,000.

Uji Warna (L)**Uji Normalitas****Tests of Normality**

Perlakuan	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
Lighness Konsenterasi 0% Spirulina menit 0	,146	6	,200*	,979	6	,945
Konsenterasi 0% Spirulina menit 3	,180	6	,200*	,912	6	,453
Konsenterasi 0% Spirulina menit 6	,377	6	,008	,718	6	,009
Konsenterasi 0% Spirulina menit 9	,265	6	,200*	,803	6	,063
Konsenterasi 0% Spirulina menit 12	,225	6	,200*	,912	6	,449
Konsenterasi 0% Spirulina menit 15	,231	6	,200*	,874	6	,242
Konsenterasi 1% Spirulina menit 0	,176	6	,200*	,947	6	,714
Konsenterasi 1% Spirulina menit 3	,136	6	,200*	,978	6	,944
Konsenterasi 1% Spirulina menit 6	,233	6	,200*	,911	6	,443
Konsenterasi 1% Spirulina menit 9	,223	6	,200*	,909	6	,427
Konsenterasi 1% Spirulina menit 12	,272	6	,189	,846	6	,146
Konsenterasi 1% Spirulina menit 15	,219	6	,200*	,907	6	,414
Konsenterasi 5% Spirulina menit 0	,216	6	,200*	,921	6	,510
Konsenterasi 5% Spirulina menit 3	,194	6	,200*	,959	6	,812
Konsenterasi 5% Spirulina menit 6	,247	6	,200*	,939	6	,655
Konsenterasi 5% Spirulina menit 9	,277	6	,169	,908	6	,425
Konsenterasi 5% Spirulina menit 12	,239	6	,200*	,894	6	,342
Konsenterasi 5% Spirulina menit 15	,168	6	,200*	,931	6	,590
Konsenterasi 10% Spirulina menit 0	,238	6	,200*	,900	6	,374
Konsenterasi 10% Spirulina menit 3	,231	6	,200*	,915	6	,473
Konsenterasi 10% Spirulina menit 6	,188	6	,200*	,973	6	,914
Konsenterasi 10% Spirulina menit 9	,200	6	,200*	,920	6	,505
Konsenterasi 10% Spirulina menit 12	,192	6	,200*	,969	6	,885
Konsenterasi 10% Spirulina menit 15	,286	6	,136	,898	6	,361

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

Post Hoc One Way Anova

L_menit_0

Duncan^a

Perlakuan	N	Subset for alpha = 0.05			
		1	2	3	4
Spirulina 10%	6	46,1167			
Spirulina 5%	6		57,7433		
Spirulina 1%	6			76,0683	
Spirulina 0%	6				90,0250
Sig.		1,000	1,000	1,000	1,000

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.

L_menit_3

Duncan^a

Perlakuan	N	Subset for alpha = 0.05		
		1	2	3
Spirulina 5%	6	32,1617		
Spirulina 10%	6	32,1680		
Spirulina 1%	6		44,4533	
Spirulina 0%	6			59,6033
Sig.		,988	1,000	1,000

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.

L_menit_6

Duncan^a

Perlakuan	N	Subset for alpha = 0.05		
		1	2	3
Spirulina 5%	6	30,9217		
Spirulina 10%	6	31,2333		
Spirulina 1%	6		43,3633	
Spirulina 0%	6			58,4650
Sig.		,167	1,000	1,000

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.

L_menit_9

Duncan^a

Perlakuan	N	Subset for alpha = 0.05		
		1	2	3
Spirulina 10%	6	30,0467		
Spirulina 5%	6	30,7300		
Spirulina 1%	6		43,0583	
Spirulina 0%	6			57,5433
Sig.		,051	1,000	1,000

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.

L_menit_12Duncan^a

Perlakuan	N	Subset for alpha = 0.05		
		1	2	3
Spirulina 10%	6	29,7767		
Spirulina 5%	6	29,8850		
Spirulina 1%	6		42,6900	
Spirulina 0%	6			57,3967
Sig.		,684	1,000	1,000

Means for groups in homogeneous subsets are displayed.


a. Uses Harmonic Mean Sample Size = 6,000.

L_menit_15Duncan^a

Perlakuan	N	Subset for alpha = 0.05		
		1	2	3
Spirulina 10%	6	29,6983		
Spirulina 5%	6	29,7483		
Spirulina 1%	6		40,7817	
Spirulina 0%	6			57,0567
Sig.		,864	1,000	1,000

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.



Uji Warna (a*)**Uji Normalitas****Tests of Normality**

Warna_a	Perlakuan	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
Warna_a	Konsentrasi 0% Spirulina menit 0	,205	6	,200*	,961	6	,830
	Konsentrasi 0% Spirulina menit 3	,204	6	,200*	,918	6	,493
	Konsentrasi 0% Spirulina menit 6	,227	6	,200*	,941	6	,666
	Konsentrasi 0% Spirulina menit 9	,231	6	,200*	,897	6	,358
	Konsentrasi 0% Spirulina menit 12	,258	6	,200*	,940	6	,659
	Konsentrasi 0% Spirulina menit 15	,250	6	,200*	,916	6	,480
	Konsentrasi 1% Spirulina menit 0	,249	6	,200*	,879	6	,263
	Konsentrasi 1% Spirulina menit 3	,181	6	,200*	,928	6	,566
	Konsentrasi 1% Spirulina menit 6	,184	6	,200*	,974	6	,917
	Konsentrasi 1% Spirulina menit 9	,278	6	,164	,903	6	,392
	Konsentrasi 1% Spirulina menit 12	,217	6	,200*	,943	6	,684
	Konsentrasi 1% Spirulina menit 15	,233	6	,200*	,925	6	,546
	Konsentrasi 5% Spirulina menit 0	,162	6	,200*	,961	6	,826
	Konsentrasi 5% Spirulina menit 3	,166	6	,200*	,951	6	,751
	Konsentrasi 5% Spirulina menit 6	,289	6	,128	,904	6	,399
	Konsentrasi 5% Spirulina menit 9	,207	6	,200*	,859	6	,187
	Konsentrasi 5% Spirulina menit 12	,252	6	,200*	,893	6	,335
	Konsentrasi 5% Spirulina menit 15	,257	6	,200*	,862	6	,196
	Konsentrasi 10% Spirulina menit 0	,220	6	,200*	,887	6	,303
	Konsentrasi 10% Spirulina menit 3	,253	6	,200*	,916	6	,479
	Konsentrasi 10% Spirulina menit 6	,299	6	,100	,895	6	,343
	Konsentrasi 10% Spirulina menit 9	,195	6	,200*	,955	6	,781
	Konsentrasi 10% Spirulina menit 12	,166	6	,200*	,976	6	,929
	Konsentrasi 10% Spirulina menit 15	,222	6	,200*	,882	6	,278

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

Post Hoc One Way Anova

a_menit_0

Duncan^a

Perlakuan	N	Subset for alpha = 0.05			
		1	2	3	4
Spirulina 10%	6	-8,4150			
Spirulina 5%	6		-7,6667		
Spirulina 1%	6			-4,8133	
Spirulina 0%	6				,3383
Sig.		1,000	1,000	1,000	1,000

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.

a_menit_3

Duncan^a

Perlakuan	N	Subset for alpha = 0.05		
		1	2	3
Spirulina 1%	6	-3,0717		
Spirulina 10%	6		-,6200	
Spirulina 5%	6		-,5983	
Spirulina 0%	6			-,2967
Sig.		1,000	,679	1,000

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.

a_menit_6

Duncan^a

Perlakuan	N	Subset for alpha = 0.05		
		1	2	3
Spirulina 1%	6	-2,6598		
Spirulina 10%	6		-,6450	
Spirulina 5%	6		-,5517	-,5517
Spirulina 0%	6			-,4067
Sig.		1,000	,379	,177

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.

a_menit_9Duncan^a

Perlakuan	N	Subset for alpha = 0.05		
		1	2	3
Spirulina 1%	6	-2,5183		
Spirulina 10%	6		-,6450	
Spirulina 5%	6		-,5367	-,5367
Spirulina 0%	6			-,4667
Sig.		1,000	,163	,361

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.

a_menit_12Duncan^a

Perlakuan	N	Subset for alpha = 0.05		
		1	2	3
Spirulina 1%	6	-2,3467		
Spirulina 10%	6		-,7483	
Spirulina 5%	6			-,5283
Spirulina 0%	6			-,4850
Sig.		1,000	1,000	,539

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.

a_menit_15Duncan^a

Perlakuan	N	Subset for alpha = 0.05		
		1	2	3
Spirulina 1%	6	-2,3117		
Spirulina 10%	6		-,7883	
Spirulina 0%	6			-,5100
Spirulina 5%	6			-,5033
Sig.		1,000	1,000	,942

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.

Uji Warna (b*)**Uji Normalitas****Tests of Normality**

Warna_b	Perlakuan	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
	Konsentrasi 0% Spirulina menit 0	,235	6	,200 [*]	,815	6	,079
	Konsentrasi 0% Spirulina menit 3	,241	6	,200 [*]	,913	6	,459
	Konsentrasi 0% Spirulina menit 6	,185	6	,200 [*]	,965	6	,860
	Konsentrasi 0% Spirulina menit 9	,236	6	,200 [*]	,935	6	,619
	Konsentrasi 0% Spirulina menit 12	,198	6	,200 [*]	,896	6	,353
	Konsentrasi 0% Spirulina menit 15	,220	6	,200 [*]	,898	6	,364
	Konsentrasi 1% Spirulina menit 0	,277	6	,169	,889	6	,314
	Konsentrasi 1% Spirulina menit 3	,177	6	,200 [*]	,975	6	,921
	Konsentrasi 1% Spirulina menit 6	,211	6	,200 [*]	,909	6	,430
	Konsentrasi 1% Spirulina menit 9	,206	6	,200 [*]	,917	6	,485
	Konsentrasi 1% Spirulina menit 12	,164	6	,200 [*]	,982	6	,962
	Konsentrasi 1% Spirulina menit 15	,246	6	,200 [*]	,880	6	,267
	Konsentrasi 5% Spirulina menit 0	,238	6	,200 [*]	,903	6	,393
	Konsentrasi 5% Spirulina menit 3	,162	6	,200 [*]	,945	6	,701
	Konsentrasi 5% Spirulina menit 6	,189	6	,200 [*]	,920	6	,504
	Konsentrasi 5% Spirulina menit 9	,228	6	,200 [*]	,898	6	,364
	Konsentrasi 5% Spirulina menit 12	,250	6	,200 [*]	,887	6	,302
	Konsentrasi 5% Spirulina menit 15	,227	6	,200 [*]	,890	6	,317
	Konsentrasi 10% Spirulina menit 0	,234	6	,200 [*]	,844	6	,140
	Konsentrasi 10% Spirulina menit 3	,246	6	,200 [*]	,932	6	,596
	Konsentrasi 10% Spirulina menit 6	,172	6	,200 [*]	,950	6	,740
	Konsentrasi 10% Spirulina menit 9	,273	6	,184	,918	6	,490
	Konsentrasi 10% Spirulina menit 12	,265	6	,200 [*]	,917	6	,483
	Konsentrasi 10% Spirulina menit 15	,227	6	,200 [*]	,874	6	,241

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

Post Hoc One Way Anova

b_Menit_0

Duncan^a

Perlakuan	N	Subset for alpha = 0.05			
		1	2	3	4
Spirulina 10%	6	-1,5317			
Spirulina 5%	6		-1,1400		
Spirulina 1%	6			-,0350	
Spirulina 0%	6				4,2033
Sig.		1,000	1,000	1,000	1,000

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.

b_Menit_3

Duncan^a

Perlakuan	N	Subset for alpha = 0.05		
		1	2	3
Spirulina 5%	6	,1767		
Spirulina 10%	6	,2950		
Spirulina 1%	6		2,7467	
Spirulina 0%	6			6,1500
Sig.		,180	1,000	1,000

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.

b_Menit_6

Duncan^a

Perlakuan	N	Subset for alpha = 0.05		
		1	2	3
Spirulina 5%	6	,4533		
Spirulina 10%	6	,4933		
Spirulina 1%	6		2,8283	
Spirulina 0%	6			6,1367
Sig.		,757	1,000	1,000

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.

b_Menit_9

Duncan^a

Perlakuan	N	Subset for alpha = 0.05			
		1	2	3	4
Spirulina 10%	6	,4217			
Spirulina 5%	6		,6050		
Spirulina 1%	6			3,1400	
Spirulina 0%	6				5,7667
Sig.		1,000	1,000	1,000	1,000

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.

b_Menit_12Duncan^a

Perlakuan	N	Subset for alpha = 0.05		
		1	2	3
Spirulina 10%	6	,5733		
Spirulina 5%	6	,6083		
Spirulina 1%	6		3,4300	
Spirulina 0%	6			5,2200
Sig.		,775	1,000	1,000

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.

b_Menit_15Duncan^a

Perlakuan	N	Subset for alpha = 0.05		
		1	2	3
Spirulina 10%	6	,6233		
Spirulina 5%	6	,7933		
Spirulina 1%	6		3,6567	
Spirulina 0%	6			5,1983
Sig.		,221	1,000	1,000

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.

Uji Protein**Uji Normalitas****Tests of Normality**

perlakuan	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
Protein konsenterasi 0% Spirulina	,302	6	,093	,775	6	,035
konsenterasi 1% Spirulina	,261	6	,200*	,928	6	,565
konsenterasi 5% Spirulina	,182	6	,200*	,960	6	,822
konsenterasi 10% Spirulina	,209	6	,200*	,904	6	,396

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

Post Hoc One Way Anova

Protein

Duncan^a

perlakuan	N	Subset for alpha = 0.05			
		1	2	3	4
konsentrasi 0% Spirulina	6	7,0623			
konsentrasi 1% Spirulina	6		7,7047		
konsentrasi 5% Spirulina	6			8,8865	
konsentrasi 10% Spirulina	6				11,0750
Sig.		1,000	1,000	1,000	1,000

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.

Uji Amilosa

Uji Normalitas

Tests of Normality

Perlakuan	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
Amilosa Kadar 0% non kukus	,287	6	,135	,870	6	,227
Kadar 0% kukus	,262	6	,200*	,900	6	,374
Kadar 1% non kukus	,318	6	,058	,790	6	,048
Kadar 1% kukus	,246	6	,200*	,881	6	,273
Kadar 5% non kukus	,242	6	,200*	,828	6	,103
Kadar 5% kukus	,259	6	,200*	,874	6	,241
Kadar 10% non kukus	,264	6	,200*	,928	6	,567
Kadar 10% kukus	,194	6	,200*	,896	6	,352

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

Post Hoc One Way Anova

Amilosa_non_kukus

Duncan^a

perlakuan	N	Subset for alpha = 0.05		
		1	2	3
Spirulina 10%	6	7,7207		
Spirulina 5%	6	7,8198	7,8198	
Spirulina 1%	6		7,9497	7,9497
Spirulina 0%	6			8,0467
Sig.		,138	,057	,147

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.

Amilosa_kukusDuncan^a

perlakuan	N	Subset for alpha = 0.05	
		1	2
Spirulina 10%	6	4,4972	
Spirulina 5%	6	4,5540	
Spirulina 1%	6		4,8063
Spirulina 0%	6		4,8648
Sig.		,385	,371

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.

Uji Amilopektin**Uji Normalitas****Tests of Normality**

Perlakuan	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
Amilopektin Kadar 0% non kukus	,286	6	,135	,871	6	,228
Kadar 0% kukus	,261	6	,200*	,900	6	,372
Kadar 1% non kukus	,317	6	,059	,790	6	,048
Kadar 1% kukus	,246	6	,200*	,882	6	,276
Kadar 5% non kukus	,243	6	,200*	,829	6	,106
Kadar 5% kukus	,261	6	,200*	,871	6	,231
Kadar 10% non kukus	,262	6	,200*	,929	6	,574
Kadar 10% kukus	,193	6	,200*	,897	6	,354

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

Post Hoc One Way Anova**Amilopektin_non_kukus**Duncan^a

perlakuan	N	Subset for alpha = 0.05		
		1	2	3
Spirulina 0%	6	91,9531		
Spirulina 1%	6	92,0505	92,0505	
Spirulina 5%	6		92,1802	92,1802
Spirulina 10%	6			92,2793
Sig.		,145	,057	,138

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.

Amilopektin_kukusDuncan^a

perlakuan	N	Subset for alpha = 0.05	
		1	2
Spirulina 0%	6	95,1351	
Spirulina 1%	6	95,1937	
Spirulina 5%	6		95,4459
Spirulina 10%	6		95,5027
Sig.		,370	,385

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.

Uji Aktivitas Antioksidan**Uji Normalitas****Tests of Normality**

Perlakuan	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df.	Sig.	Statistic	df	Sig.
Antioksidan Kadar 0% non kukus	,257	6	,200	,883	6	,285
Kadar 0% kukus	,259	6	,200	,878	6	,261
Kadar 1% non kukus	,208	6	,200	,916	6	,478
Kadar 1% kukus	,269	6	,200	,911	6	,445
Kadar 5% non kukus	,236	6	,200	,889	6	,311
Kadar 5% kukus	,259	6	,200	,895	6	,345
Kadar 10% non kukus	,170	6	,200	,967	6	,871
Kadar 10% kukus	,112	6	,200	,993	6	,996

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

Post Hoc One Way Anova**Antioksidan_non_kukus**Duncan^a

perlakuan	N	Subset for alpha = 0.05			
		1	2	3	4
Spirulina 0%	6	2,3530			
Spirulina 1%	6		3,0443		
Spirulina 5%	6			5,3042	
Spirulina 10%	6				7,1215
Sig.		1,000	1,000	1,000	1,000

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.

Antioksidan_kukusDuncan^a

perlakuan	N	Subset for alpha = 0.05		
		1	2	3
Spirulina 0%	6	,9075		
Spirulina 1%	6	1,0760		
Spirulina 5%	6		2,9915	
Spirulina 10%	6			4,8920
Sig.		,484	1,000	1,000

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.

Uji Aktivitas Viskositas**Uji Normalitas****Tests of Normality**

perlakuan	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
Viskositas konsentersasi 0% Spirulina	,121	6	,200*	,983	6	,964
Viskositas konsentersasi 1% Spirulina	,189	6	,200*	,943	6	,682
Viskositas konsentersasi 5% Spirulina	,243	6	,200*	,816	6	,082
Viskositas konsentersasi 10% Spirulina	,153	6	,200*	,957	6	,794

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

Post Hoc One Way Anova**Viskositas**Duncan^a

perlakuan	N	Subset for alpha = 0.05		
		1	2	3
konsentersasi 0% Spirulina	6	22,5667		
konsentersasi 1% Spirulina	6	23,1167		
konsentersasi 5% Spirulina	6		45,5000	
konsentersasi 10% Spirulina	6			54,3833
Sig.		,107	1,000	1,000

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.

Korelasi Warna L, a*, b* dengan Kandungan Antioksidan Adonan Sebelum Pengukusan

Correlations

		lighness	antioksidan	nilai_b	nilai_a
lighness	Pearson Correlation	1	-,959**	,899**	,954**
	Sig. (2-tailed)		,000	,000	,000
	N	24	24	24	24
antioksidan	Pearson Correlation	-,959**	1	-,786**	-,862**
	Sig. (2-tailed)	,000		,000	,000
	N	24	24	24	24
nilai_b	Pearson Correlation	,899**	-,786**	1	,985**
	Sig. (2-tailed)	,000	,000		,000
	N	24	24	24	24
nilai_a	Pearson Correlation	,954**	-,862**	,985**	1
	Sig. (2-tailed)	,000	,000	,000	
	N	24	24	24	24

** . Correlation is significant at the 0.01 level (2-tailed).

Korelasi Tekstur dengan Kandungan Amilosa Setelah Pengukusan

Correlations

		teksture	Amilosa
teksture	Pearson Correlation	1	,837**
	Sig. (2-tailed)		,000
	N	24	24
Amilosa	Pearson Correlation	,837**	1
	Sig. (2-tailed)	,000	
	N	24	24

** . Correlation is significant at the 0.01 level (2-tailed).

Korelasi Tekstur dengan Kandungan Amilopektin

Correlations

		teksture	Amilopektin
teksture	Pearson Correlation	1	-,837**
	Sig. (2-tailed)		,000
	N	24	24
Amilopektin	Pearson Correlation	-,837**	1
	Sig. (2-tailed)	,000	
	N	24	24

** . Correlation is significant at the 0.01 level (2-tailed).

Korelasi Tekstur dengan Kandungan Protein

Correlations

		teksture	Protein
teksture	Pearson Correlation	1	-,868**
	Sig. (2-tailed)		,000
	N	24	24
Protein	Pearson Correlation	-,868**	1
	Sig. (2-tailed)	,000	
	N	24	24

** . Correlation is significant at the 0.01 level (2-tailed).

Korelasi Warna L, a*, b* dengan Kandungan Antioksidan Adonan, Setelah Pengukusan

Correlations

		Antioksidan_kukus	Lighnes_Kukus	Nilai_a_Kukus	Nilai_b_kukus
Antioksidan_kukus	Pearson Correlation	1	-,775**	,368	-,870**
	Sig. (2-tailed)		,000	,077	,000
	N	24	24	24	24
Lighnes_Kukus	Pearson Correlation	-,775**	1	,000	,961**
	Sig. (2-tailed)	,000		,999	,000
	N	24	24	24	24
Nilai_a_Kukus	Pearson Correlation	,368	,000	1	-,241
	Sig. (2-tailed)	,077	,999		,256
	N	24	24	24	24
Nilai_b_kukus	Pearson Correlation	-,870**	,961**	-,241	1
	Sig. (2-tailed)	,000	,000	,256	
	N	24	24	24	24

** . Correlation is significant at the 0.01 level (2-tailed).

Korelasi Kandungan Amilosa dan Amilopektin terhadap Viskositas Adonan

Correlations

		Amilosa	Amilopektin	Viskositas
Amilosa	Pearson Correlation	1	-1,000**	-,828**
	Sig. (2-tailed)		,000	,000
	N	24	24	24
Amilopektin	Pearson Correlation	-1,000**	1	,828**
	Sig. (2-tailed)	,000		,000
	N	24	24	24
Viskositas	Pearson Correlation	-,828**	,828**	1
	Sig. (2-tailed)	,000	,000	
	N	24	24	24

** . Correlation is significant at the 0.01 level (2-tailed).

Uji Kadar Air

Uji Normalitas

Tests of Normality

Perlakuan		Kolmogorov-Smirnov ^a			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
KadarAir_Sblm_kukus	Spirulina 0%	,262	6	,200*	,863	6	,201
	Spirulina 1%	,281	6	,151	,879	6	,265
	Spirulina 5%	,278	6	,163	,843	6	,137
	Spirulina 10%	,300	6	,099	,882	6	,280

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

Tests of Normality

Perlakuan	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
KadarAir_kukus						
Spirulina 0%	,233	6	,200 [*]	,951	6	,748
Spirulina 1%	,267	6	,200 [*]	,870	6	,228
Spirulina 5%	,218	6	,200 [*]	,830	6	,108
Spirulina 10%	,191	6	,200 [*]	,916	6	,476

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

Post Hoc One Way Anova

KadarAir_Sblm_kukus

Duncan^a

Perlakuan	N	Subset for alpha = 0.05	
		1	
Spirulina 10%	6	64,0875	
Spirulina 5%	6	64,1270	
Spirulina 1%	6	64,1532	
Spirulina 0%	6	64,1607	
Sig.		,736	

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.

KadarAir_kukus

Duncan^a

Perlakuan	N	Subset for alpha = 0.05	
		1	2
Spirulina 0%	6	61,3523	
Spirulina 1%	6	61,4975	
Spirulina 5%	6		63,5805
Spirulina 10%	6		63,7217
Sig.		,631	,641

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 6,000.