

## 7. DAFTAR PUSTAKA

- Abdualrahman, M. A. Y. (2013). Chemical, In-vitro Protein Digestibility, Minerals and Amino Acids Composition of Edible Peanut Seeds (*Arachis hypogaea* L.). *Science International*, 1, 199–202. <https://scialert.net/abstract/?doi=sciintl.2013.199.202>
- Abeshu, Y., & Kefale, B. (2017). Effect of Some Traditional Processing Methods on Nutritional Composition and Alkaloid Content of Lupin Bean. *Journal of Natural Sciences Research*, 7(21), 17–21. <https://iiste.org/Journals/index.php/JNSR/article/viewFile/39929/41047>
- Abusin, S. A. E., Hassan, A. B., & Babiker, E. E. (2009). Nutritional Evaluation of Cooked Faba Bean (*Vicia Faba* L.) And White Bean (*Phaseolus Vulgaris* L.) Cultivars. *Australian Journal of Basic and Applied Sciences*, 3(3), 2484–2490. [https://www.researchgate.net/publication/274623991\\_Nutritional\\_Evaluation\\_of\\_Cooked\\_Faba\\_Bean\\_Vicia\\_Faba\\_L\\_and\\_White\\_Bean\\_Phaseolus\\_Vulgaris\\_L\\_Cultivars](https://www.researchgate.net/publication/274623991_Nutritional_Evaluation_of_Cooked_Faba_Bean_Vicia_Faba_L_and_White_Bean_Phaseolus_Vulgaris_L_Cultivars)
- Adegboye, T. T., Abberton, M. T., Abdelgadir, A. H., Dianda, M., Maziya-Dixon, B., Oyatomi, O. A., ... Babalola, O. O. (2019). Nutrient and Antinutrient Composition of Winged Bean (*Psophocarpus tetragonolobus* (L.) DC.) Seeds and Tubers. *Journal of Food Quality*, 2019, 1–8. <https://doi.org/10.1155/2019/3075208>
- Adeyeye, E. I. (2010). Effect of Cooking and Roasting on the Amino Acid Composition of Raw Groundnut (*Arachishypogaea*) Seeds. *Acta Scientiarum Polonorum Technologia Alimentaria*, 9(2), 201–216. <https://www.food.actapol.net/volume9/issue2/abstract-7.html>
- Adeyeye, E. I., & Agesin, M. O. (2012). Effects of roasting and cooking on the lipid composition of raw groundnut (*Arachis hypogaea*) seeds: dietary implications. *Elixir Food Science*, 42, 6257–6266. [https://www.researchgate.net/publication/275967705\\_Effects\\_of\\_roasting\\_and\\_cooking\\_on\\_the\\_lipid\\_composition\\_of\\_raw\\_groundnut\\_Arachis\\_hypogaea\\_seeds\\_dietary\\_implications](https://www.researchgate.net/publication/275967705_Effects_of_roasting_and_cooking_on_the_lipid_composition_of_raw_groundnut_Arachis_hypogaea_seeds_dietary_implications)
- Adhikari, B., Dhungana, S. K., Ali, M. W., Adhikari, A., Kim, I.-D., & Shin, D.-H. (2018). Resveratrol, total phenolic and flavonoid contents, and antioxidant potential of seeds and sprouts of Korean peanuts. *Food Science and Biotechnology*, 27(5), 1275–1284. <https://doi.org/10.1007/s10068-018-0364-7>
- Ahmad, A., Hayat, I., Arif, S., Masud, T., Khalid, N., & Ahmed, A. (2014). Mechanisms Involved in the Therapeutic Effects of Soybean (*Glycine Max*). *International Journal of Food Properties*, 17(6), 1332–1354. <https://doi.org/10.1080/10942912.2012.714828>
- Ali, E. A. A. M. O., Awadelkareem, A. M., Gasim, S., & Yousif, N. E. A. (2014). Nutritional Composition And Anti Nutrients Of Two Faba Bean (*Vicia faba* L.)

LINES. *International Journal of Advanced Research*, 2(12), 538–544. [https://www.researchgate.net/publication/272999127\\_NUTRITIONAL\\_COMPOSITION\\_AND\\_ANTI\\_NUTRIENTS\\_OF\\_TWO\\_FABA\\_BEAN\\_Vicia\\_faba\\_L\\_LINES](https://www.researchgate.net/publication/272999127_NUTRITIONAL_COMPOSITION_AND_ANTI_NUTRIENTS_OF_TWO_FABA_BEAN_Vicia_faba_L_LINES)

- Amarakoon, R. (2009). The effect of cooking on nutritive quality of selected legumes (doctoral dissertation). Zlin. Retrieved from <https://pdfs.semanticscholar.org/6293/02542140edd09c0c9385cd7230f119a6f743.pdf>
- Arnoldi, A., & Greco, S. (2011). Nutritional and nutraceutical characteristics of lupin protein. *Nutrafoods*, 10(4), 23–29. <https://doi.org/10.1007/bf03223356>
- Ayoola, P. B., & Adeyeye, A. (2010). Effect of Heating on the Chemical Composition and Physico - Chemical Properties of *Arachis hypogea* (Groundnut) Seed Flour and Oil. *Pakistan Journal of Nutrition*, 9(8), 751–754. <https://doi.org/10.3923/pjn.2010.751.754>
- Baloch, F. S., Karaköy, T., Demirbaş, A., Toklu, F., Özkan, H., & Hatipoğlu, R. (2014). Variation of some seed mineral contents in open pollinated faba bean (*Vicia faba* L.) landraces from Turkey. *Turkish Journal of Agriculture and Forestry*, 38, 591–602. <https://doi.org/10.3906/tar-1311-31>
- Banerjee, S., Giri, A. P., Gupta, V. S., & Dutta, S. K. (2017). Structure-function relationship of a bio-pesticidal trypsin/chymotrypsin inhibitor from winged bean. *International Journal of Biological Macromolecules*, 96, 532–537. <https://doi.org/10.1016/j.ijbiomac.2016.12.018>
- Borek, S., Pukacka, S., Michalski, K., & Ratajczak, L. (2009). Lipid and Protein Accumulation in Developing Seeds of Three Lupine Species: *Lupinus luteus* L., *Lupinus albus* L., and *Lupinus mutabilis* Sweet. *Journal of Experimental Botany*, 60(12), 3453–3466. <https://doi.org/10.1093/jxb/Erp186>
- Boschin, G., & Arnoldi, A. (2011). Legumes Are Valuable Sources of Tocopherols. *Food Chemistry*, 127(3), 1199–1203. <https://doi.org/10.1016/j.foodchem.2011.01.124>
- Buckle, K. A., & Sambudi, H. (1990). Effect of soaking and boiling treatments on the quality of winged bean seed. *Journal of the Science of Food and Agriculture*, 53(3), 379–388. <https://doi.org/10.1002/jsfa.2740530310>
- Campos-Mondragón, M. G., Medina-Juárez, L. A., Calderón De La Barca, A. M., Durán-Prado, A., Campos-Reyes, L. C., Oliart-Ros, R. M., ... Angulo, O. (2009). Nutritional composition of new Peanut (*Arachis hypogaea* L.) cultivars. *Grasas y Aceites*, 60(2), 161–167. <https://doi.org/10.3989/gya.075008>
- Carvalho, A. W. D., Natal, D. I. G., Silva, C. O. D., Dantas, M. I. D. S., Barros, E. G. D., Ribeiro, S. M. R., ... Martino, H. S. D. (2013). Heat-treatment reduces anti-nutritional phytochemicals and maintains protein quality in genetically improved

- hulled soybean flour. *Food Science and Technology*, 33(2), 310–315. <https://doi.org/10.1590/s0101-20612013005000048>
- Cazzato, E., Laudadio, V., Ceci, E., & Tufarelli, V. (2014). Effect of sulphur fertilization on fatty acid composition of faba bean (*Vicia faba* L.), white lupin (*Lupinus albus* L.) and pea (*Pisum sativum* L.) grains. *Journal of Food, Agriculture & Environment*, 12(3&4), 136–138. [https://www.researchgate.net/publication/266795153\\_Effect\\_of\\_sulphur\\_fertilization\\_on\\_fatty\\_acid\\_composition\\_of\\_faba\\_bean\\_Vicia\\_faba\\_L\\_white\\_lupin\\_Lupinus\\_albus\\_L\\_and\\_pea\\_Pisum\\_sativum\\_L\\_grains](https://www.researchgate.net/publication/266795153_Effect_of_sulphur_fertilization_on_fatty_acid_composition_of_faba_bean_Vicia_faba_L_white_lupin_Lupinus_albus_L_and_pea_Pisum_sativum_L_grains)
- Cazzato, E., Laudadio, V., Stellacci, A. M., Ceci, E., & Tufarelli, V. (2012). Influence of Sulphur Application on Protein Quality, Fatty Acid Composition and Nitrogen Fixation of White Lupin (*Lupinus Albus* L.). *European Food Research and Technology*, 235(5), 963–969. <https://doi.org/10.1007/S00217-012-1817-5>
- Černý, K., Kordylas, M., Pospíšil, F., Švábenský, O., & Zajíc, B. (1971). Nutritive value of the winged bean (*Psophocarpus palustris* Desv.). *British Journal of Nutrition*, 26(2), 293–299. <https://doi.org/10.1079/bjn19710035>
- Chai, B. C., Voort, J. R. V. D., Grofelnik, K., Eliasdottir, H. G., Klöss, I., & Perez-Cueto, F. J. A. (2019). Which Diet Has the Least Environmental Impact on Our Planet? A Systematic Review of Vegan, Vegetarian and Omnivorous Diets. *Sustainability*, 11(15), 4110. <https://doi.org/10.3390/su11154110>
- Chaieb, N., González, J. L., López-Mesas, M., Bouslama, M., & Valiente, M. (2011). Polyphenols content and antioxidant capacity of thirteen faba bean (*Vicia faba* L.) genotypes cultivated in Tunisia. *Food Research International*, 44(4), 970–977. <https://doi.org/10.1016/j.foodres.2011.02.026>
- Chay, S. Y., Salleh, A., Sulaiman, N. F., Abidin, N. Z., Hanafi, M. A., Zarei, M., & Saari, N. (2018). Blood-pressure lowering efficacy of winged bean seed hydrolysate in spontaneously hypertensive rats, peptide characterization and a toxicity study in Sprague-Dawley rats. *Food & Function*, 9(3), 1657–1671. <https://doi.org/10.1039/c7fo01769c>
- Chen, L., Teng, H., & Xiao, J. (2019). A value-added cooking process to improve the quality of soybean: Protecting its isoflavones and antioxidant activity. *Food Science and Human Wellness*, 8(2), 195–201. <https://doi.org/10.1016/j.fshw.2019.05.001>
- Chung, I.-M., Oh, J.-Y., & Kim, S.-H. (2017). Comparative study of phenolic compounds, vitamin E, and fatty acids compositional profiles in black seed-coated soybeans (*Glycine Max* (L.) Merrill) depending on pickling period in brewed vinegar. *Chemistry Central Journal*, 11(1). <https://doi.org/10.1186/s13065-017-0298-9>
- Comstock, S. S., Maleki, S. J., & Teuber, S. S. (2016). Boiling and Frying Peanuts Decreases Soluble Peanut (*Arachis Hypogaea*) Allergens Ara h 1 and Ara h 2

- But Does Not Generate Hypoallergenic Peanuts. PLOS ONE, 11(6), e0157849. <https://doi.org/10.1371/journal.pone.0157849>
- Craft, B. D., Kosińska, A., Amarowicz, R., & Pegg, R. B. (2010). Antioxidant Properties of Extracts Obtained from Raw, Dry-roasted, and Oil-roasted US Peanuts of Commercial Importance. *Plant Foods for Human Nutrition*, 65(3), 311–318. <https://doi.org/10.1007/s11130-010-0160-x>
- Crépon, K., Marget, P., Peyronnet, C., Carrouée, B., Arese, P., & Duc, G. (2010). Nutritional value of faba bean (*Vicia faba* L.) seeds for feed and food. *Field Crops Research*, 115(3), 329–339. <https://doi.org/10.1016/j.fcr.2009.09.016>
- Dalaram, I. S. (2017). Evaluation of Total Polyphenol Content and Antioxidant Capacity of Different Variety Lupin Seeds. *Potravinarstvo*, 11(1). <https://doi.org/10.5219/678>
- De Lumen, B. O., & Fiad, S. (1982). Tocopherols of winged bean (*Psophocarpus tetragonolobus*) oil. *Journal of Agricultural and Food Chemistry*, 30(1), 50–53. <https://doi.org/10.1021/jf00109a010>
- De Lumen, B. O., & Salamat, L. A. (1980). Trypsin inhibitor activity in winged bean (*Psophocarpus tetragonolobus*) and the possible role of tannin. *Journal of Agricultural and Food Chemistry*, 28(3), 533–536. <https://doi.org/10.1021/jf60229a042>
- Doyle, J. (2001). Leguminosae. *Encyclopedia of Genetics*, 1081–1085. <https://doi.org/10.1006/rwgn.2001.1642>
- Duc, G. (1997). Faba bean (*Vicia faba* L.). *Field Crops Research*, 53(1-3), 99–109. [https://doi.org/10.1016/s0378-4290\(97\)00025-7](https://doi.org/10.1016/s0378-4290(97)00025-7)
- Duval, R. (n.d.). Plant database entry for Winged Bean (*Psophocarpus tetragonolobus*) with 3 images and 27 data details. Retrieved April 1, 2020, from <https://garden.org/plants/view/120697/Winged-Bean-Psophocarpus-tetragonolobus/>
- Ekpenyong, T. E., & Borchers, R. L. (1980). Effect of Cooking on the Chemical Composition of Winged Beans (*Psophocarpus tetragonolobus*). *Journal of Food Science*, 45(6), 1559–1560. <https://doi.org/10.1111/j.1365-2621.1980.tb07562.x>
- Elsamani, M. O., Habbani, S. S., Babiker, E. E., & Ahmed, I. A. M. (2014). Biochemical, Microbial and Sensory Evaluation of White Soft Cheese Made from Cow and Lupin Milk. *LWT - Food Science and Technology*, 59(1), 553–559. <https://doi.org/10.1016/j.lwt.2014.04.027>
- Embaby, H. E.-S. (2010). Effect of Heat Treatments on Certain Antinutrients and in vitro Protein Digestibility of Peanut and Sesame Seeds. *Food Science and Technology Research*, 17(1), 31–38. <https://doi.org/10.3136/fstr.17.31>

- Embaby, H. E.-S. (2010). Effect of Soaking, Dehulling, And Cooking Methods on Certain Antinutrients And In Vitro Protein Digestibility of Bitter and Sweet Lupin Seeds. *Food Science and Biotechnology*, 19(4), 1055–1062. <https://doi.org/10.1007/S10068-010-0148-1>
- Erbas, M. (2010). The Effects of Different Debittering Methods on The Production of Lupin Bean Snack from Bitter Lupinus Albus L. Seeds. *Journal of Food Quality*, 33(6), 742–757. <https://doi.org/10.1111/J.1745-4557.2010.00347.x>
- Ertas, N., & Bilgiçli, N. (2012). Effect of Different Debittering Processes on Mineral and Phytic Acid Content of Lupin (*Lupinus Albus L.*) Seeds. *Journal of Food Science and Technology*, 51(11), 3348–3354. <https://doi.org/10.1007/S13197-012-0837-2>
- Esteves, E. A., Martino, H. S. D., Oliveira, F. C. E., Bressan, J., & Costa, N. M. B. (2010). Chemical composition of a soybean cultivar lacking lipoxygenases (LOX2 and LOX3). *Food Chemistry*, 122(1), 238–242. <https://doi.org/10.1016/j.foodchem.2010.02.069>
- Etemadi, F., Barker, A. V., Hashemi, M., Zandvakili, O. R., & Park, Y. (2018). Nutrient Accumulation in Faba Bean Varieties. *Communications in Soil Science and Plant Analysis*, 49(16), 2064–2073. <https://doi.org/10.1080/00103624.2018.1495729>
- Fabbri, A. D., & Crosby, G. A. (2016). A *review* of the impact of preparation and cooking on the nutritional quality of vegetables and legumes. *International Journal of Gastronomy and Food Science*, 3, 2–11. <https://doi.org/10.1016/j.ijgfs.2015.11.001>
- Fialkowski Revilla, M. K., Titchenall, A., Calabrese, A., Gibby, C., & Meinke, W. (2018). *Human Nutrition*. ISBN: 978-1-948027-01-4. <https://open.umn.edu/opentextbooks/textbooks/human-nutrition>
- Haddad, E., & Faed, P. (2014). *Vegetarianism and Veganism*. Reference Module in Biomedical Sciences. <https://doi.org/10.1016/b978-0-12-801238-3.00241-5>
- Hefni, M. E., Shalaby, M. T., & Witthöft, C. M. (2015). Folate content in faba beans (*Vicia faba*L.)-effects of cultivar, maturity stage, industrial processing, and bioprocessing. *Food Science & Nutrition*, 3(1), 65–73. <https://doi.org/10.1002/fsn3.192>
- Heuzé V., Thiollet H., Tran G., Bastianelli D., Lebas F., 2017. Peanut seeds. Feedipedia, a programme by INRA, CIRAD, AFZ and FAO. <https://www.feedipedia.org/node/55> . Last updated on January 25, 2017, 14:14, Retrieved April 1, 2020
- Heuzé V., Thiollet H., Tran G., Lebas F., 2017. Peanut forage. Feedipedia, a programme by INRA, CIRAD, AFZ and FAO. <https://feedipedia.org/node/695> . Last updated on January 23, 2017, 17:15, Retrieved April 1, 2020

- Heuzé V., Thiollet H., Tran G., Nozière P., Lessire M., Lebas F., 2020. White lupin (*Lupinus albus*) seeds. Feedipedia, a programme by INRA, CIRAD, AFZ and FAO. <https://www.feedipedia.org/node/279> . Last updated on February 5, 2020, 15:18, Retrieved April 1, 2020
- Heuzé V., Tran G., 2016. Soybean (general). Feedipedia, a programme by INRA, CIRAD, AFZ and FAO. <https://feedipedia.org/node/753> . Last updated on August 3, 2016, 10:37, Retrieved April 1, 2020
- Heuzé V., Tran G., Delagarde R., Lessire M., Lebas F., 2018. Faba bean (*Vicia faba*). Feedipedia, a programme by INRA, CIRAD, AFZ and FAO. <https://feedipedia.org/node/4926> . Last updated on July 17, 2018, 11:23, Retrieved April 1, 2020
- Hiwatashi, K., Shirakawa, H., Hori, K., Yoshiki, Y., Suzuki, N., Hokari, M., ... Takahashi, S. (2010). Reduction of Blood Pressure by Soybean Saponins, Renin Inhibitors from Soybean, in Spontaneously Hypertensive Rats. *Bioscience, Biotechnology, and Biochemistry*, 74(11), 2310–2312. <https://doi.org/10.1271/bbb.100328>
- Huang, G., Cai, W., & Xu, B. (2017). Improvement in beta-carotene, vitamin B<sub>2</sub>, GABA, free amino acids and isoflavones in yellow and black soybeans upon germination. *Lwt*, 75, 488–496. <https://doi.org/10.1016/j.lwt.2016.09.029>
- Huyghe, C. (1997). White lupin (*Lupinus albus* L.). *Field Crops Research*, 53(1-3), 147–160. [https://doi.org/10.1016/s0378-4290\(97\)00028-2](https://doi.org/10.1016/s0378-4290(97)00028-2)
- Hymowitz, T. (1970). On the domestication of the soybean. *Economic Botany*, 24(4), 408–421. <https://doi.org/10.1007/bf02860745>
- Interagency Taxonomic Information System. (2014). Taxonomic Hierarchy Fabaceae. Retrieved March 11, 2020, from [https://www.itis.gov/servlet/SingleRpt/SingleRpt?search\\_topic=TSN&search\\_value=500059#null](https://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=500059#null)
- Iqbal, A., Shah, F., Hamayun, M., Ahmad, A., Hussain, A., Waqas, M., ... Lee, I.-J. (2016). Allergens of *Arachis hypogaea* and the effect of processing on their detection by ELISA. *Food & Nutrition Research*, 60(1), 28945. <https://doi.org/10.3402/fnr.v60.28945>
- Ismail, A., Tiong, N. W., Tan, S. T., & Azlan, A. (2009). Antioxidant properties of selected non-leafy vegetables. *Nutrition & Food Science*, 39(2), 176 – 180. <https://doi.org/10.1108/00346650910943280>
- Jayanegara, A., Harahap, R. P., Rozi, R. F., & Nahrowi. (2018). Effects of lipid extraction on nutritive composition of winged bean (*Psophocarpus tetragonolobus*), rubber seed (*Hevea brasiliensis*), and tropical almond (*Terminalia catappa*). *Veterinary World*, 11(4), 446–451. <https://doi.org/10.14202/vetworld.2018.446-451>

- Kadam, S. S., Smithard, R. R., Eyre, M. D., & Armstrong, D. G. (1987). Effects of heat treatments of antinutritional factors and quality of proteins in winged bean. *Journal of the Science of Food and Agriculture*, 39(3), 267–275. <https://doi.org/10.1002/jsfa.2740390310>
- Kamalaja, T., Prashanthi, M., & Rajeswari, K. (2018). Evaluation of Antioxidant Activity and Bioactive Compounds on Domestic Cooking Method. *International Journal of Current Microbiology and Applied Sciences*, 7(08), 4090–4097. <https://doi.org/10.20546/ijcmas.2018.708.425>
- Kanetro, B. (2018). Amino acid profile of soybean (*Glicine max*) sprout protein for determining insulin stimulation amino acids. *International Food Research Journal* 2, 25(6), 2497–2502. [http://www.ifrj.upm.edu.my/25%20\(06\)%202018/\(34\).pdf](http://www.ifrj.upm.edu.my/25%20(06)%202018/(34).pdf)
- Karamać, M., Orak, H. H., Amarowicz, R., Orak, A., & Piekoszewski, W. (2018). Phenolic Contents and Antioxidant Capacities of Wild and Cultivated White Lupin (*Lupinus Albus L.*) Seeds. *Food Chemistry*, 258, 1–7. <https://doi.org/10.1016/J.Foodchem.2018.03.041>
- Ketnawa, S., & Ogawa, Y. (2019). Evaluation of protein digestibility of fermented soybeans and changes in biochemical characteristics of digested fractions. *Journal of Functional Foods*, 52, 640–647. <https://doi.org/10.1016/j.jff.2018.11.046>
- Khalili, R. M. A., Shafekh, S. E., Norhayati, A. H., Fatahudin, I. M., Rahimah, R., Norkamalia, H., & Azimah, A. N. (2013). Total Phenolic Content and In vitro Antioxidant Activity of Winged Bean (*Psophocarpus tetragonolobus*). *Pakistan Journal of Nutrition*, 12(5), 416–422. <https://doi.org/10.3923/pjn.2013.416.422>
- Khan, M. K., Karnpanit, W., Nasar-Abbas, S. M., Huma, Z.-E.-, & Jayasena, V. (2015). Phytochemical Composition and Bioactivities of Lupin: A Review. *International Journal of Food Science & Technology*, 50(9), 2004–2012. <https://doi.org/10.1111/ijfs.12796>
- Kim, G.-P., Lee, J., Ahn, K.-G., Hwang, Y.-S., Choi, Y., Chun, J., ... Choung, M.-G. (2014). Differential responses of B vitamins in black soybean seeds. *Food Chemistry*, 153, 101–108. <https://doi.org/10.1016/j.foodchem.2013.12.047>
- Koley, T. K., Maurya, A., Tripathi, A., Singh, B. K., Singh, M., Bhutia, T. L., ... Singh, B. (2018). Antioxidant potential of commonly consumed underutilized leguminous vegetables. *International Journal of Vegetable Science*, 25(4), 362–372. <https://doi.org/10.1080/19315260.2018.1519866>
- Kumar, V., Rani, A., Goyal, L., Vaishnav, J., Pratap, D., Dixit, A. K., & Billore, S. (2013). Assessment of Antioxidant Constituents and Anti-Oxidative Properties of Vegetable Soybean. *International Journal of Food Properties*, 17(3), 536–544. <https://doi.org/10.1080/10942912.2012.654559>

- Lepcha, P., Egan, A. N., Doyle, J. J., & Sathyanarayana, N. (2017). A Review on Current Status and Future Prospects of Winged Bean (*Psophocarpus tetragonolobus*) in Tropical Agriculture. *Plant Foods for Human Nutrition*, 72(3), 225–235. <https://doi.org/10.1007/s11130-017-0627-0>
- Li, Y., Qian, H., Sun, X., Cui, Y., Wang, H., Du, C., & Xia, X. (2014). The Effects of Germination on Chemical Composition of Peanut Seed. *Food Science and Technology Research*, 20(4), 883-889. <https://doi.org/10.3136/fstr.20.883>
- Lim, T. K. (2011). *Lupinus albus*. *Edible Medicinal and Non-Medicinal Plants*, 763–769. [https://doi.org/10.1007/978-94-007-1764-0\\_87](https://doi.org/10.1007/978-94-007-1764-0_87)
- Lim, T. K. (2011). *Psophocarpus tetragonolobus*. *Edible Medicinal and Non-Medicinal Plants*, 867–878. [https://doi.org/10.1007/978-94-007-1764-0\\_94](https://doi.org/10.1007/978-94-007-1764-0_94)
- Lim, T. K. (2011). *Vicia faba*. *Edible Medicinal And Non-Medicinal Plants*, 925–936. [https://doi.org/10.1007/978-94-007-1764-0\\_97](https://doi.org/10.1007/978-94-007-1764-0_97)
- Luo, S., Duan, H., Zou, Y., Qiu, R., & Wang, C. (2017). Quantification of Total Folate, Folate Species and Polyglutamyl Folate Distribution in Winged Beans (*Psophocarpus tetragonolobus* (L) DC) from Different Cultivars and Growth Stages by Ultra-High Performance Liquid Chromatography Tandem Mass Spectrometry. *Journal of Nutritional Science and Vitaminology*, 63(1), 69–80. <https://doi.org/10.3177/jnsv.63.69>
- Luo, Y. W., & Xie, W. H. (2013). Effect of different processing methods on certain antinutritional factors and protein digestibility in green and white faba bean (*Vicia faba*L.). *CyTA - Journal of Food*, 11(1), 43–49. <https://doi.org/10.1080/19476337.2012.681705>
- Luo, Y., & Xie, W. (2013). Effect of soaking and sprouting on iron and zinc availability in green and white faba bean (*Vicia faba* L.). *Journal of Food Science and Technology*, 51(12), 3970–3976. <https://doi.org/10.1007/s13197-012-0921-7>
- Mada, S., Garba, A., Mohammed, A., Muhammad, A., Olagunju, A., & Mohammed, H. (2012). Effects of Boiling and Roasting on Antinutrients and Proximate Composition of Local and Some Selected Improved Varieties of *Arachis hypogaea* L (Groundnut). *International Journal of Food Nutrition and Safety*, 1(1), 45–53. [https://www.researchgate.net/publication/235963420\\_Effects\\_of\\_Boiling\\_and\\_Roasting\\_on\\_Antinutrients\\_and\\_Proximate\\_Composition\\_of\\_Local\\_and\\_Some\\_Selected\\_Improved\\_Varieties\\_of\\_Arachis\\_hypogaea\\_L\\_Groundnut](https://www.researchgate.net/publication/235963420_Effects_of_Boiling_and_Roasting_on_Antinutrients_and_Proximate_Composition_of_Local_and_Some_Selected_Improved_Varieties_of_Arachis_hypogaea_L_Groundnut)
- Makeri, M. U., Mohamed, S. A., Karim, R., Ramakrishnan, Y., & Muhammad, K. (2017). Fractionation, physicochemical, and structural characterization of winged bean seed protein fractions with reference to soybean. *International Journal of Food Properties*, 1–17. <https://doi.org/10.1080/10942912.2017.1369101>



- Marsh, K. A., Munn, E. A., & Baines, S. K. (2012). Protein and vegetarian diets. *The Medical Journal of Australia*, 1(2), 7–10. <https://doi.org/10.5694/mjao11.11492>
- McGee, H. (2004). *McGee on food and cooking: an encyclopedia of kitchen science, history and culture*. London: Hodder & Stoughton. <http://wtf.tw/ref/mcgee.pdf>
- Mo, H., Kariluoto, S., Piironen, V., Zhu, Y., Sanders, M. G., Vincken, J.-P., ... Nout, M. R. (2013). Effect of soybean processing on content and bioaccessibility of folate, vitamin B12 and isoflavones in tofu and tempe. *Food Chemistry*, 141(3), 2418–2425. <https://doi.org/10.1016/j.foodchem.2013.05.017>
- Mohamed, I., El-Meihy, R., Ali, M., Chen, F., & Raleve, D. (2017). Interactive effects of biochar and micronutrients on faba bean growth, symbiotic performance, and soil properties. *Journal of Plant Nutrition and Soil Science*, 180(6), 729–738. <https://doi.org/10.1002/jpln.201700293>
- Mohanty, C. S., Pradhan, R. C., Singh, V., Singh, N., Pattanayak, R., Prakash, O., ... Rout, P. K. (2014). Physicochemical analysis of *Psophocarpus tetragonolobus* (L.) DC seeds with fatty acids and total lipids compositions. *Journal of Food Science and Technology*, 52, 3660–3670. <https://doi.org/10.1007/s13197-014-1436-1>
- Mohtar, W. A. A.-Q. I. W., Hamid, A. A., Abd-Aziz, S., Muhamad, S. K. S., & Saari, N. (2013). Preparation of bioactive peptides with high angiotensin converting enzyme inhibitory activity from winged bean [*Psophocarpus tetragonolobus* (L.) DC.] seed. *Journal of Food Science and Technology*, 51(12), 3658–3668. <https://doi.org/10.1007/s13197-012-0919-1>
- Mortuza, M. G., Hannan, M. A., & Tzen, J. T. (1970). Chemical composition and functional properties of *Vicia faba* L. from Bangladesh. *Bangladesh Journal of Botany*, 38(1), 93–97. <https://doi.org/10.3329/bjb.v38i1.5129>
- Mudgil, D., & Barak, S. (2013). Composition, properties and health benefits of indigestible carbohydrate polymers as dietary fiber: A review. *International Journal of Biological Macromolecules*, 61, 1–6. <https://doi.org/10.1016/j.ijbiomac.2013.06.044>
- Nile, S. H., & Park, S. W. (2013). Fatty Acid Composition and Antioxidant Activity of Groundnut (*Arachis hypogaea* L.) Products. *Food Science and Technology Research*, 19(6), 957–962. <https://doi.org/10.3136/fstr.19.957>
- Ningombam, R. D., Singh, P. K., & Salam, J. S. (2012). Proximate Composition and Nutritional Evaluation of Underutilized Legume *Psophocarpus tetragonolobus* (L.) DC. Grown in Manipur, Northeast India. *American Journal of Food Technology*, 7(8), 487–493. <https://doi.org/10.3923/ajft.2012.487.493>
- Noort, M. V. D. (2017). Lupin. *Sustainable Protein Sources*, 165–183. <https://doi.org/10.1016/b978-0-12-802778-3.00010-x>

- Nwosu, D. J., Olubiyi, M. R., Aladele, S. E., Apuyor, B., Okere, A. U., Lawal, A. I., ... Nwosu, E. C. (2019). Proximate and Mineral Composition of Selected Soybean Genotypes In Nigeria. *Journal of Plant Development*, 26, 67–76. <https://doi.org/10.33628/jpd.2019.26.1.67>
- Olaiya, C. O., Soetan, K. O., & Karigidi, K. O. (2018). Evaluation of In Vitro Antioxidant Capacities of Six Accessions of Winged Beans (*Psophocarpus tetragonolobus*). *EC Nutrition*, 13(8), 589–595. <https://www.semanticscholar.org/paper/Evaluation-of-In-Vitro-Antioxidant-Capacities-of-of-Olaiya-Soetan/8d196cc589d1965018a8c6b90bc1a171ca614bac>
- Olatidoye, O., & Sobowale, S. (2019). Effect of Cooking Methods on Anti-nutrient Content and Phenolic Acid Profiles of Groundnut Varieties Grown in Nigeria Antioxidant properties of nuts and pulses consumed in Nigeria View project. *International Research Journal of Food and Nutrition*, 1(1), 27–37. [https://www.researchgate.net/publication/334711029\\_Effect\\_of\\_Cooking\\_Methods\\_on\\_Anti-nutrient\\_Content\\_and\\_Phenolic\\_Acid\\_Profiles\\_of\\_Groundnut\\_Varieties\\_Grown\\_in\\_Nigeria](https://www.researchgate.net/publication/334711029_Effect_of_Cooking_Methods_on_Anti-nutrient_Content_and_Phenolic_Acid_Profiles_of_Groundnut_Varieties_Grown_in_Nigeria)
- Oomah, B. D., Luc, G., Leprelle, C., Drover, J. C. G., Harrison, J. E., & Olson, M. (2011). Phenolics, Phytic Acid, and Phytase in Canadian-Grown Low-Tannin Faba Bean (*Vicia faba* L.) Genotypes. *Journal of Agricultural and Food Chemistry*, 59(8), 3763–3771. <https://doi.org/10.1021/jf200338b>
- Prusinski, J. (2017). White Lupin (*Lupinus Albus* L.) – Nutritional and Health Values in Human Nutrition – A Review. *Czech Journal of Food Sciences*, 35(No. 2), 95–105. <https://doi.org/10.17221/114/2016-Cjfs>
- Ranilla, L. G., Genovese, M. I., & Lajolo, F. M. (2009). Isoflavones and antioxidant capacity of Peruvian and Brazilian lupin cultivars. *Journal of Food Composition and Analysis*, 22(5), 397–404. <https://doi.org/10.1016/j.jfca.2008.06.011>
- Revilla, I. (2015). Impact of Thermal Processing on Faba Bean (*Vicia faba*) Composition. *Processing and Impact on Active Components in Food*, 337–343. <https://doi.org/10.1016/b978-0-12-404699-3.00040-8>
- Robinson, G. H. J., Balk, J., & Domoney, C. (2019). Improving pulse crops as a source of protein, starch and micronutrients. *Nutrition Bulletin*, 44(3), 202–215. <https://doi.org/10.1111/nbu.12399>
- Rockland, L. B., Zaragosa, E. M., & Oracca-Tetteh, R. (1979). Quick-Cooking Winged Beans (*Psophocarpus tetragonolobus*). *Journal of Food Science*, 44(4), 1004–1007. <https://doi.org/10.1111/j.1365-2621.1979.tb03432.x>
- Ruberte, R. M., & Martin, F. W. (1978). Cooking of winged bean seed. *The Journal of Agriculture of the University of Puerto Rico*, 62(4), 321–329. <https://agris.fao.org/agris-search/search.do?recordID=US201302466600>

- Sambudi, H., & Buckle, K. A. (1991). Characteristics of winged bean (*Psophocarpus tetragonolobus*) seeds during soaking and boiling. *Journal of the Science of Food and Agriculture*, 57(4), 585–595. <https://doi.org/10.1002/jsfa.2740570411>
- Sathe, S. K., & Salunkhe, D. K. (1981). Investigations on Winged Bean [*Psophocarpus tetragonolobus* (L.) DC] Proteins and Antinutritional Factors. *Journal of Food Science*, 46(5), 1389–1393. <https://doi.org/10.1111/j.1365-2621.1981.tb04181.x>
- Schulmerich, M. V., Gelber, M. K., Azam, H. M., Harrison, S. K., Mckinney, J., Thompson, D., ... Bhargava, R. (2013). Amino Acid Quantification in Bulk Soybeans by Transmission Raman Spectroscopy. *Analytical Chemistry*, 85(23), 11376–11381. <https://doi.org/10.1021/ac402284b>
- Schumacher, H., Paulsen, H. M., Gau, A. E., Link, W., Jürgens, H. U., Sass, O., & Dieterich, R. (2011). Seed protein amino acid composition of important local grain legumes *Lupinus angustifolius* L., *Lupinus luteus* L., *Pisum sativum* L. and *Vicia faba* L. *Plant Breeding*, 130(2), 156–164. <https://doi.org/10.1111/j.1439-0523.2010.01832.x>
- Sebei, K., Gnouma, A., Herchi, W., Sakouhi, F., & Boukhchina, S. (2013). Lipids, proteins, phenolic composition, antioxidant and antibacterial activities of seeds of peanuts (*Arachis hypogaea* L) cultivated in Tunisia. *Biological Research*, 46(3), 257–263. <https://doi.org/10.4067/s0716-97602013000300006>
- Sedláková, K., Straková, E., Suchý, P., Krejcarová, J., & Herzig, I. (2016). Lupin as a perspective protein plant for animal and human nutrition – a review. *Acta Veterinaria Brno*, 85(2), 165–175. <https://doi.org/10.2754/avb201685020165>
- Settaluri, V. S., Kandala, C. V. K., Puppala, N., & Sundaram, J. (2012). Peanuts and Their Nutritional Aspects—A Review. *Food and Nutrition Sciences*, 03(12), 1644–1650. <https://doi.org/10.4236/fns.2012.312215>
- Shokunbi, O. S., Fayomi, E. T., Sonuga, O. S., & Tayo, G. O. (2012). Nutrient composition of five varieties of commonly consumed Nigerian groundnut (*Arachis hypogaea* L.). *Grasas y Aceites*, 63(1), 14–18. <https://doi.org/10.3989/gya.056611>
- Siah, S., Wood, J. A., Agboola, S., Konczak, I., & Blanchard, C. L. (2014). Effects of soaking, boiling and autoclaving on the phenolic contents and antioxidant activities of faba beans (*Vicia faba* L.) differing in seed coat colours. *Food Chemistry*, 142, 461–468. <https://doi.org/10.1016/j.foodchem.2013.07.068>
- Siger, A., Czubinski, J., Kachlicki, P., Dwiecki, K., Lampart-Szczapa, E., & Nogala-Kalucka, M. (2012). Antioxidant Activity and Phenolic Content in Three Lupin Species. *Journal of Food Composition and Analysis*, 25(2), 190–197. <https://doi.org/10.1016/j.jfca.2011.10.002>
- Singh, A., Bharadwaj, R., & Singh, I. (2014). Assessment of Nutritional Quality of Developed Faba Bean (*Vicia faba* L.) Lines. *Journal of AgriSearch*, 1(2), 96–

101.  
[https://www.researchgate.net/publication/283538549\\_Assessment\\_of\\_Nutritional\\_Quality\\_of\\_Developed\\_Faba\\_Bean\\_Vicia\\_faba\\_L\\_Lines](https://www.researchgate.net/publication/283538549_Assessment_of_Nutritional_Quality_of_Developed_Faba_Bean_Vicia_faba_L_Lines)
- Singh, M., Dubey, R., Koley, T., Maurya, A., Singh, P., & Singh, B. (2019). Valorization of winged bean (*Psophocarpus tetragonolobus* (L) DC) by evaluation of its antioxidant activity through chemometric analysis. *South African Journal of Botany*, 121, 114–120.  
<https://doi.org/10.1016/j.sajb.2018.10.026>
- Stabler, S. P. (2013). Clinical practice. Vitamin B12 deficiency. *The New England Journal of Medicine*, 368(2), 149–160. <https://doi.org/10.1056/NEJMcp1113996>
- Stalker, H. (1997). Peanut (*Arachis hypogaea* L.). *Field Crops Research*, 53(1-3), 205–217. [https://doi.org/10.1016/s0378-4290\(97\)00032-4](https://doi.org/10.1016/s0378-4290(97)00032-4)
- Stocco, B., Toledo, K. A., Fumagalli, H. F., Bianchini, F. J., Fortes, V. S., Fonseca, M. J. V., & Toloï, M. R. T. (2015). Biotransformed Soybean Extract Induces Cell Death of Estrogen-Dependent Breast Cancer Cells by Modulation of Apoptotic Proteins. *Nutrition and Cancer*, 67(4), 612–619.  
<https://doi.org/10.1080/01635581.2015.1015744>
- Thambiraj, S. R., Reddy, N., Phillips, M., & Koyyalamudi, S. R. (2019). Biological Activities and Characterization of Polysaccharides from The Three Australian Sweet Lupins. *International Journal of Food Properties*, 22(1), 522–535.  
<https://doi.org/10.1080/10942912.2019.1588298>
- Thorp, A. A., Sinn, N., Buckley, J. D., Coates, A. M., & Howe, P. R. C. (2009). Soya isoflavone supplementation enhances spatial working memory in men. *British Journal of Nutrition*, 102(9), 1348–1354.  
<https://doi.org/10.1017/s0007114509990201>
- Tobin-West, M. D., & Baraka, R. E. (2018). Effects of Methods of Processing Groundnut (*Arachis hypogaea* L.) on the Nutritional Composition and Storage Life of the Seed. *International Journal of Agriculture and Earth Science*, 4(6), 43–49.  
[https://www.researchgate.net/publication/339442247\\_Effects\\_of\\_Methods\\_of\\_Processing\\_Groundnut\\_Arachis\\_hypogaea\\_L\\_on\\_the\\_Nutritional\\_Composition\\_and\\_Storage\\_Life\\_of\\_the\\_Seed](https://www.researchgate.net/publication/339442247_Effects_of_Methods_of_Processing_Groundnut_Arachis_hypogaea_L_on_the_Nutritional_Composition_and_Storage_Life_of_the_Seed)
- Toomer, O. T. (2017). Nutritional chemistry of the peanut (*Arachis hypogaea*). *Critical Reviews in Food Science and Nutrition*, 58(17), 3042–3053.  
<https://doi.org/10.1080/10408398.2017.1339015>
- Turco, I., Ferretti, G., & Bacchetti, T. (2016). Review of the health benefits of Faba bean (*Vicia faba* L.) polyphenols. *Journal of Food & Nutrition Research*, 55(4), 283–293.  
[https://www.researchgate.net/publication/311931044\\_Review\\_of\\_the\\_health\\_benefits\\_of\\_Faba\\_bean\\_Vicia\\_faba\\_L\\_polyphenols](https://www.researchgate.net/publication/311931044_Review_of_the_health_benefits_of_Faba_bean_Vicia_faba_L_polyphenols)

- Turfan, N., Kurnaz, A., Karataşlı, M., Özer, T., & Turhan, Ş. (2019). Nutrition content assessment of *Arachis hypogaea* L. samples grown in the Mediterranean region of Turkey. *Turkish Journal of Agriculture - Food Science and Technology*, 7(6), 892. <https://doi.org/10.24925/turjaf.v7i6.892-898.2493>
- Tvrzicka, E., Kremmyda, L.-S., Stankova, B., & Zak, A. (2011). Fatty Acids As Biocompounds: Their Role In Human Metabolism, Health And Disease - A Review. Part 1: Classification, Dietary Sources And Biological Functions. *Biomedical Papers*, 155(2), 117–130. <https://doi.org/10.5507/bp.2011.038>
- USDA PLANTS. (n.d.). Classification for Kingdom Plantae Down to Species *Glycine max* (L.) Merr. Retrieved March 11, 2020, from <https://plants.usda.gov/java/ClassificationServlet?source=display&classid=GLMA4>
- USDA PLANTS. (n.d.). Classification for Kingdom Plantae Down to Species *Lupinus albus* L. white lupine. Retrieved March 12, 2020, from <https://plants.usda.gov/core/profile?symbol=LUAL22>
- USDA PLANTS. (n.d.). Classification for Kingdom Plantae Down to Species *Psophocarpus tetragonolobus* (L.) DC. winged bean. Retrieved March 12, 2020, from <https://plants.usda.gov/core/profile?symbol=PSTE10>
- USDA PLANTS. (n.d.). Classification for Kingdom Plantae Down to Species *Vicia faba* L. fava bean. Retrieved March 12, 2020, from <https://plants.sc.egov.usda.gov/core/profile?symbol=VIFA>
- USDA PLANTS. (n.d.). Classification for Kingdom Plantae Down to Species *Arachis hypogaea* L. peanut. Retrieved March 12, 2020, from <https://plants.sc.egov.usda.gov/core/profile?symbol=ARHY>
- USDA. (2019, April 1). FoodData Central Search Results Broadbeans (fava beans), mature seeds, raw. Retrieved March 12, 2020, from <https://fdc.nal.usda.gov/fdc-app.html#/food-details/175205/nutrients>
- USDA. (2019, April 1). FoodData Central Search Results Lupins, mature seeds, raw. Retrieved March 12, 2020, from <https://fdc.nal.usda.gov/fdc-app.html#/food-details/172423/nutrients>
- USDA. (2019, April 1). FoodData Central Search Results Peanuts, all types, raw. Retrieved March 12, 2020, from <https://fdc.nal.usda.gov/fdc-app.html#/food-details/172430/nutrients>
- USDA. (2019, April 1). FoodData Central Search Results Soybeans, mature seeds, raw. Retrieved March 12, 2020, from <https://fdc.nal.usda.gov/fdc-app.html#/food-details/174270/nutrients>
- USDA. (2019, April 1). FoodData Central Search Results Winged beans, mature seeds, raw. Retrieved March 12, 2020, from <https://fdc.nal.usda.gov/fdc-app.html#/food-details/174283/nutrients>

- Venkidasamy, B., Selvaraj, D., Nile, A. S., Ramalingam, S., Kai, G., & Nile, S. H. (2019). Indian pulses: A review on nutritional, functional and biochemical properties with future perspectives. *Trends in Food Science & Technology*, 88, 228–242. <https://doi.org/10.1016/j.tifs.2019.03.012>
- Vioque, J., Alaiz, M., & Girón-Calle, J. (2012). Nutritional and functional properties of *Vicia faba* protein isolates and related fractions. *Food Chemistry*, 132(1), 67–72. <https://doi.org/10.1016/j.foodchem.2011.10.033>
- Wolosiak, R., Worobiej, E., Piecyk, M., Druzynska, B., Nowak, D., & Lewicki, P. P. (2009). Activities of amine and phenolic antioxidants and their changes in broad beans (*Vicia faba*) after freezing and steam cooking. *International Journal of Food Science & Technology*, 45(1), 29–37. <https://doi.org/10.1111/j.1365-2621.2009.02099.x>
- Woumbo, C. Y., Kuate, D., & Womeni, H. M. (2017). Cooking methods affect phytochemical composition and anti-obesity potential of soybean (*Glycine max*) seeds in Wistar rats. *Heliyon*, 3(8). <https://doi.org/10.1016/j.heliyon.2017.e00382>
- Xu, X. P., Liu, H., Tian, L., Dong, X. B., Shen, S. H., & Qu, L. Q. (2015). Integrated and comparative proteomics of high-oil and high-protein soybean seeds. *Food Chemistry*, 172, 105–116. <https://doi.org/10.1016/j.foodchem.2014.09.035>
- Young, V. R., Scrimshaw, N. S., Torun, B., & Viteri, F. (1979). Soybean protein in human nutrition: An overview. *Journal of the American Oil Chemists Society*, 56(3), 110–120. <https://doi.org/10.1007/bf02671432>
- Zhong, Y., Wang, Z., & Zhao, Y. (2015). Impact of Radio Frequency, Microwaving, and High Hydrostatic Pressure at Elevated Temperature on the Nutritional and Antinutritional Components in Black Soybeans. *Journal of Food Science*, 80(12). <https://doi.org/10.1111/1750-3841.13131>
- Žilić, S., Akıllıoğlu, H. G., Serpen, A., Perić, V., & Gökmen, V. (2013). Comparisons of phenolic compounds, isoflavones, antioxidant capacity and oxidative enzymes in yellow and black soybeans seed coat and dehulled bean. *European Food Research and Technology*, 237(3), 409–418. <https://doi.org/10.1007/s00217-013-2005-y>